ANAESTHESIA AND TRANSMURAL DISPERSION OF REPOLARISATION IN CHILDREN

Simon D Whyte MBBS, Peter D Booker MD, MBBS and David G Buckley CCSO.
Department of Anaesthesia, Royal Liverpool Children's Hospital, Eaton Road, Liverpool, L12 2AP,
United Kingdom.

INTRODUCTION: Prolongation of the QT interval is associated with torsades de pointes (TdP), classically in children or young adults with long QT syndromes (LQTS). Any overt or covert predisposition to TdP may be unmasked by perioperative stress. Which anaesthetic drugs are safest to use in these patients? Susceptibility to TdP arises from increased transmural dispersion of repolarisation (TDR) across the myocardial wall, rather than QT interval prolongation *per se* [1]. Agents that don't increase TDR would be useful in minimising any increased perioperative risk of TdP. Several anaesthetic drugs prolong the QT interval, but their effect on TDR is unknown. TDR can be measured on the ECG as the time interval between the peak and end of the T wave (Tp-e) [2]. We have investigated the effects of propofol and sevoflurane on the QT and Tp-e intervals in children.

METHODS: 50 ASA I-II unpremedicated children, aged 1-14, received propofol (group P – target plasma concentration 3 µg/ml) or sevoflurane (group S - 8% for induction followed by 3%) for 15 min prior to elective surgery. Both groups breathed 40% oxygen in air. We recorded a 12 lead ECG on each patient pre-induction and 15 minutes post-induction. RR, QT and Tp-e intervals were measured by investigators blinded to the anaesthetic received and to the pre-induction/post-induction status of the recording. Corrected QT interval (QTc) was calculated as $QT/\sqrt{RR}$. Paired t-tests were used to compare QTc and Tp-e before and after propofol and sevoflurane anaesthesia; unpaired t-tests were used for between-patient comparisons of QTc and Tp-e after propofol or sevoflurane anaesthesia.

RESULTS: Age and weight were similar between groups P (n = 22) and S (n = 28). Interobserver variation was least in leads I, II and V5. Sevoflurane significantly prolonged the pre-induction QTc; propofol did so to a much lesser extent (see table). Neither agent had any significant effect on the pre-induction Tp-e.

ECG Lead	I	II	V5
<u>Propofol</u>			
QTc pre	418	423	436
QTc post	430	432	442
<i>p</i>	0.012	0.06	NS
Tp-e pre	71	75	80
Tp-e post	74	74	83
<i>p</i>	NS	NS	NS
<u>Sevoflurane</u>			
QTc pre	406	413	423
QTc post	445	449	456
<i>p</i>	<0.0001	<0.0001	<0.0001
Tp-e pre	71	71	77
Tp-e post	66	70	80
<i>p</i>	NS	NS	NS

DISCUSSION: Sevoflurane increases the duration of myocardial repolarisation in children to a greater extent than propofol. However, as the dispersion of repolarisation appears unaffected, neither agent is likely to increase the risk of torsades de pointes.

REFERENCES:

- [1] *J Cardiovasc Electrophysiol* 1999; **10**: 154-64
- [2] *Circulation* 1998; **98**: 1928-36

QTc = corrected QT interval (msec)
Tp-e = Tpeak-end interval (msec)
Pre = pre-induction ECG
Post = post-induction ECG
P = p value for paired t test