risk of anaphylaxis during anaesthesia. Abbreviated text. *Ann Fr Anesth Reanim* 2002; 21 Suppl 1: 7–23.

- 7 Joint Task Force on Practice Parameters; American Academy of Allergy, Asthma and Immunology; American College of Allergy, Asthma and Immunology; Joint Council of Allergy, Asthma and Immunology. The diagnosis and management of anaphylaxis: an updated practice parameter. *J Allergy Clin Immunol* 2005; 115(3 Suppl 2): S483–523.

Reply:

I thank Drs. Dewachter and Mouton-Faivre for their thoughtful comments on our case report. I believe, however, that certain points require comment and clarification.

First, they state that a reaction to rocuronium cannot be ruled out because of the brief time between the event and the subsequent skin testing. We agree that published recommendations¹ suggest a delay of four to six weeks from the event to allow regeneration of reactive IgE and mast cells. Due to logistical issues, this was not feasible in our situation. The patient did, however, react to succinylcholine, which, if their reasoning is true, would require antibodies to two separate neuromuscular blocking drugs (NMBDs) instead of one. This is certainly possible, and we subsequently referred the patient for repeat skin testing. Fentanyl allergy, however, is exceedingly rare, with only case reports documenting occurrences. Our hypothesis of causation by fentanyl rests partially on the fact that a false positive skin test to fentanyl should be considerably less likely than a NMBD. The work by Dhonneur et al.² highlights the hazards of interpreting a positive skin test for rocuronium or vecuronium.

The results of the patient's repeat skin testing were quite unexpected: her only reaction was to the positive control (histamine), with no reaction to any of the NMBDs or fentanyl. The patient's allergist believes that these results represent a false negative, and that she has a low level of antibodies to one of the drugs tested. This situation has been encountered previously with Hymenoptera venom allergy and is a vexing problem in the allergy literature.³ The best advice we can offer the patient at this point is to avoid fentanyl, succinylcholine, and rocuronium. If she were to require surgery with paralysis in the future, cisatracurium may be the wisest choice. One cannot, however, be entirely certain, as false-negative results have also been reported for NMBDs.⁴

Finally, I would like to comment on the assertion that we distorted the conclusions of the allergist. Due to the ambiguity of the original test results, the allergist drew a rational conclusion: avoid all paralytic agents. We

have made the patient aware of the uncertainty from the skin tests. Our position, however, is that this could pose an unnecessary hazard to the patient in case of future surgery. General anesthesia with airway management, particularly in an emergency, could be much more hazardous without the use of a NMBD. The patient's lack of response (twice) to cisatracurium is somewhat reassuring, as she did not receive this drug prior to her initial reaction. I apologize for the lack of clarity on this point in the case report. A wiser phrasing might have been: "It is likely that the patient may receive selected NMBDs, such as cisatracurium, in the future if circumstances warranted."

Kenneth C. Cummings III MD
The Cleveland Clinic, Cleveland, USA
E-mail: cummink2@ccf.org

References

- 1 Mertes PM, Laxenaire MC, Lienhard A, et al. Reducing the risk of anaphylaxis during anesthesia: guidelines for clinical practice. *J Invest Allergol Clin Immunol* 2005; 15: 91–101.
- 2 Dhonneur G, Combes X, Chassard D, Merle JC. Skin sensitivity to rocuronium and vecuronium: a randomized controlled prick-testing study in healthy volunteers. *Anesth Analg* 2004; 98: 986–9.
- 3 Kontou-Fili K. Patients with negative skin tests. *Curr Opin Allergy Clin Immunol* 2002; 2: 353–7.
- 4 Fisher MM, Mercfield D, Baldo B. Failure to prevent an anaphylactic reaction to a second neuromuscular blocking drug during anaesthesia. *Br J Anaesth* 1999; 82: 770–3.

Evaluating effectiveness of medical emergency teams

To the Editor:

Brindley et al.¹ are to be congratulated on their balanced appraisal of the MERIT study report.² Unlike other commentators^{3,4} who have used its results to question the value of Medical Emergency Teams (MET), they rightly draw attention to the limitations of the study, which make it difficult to draw firm conclusions from its results. MERIT failed to find a clear advantage to the introduction of MET in a cluster randomized prospective trial, but the duration of the study period was short (only six months), and there was possibly cross-contamination between the study groups. Such teams are complex, introduction is labour-intensive and may be difficult, and after only