exchange during routine fiberoptic bronchoscopy. *Chest* 1984; 86: 184–8.

- 9 Gal TJ, Suratt PM. Resistance to breathing in healthy subjects after endotracheal intubation under topical anesthesia. *Anesth Analg* 1980; 59: 27–4.
- 10 Rodenstein DO, Stanescu DC, Francis C. Demonstration of failure of body plethysmography in airway obstruction. *J Appl Physiol* 1982; 52: 949–54.

REPLY

Dr. T. Gal, who is well versed in respiratory muscle and small airway physiology, raises very specific points concerning the mechanism of reduced FRC during anaesthesia. Before answering, two considerations must be clarified:

- In a review of the literature designed for clinicians, as opposed to respiratory physiologists, the author often has to make hard decisions about what to include or exclude and how to present these data in the general context. Thus, some readers may disagree.
- Secondly, there are many instances, as will be seen below, when similar studies have yielded very different results. The difficulty then is to reconcile these different results.

The first point raised concerns the effects of d-tubocurarine on FRC. DeTroyer's¹ work was quoted only to explain the role of inspiratory muscle tone in determining FRC and not to discuss the effects of curarization as such on FRC. It is true that curarization reduces FRC only if the subject is seated. I regret this misunderstanding. The reader is referred to a lucid editorial which reconciled the differences between DeTroyer's data and that which Dr. Gal quotes above.² The reconciliation allows the conclusion that the change in FRC is primarily due to loss of inspiratory muscle tone.

Next is the question of gas trapping. Rehder et al. reported a clear-cut relationship between closing volume and the degree of reduced FRC.³ Later work, however, did not confirm that: an example of differing results we sometimes read.

I have looked very carefully at the data reported by the Mayo Clinic group⁴ and found no correlation between either the magnitude of change or the intraoperative FRC and closing volume. For that reason I was very careful in stating that gas trapping may be an additional contributing factor. Another example of the difficulties encountered is in reconciling Hickey's data⁵ (FRC is reduced more in patients with COPD) and Gunnarsson's⁶ work (no dimensional changes in CT scan in similar patients). The difference is, logically, due to trapped gas. It is in accepting this explanation that the bits of information can be placed in the general framework. I see Dr. Gal's points and I can but reiterate that gas trapping can only be an additional mechanism.

Richard W.M. Wahba MB BCH SCFRCP
Montréal, Que.

REFERENCES

- 1 DeTroyer A, Bastenier J, Delhez L. Function of respiratory muscles during partial curarization in humans. *J Appl Physiol* 1980; 49: 1049–56.
- 2 DeTroyer A, Martin JG. Respiratory muscle tone and the control of functional residual capacity. *Chest* 1983; 84: 3–4.
- 3 Rehder K, Mallow JE, Fibush EE, Krabill DR, Sessler AD. Effects of isoflurane anesthesia and muscle paralysis on respiratory mechanics in normal man. *Anesthesiology* 1974; 44: 477–85.
- 4 Juno P, Marsh M, Knopp TJ, Rehder K. Closing capacity in awake and anesthetized-paralyzed man. *J Appl Physiol* 1978; 44: 238–44.
- 5 Hickey RF, Visick WD, Fairley HB, Fourcade HE. Effects of halothane anesthesia on functional residual capacity and alveolar-arterial oxygen tension difference. *Anesthesiology* 1973; 38: 20–4.
- 6 Gunnarsson L, Tokics L, Lundquist H et al. Chronic obstructive pulmonary disease and anaesthesia – formation of atelectasis and gas exchange impairment. *Acta Anaesthesiol Scan* (In Press).

Uterine relaxation can be life saving

To the Editor:

Uterine relaxation may be required in complicated obstetrical situations. Classically amyl nitrite, diethyl ether and halogenated hydrocarbons have been used. Recently, betamimetic drugs are being used.^{1–3} There are some reports of manual extraction of retained placentae using intravenous nitroglycerin to relax the uterus,^{4,5} but none, to our knowledge, of its use at Caesarean delivery to induce rapid uterine relaxation as a life-saving manoeuvre.

The patient presented at term, with a double footling breech in active labour. A Caesarean section was performed under epidural anaesthesia. At delivery, the baby's head was trapped in the hypertonic upper segment. While we did not document the duration of these events it became obvious that the baby was in jeopardy. The anaesthetist gave 1000 µg (1 mg) of nitroglycerin intravenously. It was only then that the uterus became sufficiently relaxed to allow delivery of the head. The patient remained asymptomatic although her blood pressure did decrease to 70/30 but this responded to 36 mg of ephedrine in divided doses.

The Apgar scores were five and nine at one and five minutes respectively. The infant breathed room air by two minutes of age and weighed 3090 g. Cord blood gas analysis was not measured.

This patient's obstetrical problem was life-threatening and its management unique. Rapid uterine relaxation was essential. All the methods used to induce uterine relaxation have potential problems – e.g., delay in onset of action, increased bleeding, risks of emergency anaesthetic induction and subsequent deep general anaesthesia.

It is our impression that intravenous nitroglycerin relaxes the uterus quickly, effectively and safely. Further