9. Hayward JR et al. (1981) Central cavernous hemangioma of the mandible :report of four cases . J Oral Surg. ;39:526.
10. Halazonites JA ,Kountourise J, Halazonites NA.(1982) Arterio-venous aneurysms of the mandible :report of a case. Oral Surg Oral Med Oral Pathol. ;53: 454.
11. James W Smith, Sherrell J Aston, eds. Grabb and Smith's (1991) Plastic Surgery Boston / Toronto /London Little Brown and Company. :805-18
12. Hayton -Williams DS. Arterio -venous aneurysm simulating a mandibular cyst. Oral Surg. 195: 8:21
13. Cook TJ,Zbar MI.Arterio-venous aneurysms of the mandible. Oral Surg. (1962) ; 15: 442.
14. Kelly DE,Terry BC, Small EW. (1977) Arterio -venous malformation on the mandible:report of a case . J Oral Surg. ; 35 :387.
15. Whear DM.(1991) Condylar hemangioma; a case report and review of the literature. Br J Oral and Maxillofacial Surg. ; 29 : 44.
16. Sadowsky D.Rossnberg Rd, Kanfman j, Levin BC ,Friedman JM. 1981 Central hemangioma of the mandible. Oral Surg Oral Med Oral Pathol; 52 : 447.
17. Frame JW ,Putman G,Eake MJ,Rolfe EB . (1987) Therapeutic embolization of vascular lesions in the Maxillofacial region. Br J Oral and Maxillofacial Surg. ; 25: 181.
18. Yeoman CM. The management of hemangioma involving facial, mandibular and pharyngeal structures. (1987) Br J Oral and Maxillofacial Surg. ; 25 : 181.
19. Thompson JN. Fierstein BB,Kohut RI. Embolization techniques in vascular tumors of the head and neck (1979). Head and Neck Surg. ; 2 : 25.
20. Guilbert-Trainer F, Piton J,Riche MC,Merland II (1982). Vascular malformations of the mandible (intraosseous hemangiomas). The importance of preoperative embolization : a Study of nine cases. Europ J Radiol. ; 2:257.

CORRIGENDUM

Non-Sporing Anaerobic Bacteria in ENT Infection – B. S. Nagoba et al Vol No. 3 1994. Please read the following few lines in its material and method section in continuity with the already published matter as its last few lines.

Sheep blood agar and plain sheep blood agar. In the area of primary streaking a disc of gentamicin (10 µg) was placed and in the area of secondary streaking a disc of metronidazole (5 µg) was placed for presumptive identification of anaerobic bacteria. The plates were incubated in Dynox anaerobic jar, at 37°C for 48 Hrs. or more. After incubation the bacterial colony resistant to gentamicin and sensitive to metronidazole was presumptively identified as anaerobe and further studies in detail for identification using conventional anaerobic techniques. (Nagoba et. al 1991, Sutter et. al. 1980).