Recent Advances in the Chemistry of Gold(I) Complexes with C-, N- and S-Donor Ligands

Part I: Alkynyl, Amino, Imino and Nitrido Derivatives

José Vicente, M Teresa Chicote, M Dolores Abrisqueta, Pablo González-Herrero, and Rita Guerrero*

Grupo de Química Organometálica, Departamento de Química Inorgánica, Facultad de Química, Universidad de Murcia, Apartado 4021, Murcia, 30071 Spain

In this paper, and in Part II to be published later, we give an account of the chemistry of gold(I) complexes with C-, N-, and S-donor ligands. In this first part, the synthesis of gold(I) complexes with alkynyl and N-donor ligands is reported.

INTRODUCTION

Gold(I) complexes are interesting from both basic and applied points of view (1-5). Thus, many gold(I) complexes show Au...Au interactions that are weaker than normal covalent bonds but stronger than van der Waals forces. These interactions are called *aurophilic* because it was in gold complexes that they were first found (6, 7). Such interactions determine the supramolecular structure of many gold(I) complexes as well as the formation of rare hypercoordinate complexes (8-12). Much of the present basic research devoted to gold(I) complexes is aimed at the synthesis and structural characterization of new complexes (particularly hypercoordinate) in order to better understand the *aurophilic* interaction.

Because gold(I) has a high preference for S- and P-donor ligands, *ie* soft ligands, to give linear two-coordinate complexes, the synthesis of complexes with N- and O-donor ligands or with coordination numbers three and four (13) are other areas of fundamental importance.

The search for gold(I) complexes with potential applications is also an important research area (5). In this respect, the synthesis of complexes with S-donor ligands is probably one of the more active areas because it is well known that such complexes find important applications for the treatment of rheumatoid arthritis (14-16).

However, although other potential medical applications of these complexes have been reported (16-22), as yet no gold compound has entered into clinical trials.

Some rod-like isocyanide or alkynyl-isocyanidegold(I) complexes have mesomorphic properties (23-28). In general, the usual linear coordination of gold(I) makes its complexes potentially useful metal-based liquid crystalline materials (*metallomesogens*) (29). In some cases, a relationship between the *aurophilic* interaction and potentially interesting properties has been suggested, as in the case of the solvoluminescence of some trinuclear complexes (30, 31) and luminescence of some alkynyl or alkynyl-isocyanidegold(I) complexes (32, 33). Some alkynylgold(I) complexes show non-linear optical properties (34, 35) and isonitrilegold(I) nitrates are effective precursors for chemical deposition of gold on iron oxide to give catalysts which efficiently oxidize carbon monoxide in air at low temperatures (36).

In the first part of this account, we report our recent results in the synthesis of alkynylgold(I) complexes containing isocyanide, carbene, halide, amine, phosphine or aryl ligands and also ammine and other N-donor derivatives of gold(I). Only a few of these complexes display *aurophilic* interactions but those that do so are of unprecedented nature. In the second part of this account we will report new gold(I) complexes with sulfur containing ligands.

ALKYNYLGOLD(I) COMPLEXES

As stated above, the rapidly growing interest in alkynyl metal complexes is based mainly on their non-linear optical, liquid-crystalline, photophysical or photochemical properties. Most alkynyl gold(I) complexes are neutral, and have the formulae $[\text{Au}(\text{C}\equiv\text{CR})]_n$, $[\text{Au}(\text{C}\equiv\text{CR})(\text{L})]$ or $[\text{Au}_2(\mu\text{-C}\equiv\text{C})\text{L}_2]$ with L = phosphine (37-44), although a few complexes with isocyanide (25, 28, 45-47), amine (48, 49), or ylide ligands (50) have also been reported.

We have prepared a wide variety of anionic and neutral alkynyl gold(I) complexes by various synthetic routes. The use of the 'acac method' (acac = acetylacetonate) based on the general reactions (1) and (2) allowed us to prepare gold(I) complexes with ylide, thiolato, phosphide, bis(diphenylphosphino) methanide or alkyl ligands (12, 51-69):

Thus, by reacting $[\text{Au}(\text{acac})(\text{PPh}_3)]$ with the phosphonium salt $[\text{Ph}_3\text{PCH}_2\text{C}(\text{O})\text{Me}]\text{ClO}_4$, the ylide complex $[\text{Au}\{\text{CH}(\text{PPh}_3)\text{C}(\text{O})\text{Me}\}(\text{PPh}_3)]$ can be obtained. The extension of this method to prepare new alkynyl gold(I) complexes, by reacting $\text{PPN}[\text{Au}(\text{acac})_2]$ ($\text{PPN} = \text{N}(\text{PPh}_3)_2$) or $\text{PPN}[\text{Au}(\text{acac})(\text{R})]$ or $[\text{Au}(\text{acac})(\text{PR}_3)]$ with terminal alkynes (see Box 1), provides further examples of its versatility (68, 70, 71). The use of acetylene allowed the synthesis of the first family of ethynylgold(I) complexes (68). Only one well characterized ethynyl gold(I) complex had previously been reported, *ie* $[\text{Au}(\text{C}\equiv\text{CH})(\text{PPr}^i_3)]$ (72).

Ligand exchange reactions between equimolar amounts of $\text{PPN}[\text{Au}(\text{C}\equiv\text{CH})_2]$ (1) and $\text{PPN}[\text{AuX}_2]$ or $[\text{Au}(\text{PR}_3)_2]\text{ClO}_4$ lead to ethynyl(halo)aurate(I) (4) or ethynyl(phosphine)gold(I) (5) complexes, respectively (see Box 1) (68, 71). The only previously reported complexes of the type 4 are the phenylethynyl derivatives obtained by reacting $[\text{Au}(\text{C}\equiv\text{CPh})]_n$ with a halide (73, 74). Complexes $[\text{Au}(\text{C}\equiv\text{CR})(\text{PR}'_3)]$ analogous to 5 are among the best known alkynyl gold(I) derivatives, and they have been prepared by various methods (45, 48, 72, 75-79).

An alternative synthetic route to alkynylgold(I) complexes is the reaction of chlorogold(I) complexes with the corresponding alkyne in the presence of a base (see Box 2) (71). Although this reaction had been reported (75) to produce some neutral alkynyl(phosphine)gold(I) complexes in the presence

of CuCl , we have proved that the method is more general and that CuCl is not needed. Diethyl or triethylamine can be used to prepare alkynyl complexes with amine (6), phosphine (7, 8) (71) or pentafluorophenyl (10) ligands (80). However, when starting from $[\text{AuCl}(\text{CNBu}^i)]$, NEt_3 must be used to produce the alkynyl(isocyanide)gold complexes (11) (Box 3) because in the presence of NH_4Et_2 , the corresponding alkynyl(carbene)gold complexes (12) form as a result of the reaction of the isocyanide ligand with the secondary amine (71). Although this is a well-known method for the preparation of carbenegold complexes (81-86), we are not aware of the previous synthesis of any alkynyl(carbene)gold(I) complexes like 12 or 14. The use of terminal diynes leads to dinuclear complexes with bridging dialkynyl ligands (8, 13, 14, Boxes 2 and 3) (71). The crystal structure of $[\text{Au}(\text{C}\equiv\text{CSiMe}_3)(\text{CNBu}^i)]$ shows an unprecedented tetrameric aggregate consisting of a central molecule with three others connected to it by aurophilic interactions (70).

The labile alkynyl(diethylamine)gold(I) complexes (6) (Box 2) can only be obtained from $[\text{AuClL}]$ ($\text{L} = \text{tetrahydrothiophene, tht, or AsPh}_3$) and can be used to prepare new alkynylgold derivatives such as 7 or 9. Complexes of the type 6 have previously been prepared by reacting $[\text{Au}(\text{C}\equiv\text{CR})]_n$ with amine (48, 49) or the alkyne with $[\text{Au}(\text{amine})_2]^+$ (49).

GOLD(I) COMPLEXES WITH N-DONOR LIGANDS.

Gold(I) complexes with neutral N-donor ligands are much less common than those with P-donor ligands due to the soft acid nature of the metal centre. We have extended the 'acac method' to the synthesis of complexes with nitrogen donor ligands. Thus, the reactions of [Au(acac)(PPh₃)] with primary, secondary or tertiary ammonium salts [NHRR'₂]⁺X⁻ (1:1), lead to a family of cationic complexes [Au(PPh₃)(NRR'₂)]⁺X⁻ (**15**) (see Box 4) (56). The results of these reactions depend on the nature of the solvent. Acetone can only be used in the synthesis of the tertiary amine complex (**15a**). In all other cases, acetone must be excluded to prevent the formation of iminogold(I) complexes (see below). Diethyl ether was chosen to ensure rapid precipitation of the complexes thus avoiding the formation of oxonium salts [(Au(PR₃)₃O)]⁺ (**16**) due to the presence of adventitious water. All attempts to prepare [(AuPR₃)₂(μ₂-NR'₂)]⁺ (R' = Ph, C₆H₄OMe-4) (see Box 4), even using apparently dry solvents, failed and only the oxonium salts **16**, resulting after hydrolysis, could be isolated (87).

The use of (NH₄)ClO₄ allowed us to synthesize mono-, di- and tetraaurated ammonium salts (Box 5) (87). Depending on the reaction conditions, the amino complexes [Au(NH₃)L]ClO₄ (**17**), or the first amido complexes [(AuL)₂(μ₂-NH₂)]ClO₄ (**18**), or nitrido complexes [(AuL)₄(μ₄-N)]ClO₄ (**19**) can be prepared. This method of synthesis of nitrido gold(I) complexes is simpler and gives a higher yield than those previously reported (88, 89).

Complex [Au(NH₃)₂]Cl (**20**) can be easily prepared by reacting ammonia with [AuCl(tht)] (tht = tetrahydrothiophene) (see Box 6) (87). Other salts of **20** were prepared using [Au(NCPh)₂]⁺ as the starting material (90). Our method is simpler and uses a classical starting material for the synthesis of gold complexes such as [AuCl(tht)] (91, 92). Different salts of **20** can be prepared by reacting **20**·Cl with the appropriate silver or thallium salts. Complex **20** reacts very slowly with acetone to give the first iminogold(I) complexes [Au(NH=CMe₂)₂]⁺X⁻ (**21**·X). The crystal structure of **21**·O₃SCF₃ shows a polymeric chain in which aurophilic contacts along with N-H...O hydrogen bonds could compensate repulsions among the cationic fragments (70). We have also prepared other imino complexes of the types [Au(NH=CMe₂)Cl], [Au{NR=C(R')Me}(PR'₃)₃]⁺ (R = H, C₆H₄OMe-4; R' = Me, Et; R'' = Ph, C₆H₄OMe-4), and [Au(NH=CMe₂)Cl₃] (93). Related complexes [Au(NH=CR₂)Cl] and [Au(NH=CR₂)(PR₃)₃] (R = Ph, NMe₂) have recently been prepared (94, 95).

CONCLUSIONS

We have reported a number of different methods for the synthesis of gold(I) complexes. Thus, the 'acac method' allows the preparation of neutral, anionic or cationic complexes by reacting $[\text{Au}(\text{acac})\text{L}]$ or $[\text{Au}(\text{acac})_2]^-$ with protic acids such as alkynes or ammonium salts. Ligand exchange reactions lead to complexes with two different ligands by reacting symmetrical complexes (eg $[\text{Au}(\text{C}\equiv\text{CH})_2]^- + [\text{AuX}_2]^- \rightarrow 2 [\text{AuX}(\text{C}\equiv\text{CH})]^-$). Several methods have been applied to the synthesis of alkynyl complexes. Replacement of weakly bonded ligands, such as amines or tetrahydrothiophene, enabled the preparation of new alkynyl derivatives or complexes with ammine ligands.

ACKNOWLEDGMENTS

The authors wish to express their appreciation to Professor Peter G Jones for solving the crystal structure of many of the complexes reported here and DGES (Spain) for financial support.

ABOUT THE AUTHORS

Professors María Teresa Chicote and Jose Vicente received their PhD degrees from Zaragoza University under the supervision of Professor Rafael Usón. They spent the year 1976 in Bristol University where they worked with Professor F Gordon A Stone. Since 1980 they have been at Murcia University.

Dr María Dolores Abrisqueta and Dr Pablo González Herrero received their PhD's this year from Murcia University and Rita Guerrero is now studying for her PhD. They have been working under the supervision of Professors Vicente and Chicote on the chemistry of gold(I) complexes described in this paper.

REFERENCES

- 1 P.D. Akrivos, H.J. Katsikis and A. Kourmoutsis, *Coord. Chem. Rev.*, 1997, **167**, 95
- 2 K.I. Grandberg and V.P. Dyadchenko, *J. Organomet. Chem.*, 1994, **474**, 1
- 3 R.V. Parish, *Gold Bull.*, 1998, **31**, 14
- 4 R.V. Parish, *Gold Bull.*, 1997, **30**, 3
- 5 R.V. Parish, *Gold Bull.*, 1997, **30**, 55
- 6 H. Schmidbaur, *Gold Bull.*, 1990, **23**, 11
- 7 O. Steigelmann, P. Bissinger and H. Schmidbaur, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 1399
- 8 H. Schmidbaur, *Chem. Soc. Rev.*, 1995, 391
- 9 F.P. Gabbaï, A. Schier, J. Riede and H. Schmidbaur, *Chem. Ber.*, 1997, **130**, 111
- 10 J.M. Lopez de Luzuriaga, A. Schier and H. Schmidbaur, *Chem. Ber.*, 1997, **130**, 221
- 11 R.E. Bachman and H. Schmidbaur, *Inorg. Chem.*, 1996, **35**, 1399
- 12 J. Vicente, M.T. Chicote, R. Guerrero and P.G. Jones, *J. Am. Chem. Soc.*, 1996, **118**, 699
- 13 M.C. Gimeno and A. Laguna, *Chem. Rev.*, 1997, **97**, 511
- 14 S.J. Berners-Price and P.J. Sadler, *Coord. Chem. Rev.*, 1996, **151**, 1
- 15 S.P. Fricker, *Gold Bull.*, 1996, **29**, 53
- 16 S.L. Best and P.J. Sadler, *Gold Bull.*, 1996, **29**, 87
- 17 S.P. Fricker, *Trans. Met. Chem.*, 1996, **21**, 377
- 18 M. Viotte, B. Gautheron, M.M. Kubicki, I.E. Nifant'ev and S.P. Fricker, *Met.-Based Drugs*, 1995, **2**, 311
- 19 C.K. Mirabelli, R.K. Johnson, C.M. Sung, L. Faucette, K. Muirhead and S.T. Crooke, *Cancer Res.*, 1985, **45**, 32
- 20 S.Y.M. Chooi, P.H. Leung, K.Y. Sim, K.S. Tan and O.L. Kon, *Tetrahedron: Asymmetry*, 1994, **5**, 49
- 21 T. Okada, B.K. Patterson, S.-Q. Ye and M.E. Gurney, *Virology*, 1993, **192**, 631
- 22 T.M. Simon, D.H. Kunishima, D.H. Vibert and A. Lorber, *Cancer Res.*, 1981, **41**, 94
- 23 R. Bayon, S. Coco, P. Espinet, C. Fernandez-Mayordomo and J.M. Martin-Alvarez, *Inorg. Chem.*, 1997, **36**, 2329
- 24 M. Benouazzane, S. Coco, P. Espinet and J.M. Martin-Alvarez, *J. Mater. Chem.*, 1995, **5**, 441
- 25 P. Alejos, S. Coco and P. Espinet, *New J. Chem.*, 1995, **19**, 799
- 26 S. Coco, P. Espinet, S. Falagan and J.M. Martin-Alvarez, *New J. Chem.*, 1995, **19**, 959
- 27 T. Kaharu, R. Ishii and S. Takahashi, *J. Chem. Soc., Chem. Commun.*, 1994, 1349
- 28 M.J. Irwin, G.C. Jia, N.C. Payne and R.J. Puddephatt, *Organometallics*, 1996, **15**, 51
- 29 J.L. Serrano, *Metallomesogens*, VCH: Weinheim, Germany, 1996
- 30 E.Y. Fung, M.M. Olmstead, J.C. Vickery and A.L. Balch, *Coord. Chem. Rev.*, 1998, **171**, 151
- 31 L.H. Gade, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1171
- 32 M.J. Irwin, J.J. Vittal and R.J. Puddephatt, *Organometallics*, 1997, **16**, 3541
- 33 H. Xiao, K.-K. Cheung and C.-M. Che, *J. Chem. Soc., Dalton Trans.*, 1996, 3699
- 34 I.R. Whittall, M.G. Humphrey, M. Samoc and B. Lutherdavis, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 370
- 35 I.R. Whittall, M.G. Humphrey, S. Houbrechts, J. Maes, A. Persoons, S. Schmid and D.C.R. Hockless, *J. Organomet. Chem.*, 1997, **544**, 277
- 36 T.J. Mathieson, A.G. Langdon, N.B. Milestone and B.K. Nicholson, *Chem. Commun.*, 1998, 371
- 37 X. Hong, K.K. Cheung, C.X. Guo and C.M. Che, *J. Chem. Soc., Dalton Trans.*, 1994, 1867
- 38 V.W.W. Yam, S.W.K. Choi and K.K. Cheung, *J. Chem. Soc., Dalton Trans.*, 1996, 3411
- 39 V.W.W. Yam and S.W.K. Choi, *J. Chem. Soc., Dalton Trans.*, 1996, 4227
- 40 B.C. Tzeng, W.C. Lo, C.M. Che and S.M. Peng, *Chem. Commun.*, 1996, 181
- 41 D. Li, X. Hong, C.M. Che, W.C. Lo and S.M. Peng, *J. Chem. Soc., Dalton Trans.*, 1993, 2929
- 42 V.W.W. Yam, S.W.K. Choi and K.K. Cheung, *Organometallics*, 1996, **15**, 1734
- 43 N.C. Payne, R. Ramachandran and R.J. Puddephatt, *Can. J. Chem.*, 1995, **73**, 6
- 44 G.C. Jia, R.J. Puddephatt, J.D. Scott and J.J. Vittal, *Organometallics*, 1993, **12**, 3565
- 45 H. Xiao, Y.X. Weng, S.M. Peng and C.M. Che, *J. Chem. Soc., Dalton Trans.*, 1996, 3155
- 46 G.C. Jia, N.C. Payne, J.J. Vittal and R.J. Puddephatt, *Organometallics*, 1993, **12**, 4771

- 47 G.C. Jia, R.J. Puddephatt, J.J. Vittal and N.C. Payne, *Organometallics*, 1993, **12**, 263
- 48 G.E. Coates and C. Parkin, *J. Chem. Soc., Chem. Commun.*, 1962, 1787
- 49 J. Yau and D.M.P. Mingos, *J. Chem. Soc., Dalton Trans.*, 1997, 1103
- 50 C.J. Aguirre, M.C. Gimeno, A. Laguna, M. Laguna, J.M.L. De Luzuriaga and F. Puente, *Inorg. Chim. Acta*, 1993, **208**, 31
- 51 J. Vicente, M.T. Chicote, I. Saura-Llamas and M.C. Lagunas, *Chem. Commun. (Cambridge)*, 1992, 915
- 52 J. Vicente, M.T. Chicote and M.C. Lagunas, *Inorg. Chem.*, 1993, **32**, 3748
- 53 J. Vicente, M.T. Chicote and P.G. Jones, *Inorg. Chem.*, 1993, **32**, 4960
- 54 J. Vicente, M.T. Chicote, P. González-Herrero, P.G. Jones and B. Ahrens, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1852
- 55 J. Vicente, M.T. Chicote, P. González-Herrero and P.G. Jones, *J. Chem. Soc., Dalton Trans.*, 1994, 3183
- 56 J. Vicente, M.T. Chicote, P. González-Herrero and P.G. Jones, *J. Chem. Soc., Chem. Commun.*, 1995, 745
- 57 J. Vicente, M.T. Chicote and C. Rubio, *Chem. Ber.*, 1996, **129**, 327
- 58 E.J. Fernández, M.C. Gimeno, P.G. Jones, A. Laguna, M. Laguna and J.M. Lopez de Luzuriaga, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 87
- 59 E.J. Fernández, M.C. Gimeno, P.G. Jones, A. Laguna, M. Laguna and J.M. Lopez de Luzuriaga, *J. Chem. Soc., Dalton Trans.*, 1992, 3365
- 60 M.C. Gimeno, A. Laguna, M. Laguna, F. Sanmartin and P.G. Jones, *Organometallics*, 1993, **12**, 3984
- 61 M.C. Gimeno, A. Laguna, M. Laguna, F. Sanmartin and P.G. Jones, *Organometallics*, 1994, **13**, 1538
- 62 J. Vicente, M.T. Chicote, J.A. Cayuelas, J. Fernández-Baeza, P.G. Jones, G.M. Sheldrick and P. Espinet, *J. Chem. Soc., Dalton Trans.*, 1985, 1163
- 63 J. Vicente, M.T. Chicote, I. Saura-Llamas, J. Turpin and J. Fernández-Baeza, *J. Organomet. Chem.*, 1987, **333**, 129
- 64 J. Vicente, M.T. Chicote, I. Saura-Llamas, P.G. Jones, K. Meyer-Bäse and C.F. Erdbrügger, *Organometallics*, 1988, **7**, 997
- 65 J. Vicente, M.T. Chicote and I. Saura-Llamas, *J. Chem. Soc., Dalton Trans.*, 1990, 1941
- 66 J. Vicente, M.T. Chicote, M.C. Lagunas and P.G. Jones, *J. Chem. Soc., Dalton Trans.*, 1991, 2579
- 67 J. Vicente, M.T. Chicote, M.C. Lagunas and P.G. Jones, *Chem. Commun. (Cambridge)*, 1991, 1730
- 68 J. Vicente, M.T. Chicote and M.D. Abrisqueta, *J. Chem. Soc., Dalton Trans.*, 1995, 497
- 69 J. Vicente, M.T. Chicote, R. Guerrero and P.G. Jones, *J. Chem. Soc., Dalton Trans.*, 1995, 1251
- 70 J. Vicente, M.T. Chicote, M.-D. Abrisqueta, R. Guerrero and P.G. Jones, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1203
- 71 J. Vicente, M.T. Chicote, M.D. Abrisqueta and P.G. Jones, *Organometallics*, 1997, **16**, 5628
- 72 H. Werner, H. Otto, T. Ngo-Kha and C. Burschka, *J. Organomet. Chem.*, 1984, **262**, 123
- 73 O.M. Abu-Salah and A.R. Al-Ohaly, *Inorg. Chim. Acta*, 1983, **77**, L159
- 74 O.M. Abu-Salah, A.R. Al-Ohaly, S.S. Al-Showiman and I.M. Al-Najjar, *Transition Met. Chem.*, 1985, **10**, 207
- 75 M.I. Bruce, E. Horn, J.G. Matison and M.R. Snow, *Aust. J. Chem.*, 1984, **37**, 1163
- 76 R.J. Cross and M.F. Davidson, *J. Chem. Soc., Dalton Trans.*, 1986, 411
- 77 C.M. Mitchell and F.G.A. Stone, *J. Chem. Soc., Dalton Trans.*, 1972, 102
- 78 A. Johnson and R.J. Puddephatt, *J. Chem. Soc., Dalton Trans.*, 1977, 1384
- 79 F. Bonati, A. Burini, B.R. Pietroni, E. Giorgini and B. Bovio, *J. Organomet. Chem.*, 1988, **344**, 119
- 80 J. Vicente, M.T. Chicote and M.D. Abrisqueta, unpublished results
- 81 J.A. McCleverty and M.M.M. Da Mota, *J. Chem. Soc., Dalton Trans.*, 1973, 2571
- 82 G. Minghetti, L. Baratto and F. Bonati, *J. Organomet. Chem.*, 1975, **102**, 397
- 83 F. Bonati and G. Minghetti, *J. Organomet. Chem.*, 1973, **59**, 403
- 84 J.E. Parks and A.L. Balch, *J. Organomet. Chem.*, 1974, **71**, 453
- 85 R. Usón, A. Laguna, J. Vicente, J. García, B. Bergareche and P. Brun, *Inorg. Chim. Acta*, 1978, **28**, 237
- 86 R. Usón, A. Laguna, J. Vicente, J. García and B. Bergareche, *J. Organomet. Chem.*, 1979, **173**, 349
- 87 J. Vicente, M.T. Chicote, R. Guerrero, P.G. Jones and M.C. Ramírez de Arellano, *Inorg. Chem.*, 1997, **36**, 4438
- 88 K. Angermaier and H. Schmidbaur, *Chem. Ber.*, 1995, **128**, 817
- 89 E.G. Perevalova, E.I. Smyslova, V.P. Djadchenko, K.I. Grandberg and A.N. Nesmeyanov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1980, 1455. *Chem. Abstr.*, 1980, **93**, 178615h
- 90 D.M.P. Mingos, J. Yau, S. Menzer and D.J. Williams, *J. Chem. Soc., Dalton Trans.*, 1995, 319
- 91 R. Usón, J. Laguna and J. Vicente, *J. Chem. Soc., Chem. Commun.*, 1976, 353
- 92 R. Usón, J. Laguna and J. Vicente, *J. Organomet. Chem.*, 1977, **131**, 471
- 93 J. Vicente, M.T. Chicote and R. Guerrero, unpublished results
- 94 W. Schneider, A. Bauer, A. Schier and H. Schmidbaur, *Chem. Ber.*, 1997, **130**, 1417
- 95 W. Schneider, A. Bauer and H. Schmidbaur, *J. Chem. Soc., Dalton Trans.*, 1997, 415