ERRATUM

EUROPEAN JOURNAL OF DRUG METABOLISM AND PHARMACOKINETICS

1995, Vol. 20, No. 1, pp. 73–77

Effect of sulfated xylans during the interaction of [¹²⁵I]-thrombin with antithrombin III or heparin cofactor II of human plasma

R.B. SIMMONS, G.R. NEWTON and V.M. DOCTOR

Chemistry Department and Cooperative Agriculture Research Center, Prairie View A&M University, Prairie View, Texas, USA

Fig 1 : Effect of OSXS and heparin on the complexation of [¹²⁵I]-thrombin with AT-III and HC-II of human plasma. Please refer to Materials and Methods for details. Lane A and B represented 40 µg and 8 µg/ml of heparin, respectively, while lanes D and E represented 40 µg and 8 µg/ml of OSXS, respectively. The plasma control (lane C) did not show complex formation with [¹²⁵I]-thrombin. The mixture was incubated for 30 s before heating with the reducing mixture.

The upper legends to Figure 1 in this manuscript were incorrectly positioned (p.75).

The legend 'Thr–HC–II' should have been above 'Thr–AT–III' instead of below. This agrees with the correct results as given in Table I of the published article : i.e. OSXS (Lanes D & E) enhanced the interaction of [¹²⁵I]-thrombin with **both** AT–III and HC–II, whereas heparin (Lanes A & B) only enhanced the interaction of [¹²⁵I]-thrombin with AT–III.

We apologise for any problems this may have caused.