113. Halperin JR, Murphy B. Extrapyramidal reaction to ondansetron [letter]. *Cancer* 1992; 69: 1275
114. de Bruijn KM. The development of tropisetron in its clinical perspective. *Ann Oncol* 1993; 4 Suppl. 3: S19-S23
115. Gralla RJ. Antiemetic drugs for chemotherapeutic support. Current treatment and rationale for development of newer agents. *Cancer* 1992; 70 Suppl.: 1003-6
116. Bourke JP. Does granisetron share the cardiac arrhythmogenic potential of other 5-HT₃ receptor antagonists? [abstract]. Antiemetic control: maximizing the benefits. Satellite symposium, 14 November, 1993, 7th European Conference on Clinical Oncology and Cancer Nursing (ECCO), Jerusalem, Israel
117. Leigh TJ, Link CG, Fell GL. Effects of granisetron and lorazepam, alone and in combination, on psychometric performance. *Br J Clin Pharmacol* 1991; 31 (3): 333-6
118. Link CGG, Leigh TJ, Fell GL. Effects of granisetron and lorazepam, alone and in combination, on the EEG of human volunteers. *Br J Clin Pharmacol* 1991; 31: 93-7
119. Milne RJ, Heel RC. Ondansetron. Therapeutic use as an antiemetic. *Drugs* 1991; 41: 574-95
120. Pinkerton CR, on behalf of the Granisetron Study Group. Experience with granisetron in children [abstract]. Criteria for evaluation of the ideal antiemetic. Satellite symposium, 8 November, 1992, 17th Congress of the European Society for Medical Oncology, Lyon, France
121. Leigh TJ, Link CGG, Fell GL. Effects of granisetron and haloperidol, alone and in combination, on psychometric performance and the EEG. *Br J Clin Pharmacol* 1992; 34: 65-70
122. Allen JC, Gralla R, Reilly L, et al. Metoclopramide: dose-related toxicity and preliminary anti-emetic studies in children receiving cancer chemotherapy. *J Clin Oncol* 1985; 3 (8): 1136-41
123. Terrin BN, McWilliams NB, Maurer HM. Side effects of metoclopramide as an anti-emetic in childhood cancer chemotherapy. *J Paediatr Child Health* 1984; 104: 138-40
124. Gandara DR, Harvey WH, Monaghan GG, et al. The delayed-emesis syndrome from cisplatin: phase III evaluation of ondansetron versus placebo. *Semin Oncol* 1992; 19 Suppl. 10: 67-71
125. Jones AL, Lee GJ, Bosanquet N. The budgetary impact of 5-HT₃ receptor antagonists in the management of chemotherapy-induced emesis. *Eur J Cancer* 1992; 29A: 51-6
126. Kirchner V, Aapro M, Terrey J-P. Antiemetic treatment costs in chemotherapy; the economic evaluation of granisetron [abstract]. *Ann Oncol* 1992; 3 Suppl. 5: 185
127. FDC Reports, The Pink Sheet, T&G-3, May 30, 1994

Correspondence: Yvonne E. Yarker, Adis International Limited, 41 Centorian Drive, Private Bag 65901, Mairangi Bay, Auckland 10, New Zealand.

Erratum and Addendum

Vol. 48, No. 3, p. 445: In table V in the section on ear, nose and throat disorders, the dosage of naproxen (NAP) in the study by Nouri and Monti^[94] should be 500 mg/day.

[R Davis, RN Brogden. Nimesulide: an update of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy. *Drugs* 1994; 48 (3): 431-454]

Vol. 48, No. 2, p. 304: The first paragraph of section 1.3.1 cites a single *in vitro* study that showed roxithromycin to be active against *Mycobacterium tuberculosis*. However, since the completion of this manuscript it has been drawn to our attention that more recent data indicate that this is not the case. Accordingly, the following sentence should be added at the end of the first paragraph in section 1.3.1:

'More recent data from unpublished studies have shown roxithromycin to be inactive against *M. tuberculosis* (MIC range 16 to >64 mg/L) [data on file, Roussel Uclaf].'

[A Markham, D Faulds. Roxithromycin: an update of its antimicrobial activity, pharmacokinetic properties and therapeutic use. *Drugs* 1994; 48 (2): 297-326]