87. Friedel HA, Langtry HD, Buckley MM. Nabumetone: a reappraisal of its pharmacology and therapeutic use in rheumatic diseases. *Drugs* 1993; 45: 131-56
88. Balfour JA, Buckley MM-T. Etodolac: a reappraisal of its pharmacology and therapeutic use in rheumatic diseases and pain states. *Drugs* 1991; 42: 274-99
89. Lanza FL. Gastrointestinal toxicity of newer NSAIDs. *Am J Gastroenterol* 1993; 88: 1318-23
90. Skeith KJ, Wright M, Davis P. Differences in NSAID tolerability profiles: fact or fiction? *Drug Saf* 1994; 10: 183-95
91. Rees Willet L, Carson JL, Strom BL. Epidemiology of gastrointestinal damage associated with nonsteroidal anti-inflammatory drugs. *Drug Saf* 1994; 10: 170-81

Correspondence: C. Rhoda Lee, Adis International Limited, 41 Centorian Drive, Mairangi Bay, Private Bag 65901, Auckland 10, New Zealand.

Errata

Vol. 46, No. 6, 1993, page 1026: The third sentence in the first paragraph of the Pharmacological Properties Summary should read, 'The beneficial effects of acarbose on postprandial glucose levels have been confirmed in patients with insulin-dependent or non-insulin-dependent diabetes mellitus (IDDM or NIDDM), and postprandial insulin and triglyceride levels were also lowered.'

page 1027: The second sentence of paragraph 2 of the Tolerability Summary should read,

'However, analysis of data from phase III US studies indicated that anaemia and elevated transaminase levels were significantly more common in patients receiving acarbose (at a dosage greater than the maximum currently recommended) than in placebo recipients.'

page 1028: The fourth sentence in the first paragraph of the Dosage and Administration Summary should read:

'Contraindications to acarbose use include age less than 12 years, inflammatory bowel disease . . . '.

page 1047: The following sentence should be added at the end of the fourth paragraph of section 3.2 Systemic Effects:

'Furthermore, patients in these trials received acarbose at a dosage greater than the maximum currently recommended.'

page 1048: The first sentence in the third paragraph of section 5 Dosage and Administration should read:

'Acarbose is contraindicated in children aged less than 12 years, and in patients with inflammatory bowel disease . . . '.

[Balfour JA, McTavish D. Acarbose: an update of its pharmacology and therapeutic use in diabetes mellitus. *Drugs* 1993; 46 (6): 1025-1054]

Vol. 47, No. 5, 1994, page 847: The last sentence of the fifth paragraph of section 5 Dosage and Administration should read:

'Serum teicoplanin concentrations should be monitored in patients with creatinine clearance ≤ 20 ml/min to ensure that therapeutic dosage is achieved . . . '.

[Brogden RN, Peters DH. Teicoplanin: a reappraisal of its antimicrobial activity, pharmacokinetic properties and therapeutic efficacy. *Drugs* 1994; 47 (5): 823-854]