ported drug-related adverse effects were vaginitis, nausea and vomiting, most of which were only mild or moderate in severity.

All of these findings indicate that grepafloxacin will be useful in the treatment of genitourinary tract infections caused by *N. gonorrhoeae*, *C. trachomatis* or both. ▲

References

1. Zenilman JM, Neumann T, Patton M, et al. Antibacterial activities of OPC-17116, ofloxacin, and ciprofloxacin against 200 isolates of *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 1993 Oct; 37: 2244-6
2. Wise R, Andrews JM, Brenwald N. The in-vitro activity of OPC-17116, a new 5-methyl substituted quinolone. *J Antimicrob Chemother* 1993 Apr; 31: 497-504
3. Mroczkowski TF, Millikan LE, Bean KA, et al. Randomized open-label trial of OPC-17116 vs cefixime in the treatment of uncomplicated gonococcal cervicitis [abstract]. *International Union Against Venereal Diseases and Treponematoses World STD/AIDS Congress*, Singapore, 19-23 Mar 1995: 159
4. Hook EW, McCormack WM, Martin DH, et al. Comparison of single-dose oral grepafloxacin with cefixime for treatment of uncomplicated gonorrhea in men [poster no. LM19]. *36th Interscience Conference on Antimicrobial Agents and Chemotherapy* 1996; New Orleans (LA); 15-18 Sep 1996
5. Mroczkowski TF, Bean KA, Martin DH. A dose ranging study of a new quinolone, OPC-17116, for the treatment of *Chlamydia trachomatis* infection in women [abstract]. *Sex transm Dis* 1994 Mar-Apr; 21 Suppl.: 106-7

Errata

Vol. 53, No. 3, page 512: In the first paragraph of the Pharmacological Properties section of the summary, the second sentence should read, 'Adapalene has a high affinity for the subtypes $RAR\beta$ and $RAR\gamma$, the latter being found predominantly in the epidermis'.

page 513: In the Application section of the summary, the last paragraph should read, 'Although adapalene was not detected in the plasma of patients treated topically with the drug, it *should not be used during pregnancy*'.

page 518: In section 5, the last paragraph should read, '... has not been ruled out.^[26] *It is recommended that adapalene should not be used during pregnancy.*'

[Brogden RN, Goa KL. Adapalene: A review of its pharmacological properties and clinical potential in the management of mild to moderate acne. *Drugs* 1997 Mar; 53 (3): 511-9]