- after cardiac surgery: effects of prophylaxis. *Br Heart J* 1994; 72: 186-9
18. Jahr JS, Burckart G, Smith SS, et al. Effects of famotidine on gastric pH and residual volume in paediatric surgery. *Acta Anaesthesiol Scand* 1991; 35: 457-60
19. Miyake S, Yamada M, Iwamoto H, et al. Effect of a new H₂-blocker, famotidine, in reflux esophagitis among severely handicapped children. *Clin Ther* 1987; 9 (5): 548-58
20. Carroccio A, Pardo F, Montalto G, et al. Use of famotidine in severe exocrine pancreatic insufficiency with persistent maldigestion on enzymatic replacement therapy: a long-term study in cystic fibrosis. *Dig Dis Sci* 1992; 37 (9): 1441-6
21. Chalmers DM, Brown RC, Miller MD, et al. The influence of long-term cimetidine as an adjuvant to pancreatic enzyme therapy in cystic fibrosis. *Acta Paediatr Scand* 1985; 74: 114-7

Correspondence and reprints: Dr *Laura P. James*, Division of Pediatric Clinical Pharmacology, Arkansas Children's Hospital, 800 Marshall Street, Little Rock, AR 72202, USA.

Errata

Vol. 30, No. 6, page 468: The key for figure 2 shows filled circles as levodopa plasma concentrations, this should be *tapping scores*.

[Contin M, Riva R, Albani F, Baruzzi A. Pharmacokinetic optimisation in the treatment of Parkinson's disease. *Clin Pharmacokinet* 1996 Jun; 30 (6): 463-481]