26. Pye JK, Mansell RE, Hughs LE. Clinical experience of drug treatments for mastalgia. *Lancet* 1985; II (8451): 373-7
27. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52 705 women with breast cancer and 108 411 women without breast cancer. *Lancet* 1997; 350 (9084): 1047-59
28. Ettinger B, Genant HK, Cann CE. Long-term estrogen replacement therapy prevents bone loss and fractures. *Ann Intern Med* 1985; 102: 319-24
29. Quigley MET, Martin PL, Burnier AM et al. Estrogen therapy arrests bone loss in elderly women. *Am J Obstet Gynecol* 1987; 156: 1516-23
30. Draper MW, Flowers DE, Huster Wj et al. Effects of raloxifene (LY 139481) on biochemical markers of bone and lipid metabolism in healthy postmenopausal women. In: Christiansen C, Riis B, editors. *Proceedings of the Fourth International Symposium on Osteoporosis and Consensus Development Conference*. Aalborg, Denmark: Handelstrykkeriet Aalborg Aps, 1993: 119-21
31. Wilcox G, Wahlquist ML, Burger HG et al. Oestrogen effects of plant derived foods in postmenopausal women. *BMJ* 1990; 301: 905-6

Correspondence and reprints: Dr K.A. McKinney, Department of Obstetrics and Gynaecology, Queen's University of Belfast, Institute of Clinical Science, Grosvenor Road, Belfast BT12 6BJ, Northern Ireland.
E-mail: w.thompson@qub.ac.uk

Errata

Vol. 54, No. 4, page 574: In figure I, the dosage of furosemide in the 'Ascites grade 3' and 'Refractory ascites' columns should read 40-160 mg/day and 160 mg/day, respectively.

[Bataller R, Ginès P, Arroyo V. *Practical recommendations for the treatment of ascites and its complications*. *Drugs* 1997 Oct; 54 (4): 571-80]

Vol. 55, No. 6, page 762: In the first paragraph of section 2.2, some reference citations were inadvertently omitted. Lines 8 to 16 of that paragraph should read: 'Early clinical studies with live oral heterologous (bovine or rhesus) vaccines showed good efficacy against severe rotavirus gastroenteritis in Scandinavia and the US,^[15,16] but the protection was insufficient in developing countries.^[17] Therefore, these vaccines were not developed further. Similarly, human rotavirus (neonatal) strains also failed to induce protection against diarrhoeal illness when used as vaccines.^[18]'

[Eskola JJ. *Childhood immunisation today*. *Drugs* 1998 Jun; 55 (6): 759-66]