Therapy, Pharmacoeconomics and Pharmacovigilance

11.3 Ambulatory Monitoring of Blood Pressure (AMBP) in Young People with Obstructive Sleep Apnoea Syndrome (OSAS)

R. Muschera' (1), M. D'avino (1), G. Caruso (1), E. Anastasio (1), D. Caruso (1), F. De Michele (1), on behalf of R. Muschera'
(1)AORN Cardarelli, Napoli, Italy

Introduction. Obstructive Sleep Apnoea Syndrome (OSAS) is often associated with hypertension and is a new cardiovascular risk factor independent of age, gender and body mass index (NEJM 342: 1378-84,2000). OSAS occurs in all age groups and both genders and hypertensive target organ damage in the young people is not known. We have calculated the cardiovascular risk in a young people with serious OSAS (RDI>30).

Methods. 20 young people (mean age: 39± 6,9; Male: 18 Female; We were evaluated with office blood pressure measurements (OBPM) and AMBP, ECG, echocardiography, carotid echography and plasma total LDL Cholesterol.

Results. 14 young people are obese with mean BMI: 37,6±6, 16 smoker. Familiarity hypertension is in 15/20 young people. At clinic blood pressure mean SBP 132±10 mmHg and mean DBP 86± 12 mmHg, 8/20 people with hypertension. At ambulatory monitoring of blood pressure the mean PAS= 145± 16 mmHg and PAD: 92±12 mmHg, 14/20 people with hypertension and 10 people no dippers. At the ECG the Romhilt-Estes point 2±1. At the echocardiography the LV mass/ h2,7 41,35±6,5 gr/m with diastolic dysfunction in 8/20 people. The carotid artery intima/media thickness is 0,96±0,2 mm. Mean LDL Cholesterol levels are 124±36mg/dl.

Conclusions. The clinic blood pressure in young people with OSAS isn't good method for to diagnose the hypertension in young people with OSAS because there are many people with masked hypertension. In these people, the best method of monitoring blood pressure is the AMBP because it diagnoses the alteration of circadian profile of blood pressure. The masked nocturnal hypertension went up with an elevated nocturnal BP and normal OBP levels had a bigger target organ damage.