

Lie n -algebras of BPS charges

Hisham Sati^{a,b} and Urs Schreiber^c

^a *University of Pittsburgh,
Pittsburgh, PA, 15260 U.S.A.*

^b *Mathematics Program, Division of Science and Mathematics, New York University Abu Dhabi,
Saadiyat Island, Abu Dhabi, UAE*

^c *Mathematics Institute of the Academy,
Žitna 25, Praha 1, 115 67 Czech Republic¹*

E-mail: hsati@pitt.edu, schreiber@math.cas.cz

ABSTRACT: We uncover higher algebraic structures on Noether currents and BPS charges. It is known that equivalence classes of conserved currents form a Lie algebra. We show that at least for target space symmetries of higher parameterized WZW-type sigma-models this naturally lifts to a Lie $(p + 1)$ -algebra structure on the Noether currents themselves. Applied to the Green-Schwarz-type action functionals for super p -brane sigma-models this yields super Lie $(p + 1)$ -algebra refinements of the traditional BPS brane charge extensions of supersymmetry algebras. We discuss this in the generality of higher differential geometry, where it applies also to branes with (higher) gauge fields on their worldvolume. Applied to the M5-brane sigma-model we recover and properly globalize the M-theory super Lie algebra extension of 11-dimensional superisometries by 2-brane and 5-brane charges. Passing beyond the infinitesimal Lie theory we find cohomological corrections to these charges in higher analogy to the familiar corrections for D-brane charges as they are lifted from ordinary cohomology to twisted K-theory. This supports the proposal that M-brane charges live in a twisted cohomology theory.

KEYWORDS: Differential and Algebraic Geometry, p-branes

ARXIV EPRINT: [1507.08692](https://arxiv.org/abs/1507.08692)

¹On leave at Bonn.

Contents

1	Introduction	1
2	Poisson bracket Lie n-Algebras	4
3	Truncation to ordinary Lie algebras	10
4	Transgression to ordinary Lie algebras	13
5	Application to brane charges	16

1 Introduction

The correspondence between infinitesimal symmetries of a local Lagrangian form and conserved currents of its induced Euler-Lagrange equations, i.e. Noether's first variational theorem, is one of the cornerstones of mathematical field theory. If the Lagrangian under consideration is preserved only up to an exact term, as is the case for sigma-models of Wess-Zumino-Witten-type (WZW), then the Lie algebra of conserved currents is an extension of the symmetry algebra (see for instance [3, section 8] for review from a perspective that will be useful here). Furthermore, the extensions arising this way are interesting enough that they acquired a life of their own in pure mathematics: for the 2-dimensional WZW model they transgress to the *current algebras* from affine Lie algebra theory (see e.g. [28] for a survey).

Key examples of higher-dimensional WZW models are the Green-Schwarz-type sigma models describing the worldvolume theory of all super p -branes [13, 21, 23]. Their extended current algebras have been argued in [2] to yield extensions of superisometry algebras of curved superspacetimes by brane charges [24, 41]. These constitute what characterizes BPS states of supergravity and hence of supersymmetric field theories (e.g. [12]).

However, in traditional constructions one considers a Lie bracket only on equivalence classes of current forms modulo exact forms, the Dickey bracket [14], [8, section 3]. Whenever one is in such a situation where an algebraic structure exists on the homology of a chain complex, it is natural to ask whether a homotopy-theoretic refinement of the algebraic structure may exist on the chain complex itself, where algebraic conditions such as Jacobi identities may hold only up to higher coherent homotopies. For the case of Lie algebras this means lifting to L_∞ -algebras (see [17, 36] for review and references), called Lie $(p + 1)$ -algebras when the underlying chain complex is concentrated in degrees 0 up to p .

For the case of Lie algebras of conserved Noether currents, a strategy to lift to an L_∞ -algebra had been proposed in [7, 27]. However, the construction there only applies

when all relevant higher charge cohomology groups vanish, resulting in an L_∞ -algebra that is in fact quasi-isomorphic to the plain Lie algebra of Noether currents. Hence the general question remained open.

What we do here for field theories of higher dimensional and parameterized WZW-type is the following:

1. We identify the correct higher current algebra for higher WZW-type Lagrangians as the higher Poisson bracket L_∞ -algebra of local observables considered in [32], for the pre- $(p + 1)$ -plectic form being the higher WZW curvature form on target spacetime. In fact we find two different but quasi-isomorphic L_∞ -algebras of higher WZW-type Noether currents, following [17], corresponding to the two different incarnations of the Dickey bracket observed in [8, (3.10)–(3.11)].
2. We show that the L_∞ -Heisenberg-Kostant-Souriau extension of [17, theorem 3.3.1] is the homotopy-theoretic lift of the central extension of infinitesimal symmetries via Noether currents by de Rham cohomology classes (“topological currents”) as in [42, p. 203].
3. We indicate how the higher stacky Heisenberg-Kostant-Souriau extension of [16] generalizes this to higher gauged WZW models (such as the Green-Schwarz-type sigma-model for the M5-brane) and produces cohomological corrections to the naive brane charges.

Acknowledgments

We thank Domenico Fiorenza for discussion of truncation of L_∞ -algebra extensions in section 3, and we thank Jim Stasheff for historical comments. U.S. thanks Igor Khavkine for discussion of higher WZW charge algebras in traditional variational calculus; a joint account dedicated to this aspect is in preparation. U.S. was supported by RVO:67985840.

We review some of the background material in the context of local observables in section 2. The algebras we discuss in this letter form a hierarchy of the following form (see [17, 36] for discussion and references).

- L_∞ -algebras $\in L_\infty\text{Alg}$. These have k -ary brackets for all $k \in \mathbb{N}$, $k \geq 2$ on a chain complex;
- dg-algebras $\in \text{dgLieAlg} \hookrightarrow L_\infty\text{Alg}$. These are L_∞ -algebras with at most binary brackets, hence Lie algebras in the category of chain complexes;
- Lie n -algebras $\in \text{Lie}_n\text{Alg} \hookrightarrow L_\infty\text{Alg}$. These are L_∞ -algebras whose underlying chain complex is concentrated in degree 0 to $(n - 1)$, also called n -term L_∞ -algebras;
- Lie algebras $\in \text{LieAlg} = \text{Lie}_1\text{Alg}$. These are the usual Lie algebras, i.e. Lie 1-algebras, viewed as L_∞ -algebras or dg-Lie algebras whose underlying chain complex is concentrated in degree 0.

This concept of *Lie n -algebras* in the homotopy theoretic sense of n -term L_∞ -algebras, originally due to Stasheff [25], is different from a vaguely similar definition due to Filippov that has come to be known as *n -Lie algebras*. But the Filippov 3-Lie algebras found on the worldvolume of coincident M2-branes are a special case of metric L_∞ -algebras [31, section 2.5]. Notice that (super-) L_∞ -algebras play a role in supergravity and super p -brane sigma-models already since [1], where they appear in the dual but equivalent guise of their Chevalley-Eilenberg dg-algebras (“FDA”s), as explained in [36].

The key motivation for finding an L_∞ -algebra of currents is that it retains more local information about the system at hand. This is in the general spirit of refining field theories to what are called “extended” or “multi-tiered” field theories; we refer the reader to the introductions of [17, 20] for more on this general background. The urge to refine Lie 1-algebras of infinitesimal symmetries to L_∞ -algebras becomes more pronounced as one aims to pass from infinitesimal to finite symmetries. For the example of field theories of WZW type, the Lagrangian is not a globally defined differential form, but is rather a connection on a higher bundle (higher gerbe) on field space. Here the only way even to see the full structure of the finite symmetries is to work in homotopy theory.

Once we have the higher algebras characterized as above, we will describe two ways of getting back to ‘lower’ algebras (in the categorical sense), including ordinary Lie algebras:

1. *Truncation*: in section 3 we consider appropriate truncations of higher algebras down to more classical structures, by controlling the length of the corresponding chain complex. Given any homotopy-theoretic higher structure, hence a homotopy n -type, there its “truncation” down to a 0-type, obtained by quotienting out all symmetries. This operation in general forgets much relevant structure, but the resulting object is sometimes more easily recognized and compared to more traditional concepts. We characterize the corresponding current algebra obtained by truncating the higher Poisson algebras.
2. *Transgression*: we explain in section 4 how the full L_∞ -algebra of conserved currents admits maps to all its transgressions, hence to the results of evaluating conserved currents on cycles of any dimension. For example, the current Lie 2-algebra of the 2-dimensional WZW model transgresses to the “affine” centrally extended loop Lie algebra of currents by integrating currents over a circle. The problem of obtaining the 1-algebraic structure of loop spaces from a 2-algebraic (stacky, gerby) structure on the original base spaces by transgression had been highlighted prominently in [11, p. 249].

Then in section 5 we discuss the application of the above construction to super p -brane sigma-models.

- Application to the Green-Schwarz-type action functionals for super p -brane sigma-models yields super Lie $(p+1)$ -algebra refinements of the traditional BPS brane charge central extensions of supersymmetry algebras. We discuss this in the generality of higher geometry where it applies also to branes with (higher) gauge fields on their worldvolume.

- Specifically, application to the M5-brane sigma-model recovers the M-theory super Lie algebra extension of 11-dimensional superisometries by 2-brane and 5-brane charges.
- Passing beyond the infinitesimal Lie theory we find cohomological corrections to the above charges in higher analogy to the familiar corrections for D-brane charges as they are lifted from ordinary cohomology to twisted K-theory. This supports the proposal that M-brane charges live in a twisted cohomology theory [33–35] and is also supported by the companion analysis in [22].

2 Poisson bracket Lie n -Algebras


We start by briefly reviewing the pertinent concepts and results from [17] in a way that makes manifest their relation to conserved currents. In order to see the origin of the higher symmetry in current algebras fully transparently, it is useful to adopt the following general perspective.

Traditionally a local Lagrangian for a $(p + 1)$ -dimensional field theory is thought of as a suitable differential $(p + 1)$ -form \mathbf{L} on the jet bundle of the field bundle of the theory. (If a reference volume form \mathbf{vol} on spacetime/worldvolume is chosen, then \mathbf{L} is proportional to that form, $\mathbf{L} = L\mathbf{vol}$, and the coefficient function L is what in many textbooks is regarded as the Lagrangian.) However, it is well known, though perhaps not widely appreciated, that for some key field theories of interest (locally variational theories [15]) it is not in fact globally true that the Lagrangian is a differential $(p + 1)$ -form. Rather, in general it is only locally so, but is globally a $(p + 1)$ -form connection on a “ $(p + 1)$ -bundle” or “ p -gerbe”. This is notably the case for field theories of Wess-Zumino-Witten type; see [21] for review of this fact and for pointers to the literature.

Even if \mathbf{L} happens to be given by a globally defined $(p + 1)$ -form, it is still useful to regard this as the $(p + 1)$ -connection form on a trivial $(p + 1)$ -bundle over field space, because it is this perspective that makes immediately clear that and how there are higher symmetries-of-symmetries.

In order to capture this state of affairs mathematically, it is useful to consider a simple object with a pretentious name: the universal smooth moduli stack of $(p + 1)$ -connections, which we denote by $\mathbf{B}^{p+1}\mathbf{U}(1)_{\text{conn}}$. The reader may find exposition and introduction for such objects in the context of field theory in [20], but this object is easily described and understood already by its defining universal property. That is, being a *smooth universal moduli stack for higher connections* means precisely that it is a generalized smooth space of sorts, with the following characterizing properties: for X any smooth manifold, then

1. smooth functions $\nabla: X \rightarrow \mathbf{B}^{p+1}\mathbf{U}(1)_{\text{conn}}$ correspond equivalently to $(p + 1)$ -form connections on X ;



equivalently to gauge transformations between two such connections ∇_1 and ∇_2 ;

- 3. smooth homotopies-of-homotopies correspond to gauge-of-gauge transformations, and so forth, up to p -fold homotopies-of-homotopies.

When the connections ∇_i are given by globally defined differential $(p+1)$ -forms θ_i , then a gauge transformation between these is nothing but a p -form Δ such that $\theta_2 = \theta_1 + \mathbf{d}\Delta$, where \mathbf{d} denotes the de Rham differential. A gauge-of-gauge transformation is accordingly $(p-1)$ -form, etc. up to order- p gauge-of-gauge transformation which are (not 0-forms but) $U(1)$ -valued functions.

Remark 2.1. (i) One immediate advantage of “modulating” connections and differential forms by maps into an object such as $\mathbf{B}^{p+1}U(1)_{\text{conn}}$ is that it makes their “contravariance” manifest while keeping track of higher gauge transformations. Namely, given a $(p+1)$ -form connection ∇ on X and given a map $\phi : Y \rightarrow X$ between spaces, then the pullback connection $\phi^*\nabla$ is simply the one modulated by the composite map

$$\phi^*\nabla : Y \xrightarrow{\phi} X \xrightarrow{\nabla} \mathbf{B}^{p+1}U(1)_{\text{conn}} .$$

- (ii) So suppose then that X is some space of fields of a field theory, typically the jet bundle to the field bundle. Then it is clear what a *symmetry* of such ∇ should be: namely a diffeomorphism

$$X \xrightarrow[\simeq]{\phi} X$$

of the field space, such that ∇ is preserved by this up to a specified gauge transformation $\eta : \phi^*\nabla \xrightarrow{\simeq} \nabla$.

- (iii) In the diagrammatic language that we have set up, this just means that a symmetry is a pair (ϕ, η) labeling a diagram of the form

$$\begin{array}{ccc} X & \xrightarrow{\phi} & X . \\ & \searrow \nabla & \swarrow \nabla \\ & \mathbf{B}^{p+1}U(1)_{\text{conn}} & \end{array} \quad \begin{array}{c} \eta \\ \simeq \end{array}$$

This is equivalent to saying that if ∇ happens to be given by a globally defined $(p+1)$ -form θ , then the homotopy is given by a p -form η such that

$$\phi^*\theta - \theta = \mathbf{d}\eta .$$

Moreover, if we have a smooth 1-parameter flow $t \mapsto (\phi(t), \Delta(t))$ of such symmetries, then differentiating this relation with respect to t yields the expression of an infinitesimal symmetry of ∇ :

$$\mathcal{L}_v\theta = \mathbf{d}\Delta_v ,$$

where v is the vector field of the flow $t \mapsto \phi(t)$, with \mathcal{L}_v its Lie differentiation, and where $\Delta_v = \frac{d}{dt}\eta$. Notice that if one thinks of $\theta = \mathbf{L}$ as being the Lagrangian form of a field theory, then this is precisely the expression for a “weak” symmetry of the Lagrangian, i.e. a transformation that leaves the Lagrangian invariant up to addition of an exact term $\mathbf{d}\Delta_v$.

It is useful to equivalently rewrite this as follows: let $\omega = F_{\nabla}$ denote the curvature $(p+2)$ -form of the $(p+1)$ -form connection. If the connection is given by a globally defined differential $(p+1)$ -form θ , then this is simply $\omega = \mathbf{d}\theta$. Cartan's formula then gives that the contraction is exact

$$\iota_v \omega = -\mathbf{d}J_v$$

with

$$J_v := \iota_v \theta - \Delta_v$$

a kind of Legendre transform of the weakening term.

If we do think of $\mathbf{L} = \theta$ as being a WZW-type local Lagrangian, then the expression $J_v = \iota_v \theta - \Delta_v$ is the conserved Noether charge induced by the symmetry v . This is the interpretation that we are interested in here, as it is what connects to the conceptualization of BPS charges from Noether currents of higher WZW terms in [2].

(More generally, when \mathbf{L} is not just a WZW term, then one has to refine the discussion, take X above to be a jet bundle and incorporate the horizontal/vertical decomposition of the variational bicomplex of forms on that bundle. This further generality will be discussed elsewhere, here we focus on topological terms of higher WZW type.)

For $p = 0$ the above relations are well-known from symplectic geometry, for the case that ω is a (pre-)symplectic form. Since this is the correct analogy, we also say that a closed $(p+2)$ -form ω is a *multisymplectic* $(p+2)$ -form or a $(p+1)$ -plectic form [32]. In view of this it is sensible to agree on the following terminology, modeled directly on the traditional case given by $p = 0$.

Definition 2.2. (i) A closed $(p+2)$ -form ω on a manifold X is also called a *pre- $(p+1)$ -plectic form*, and the pair (X, ω) a *pre- $(p+1)$ -plectic manifold*.

(ii) Given such, then a vector field v such that $\mathcal{L}_v \omega = 0$ called a *pre- $(p+1)$ -plectic vector field*.

(iii) If, moreover, there exists a p -form J with $\iota_v \omega = -\mathbf{d}J$ then v is called a *Hamiltonian vector field* and J is called a *Hamiltonian form* for v . The pair (v, J) is then a *Hamiltonian pair*.

(iv) If ω is non-degenerate in that the kernel of the contraction $\iota_{(-)}\omega : \text{Vect}(X) \rightarrow \Omega^{p+1}(X)$ vanishes, then v is uniquely determined by its current J ; in this case we call ω *$(p+1)$ -plectic* (instead of just *pre- $(p+1)$ -plectic*).

This is the case that is often restricted to in the literature, but the whole theory goes through in the pre-plectic case using such Hamiltonian pairs instead of just Hamiltonian forms.

Definition 2.3. (i) We write

$$\Omega_{\omega}^p(X) := \{(v, J) \in \text{Vect}(X) \oplus \Omega^p(X) \mid \iota_v \omega = -\mathbf{d}J\}$$

for the linear space of the above pairs and speak of *local Hamiltonian observables*.

(ii) We write

$$\text{Vec}_{\text{Ham}}(X, \omega) \hookrightarrow \text{Vect}_{\text{Symp}}(X, \omega) \hookrightarrow \text{Vect}(X)$$

for the subspaces of *symplectic* and *Hamiltonian vector fields*.

Notice that both inclusions above are sub-Lie algebras under the canonical Lie bracket of vector fields. When regarding $\theta = \mathbf{L}$ as a local Lagrangian, one then finds that the contraction $\iota \dots \omega$ vanishes on tangents to field trajectories which solve the corresponding equations of motion. Consequently, J_v is an on-shell conserved current, induced by the given symmetry. This is a special case of the general $(p + 1)$ -plectic Noether theorem [38].

Next, symmetries as above form a group in an evident way, where the composition operation is the pasting of diagrams

$$\begin{array}{ccccc}
 X & \xrightarrow{\phi_1} & X & \xrightarrow{\phi_2} & X \ . \\
 & \searrow \eta_1 & \downarrow \eta_2 & \swarrow \eta_1 & \\
 & \nabla & \mathbf{B}^{p+1} \text{U}(1)_{\text{conn}} & \nabla &
 \end{array} \tag{2.1}$$

Working out the Lie bracket of this Lie group one finds (as a special case of what is generally given by definition 2.6 below) that the bracket on these Hamiltonian pairs (v, Δ_v) (from definition 2.2) is given by

$$[(v_1, \Delta_{v_1}), (v_2, \Delta_{v_2})] = ([v_1, v_2], \mathcal{L}_{v_1} \Delta_{v_2} - \mathcal{L}_{v_2} \Delta_{v_1}) .$$

Remark 2.4. In the case that ∇ is given by a globally defined form θ and we think of this as being a local Lagrangian as before, so that the Δ s are its conserved Noether currents, then after passing to de Rham cohomology classes, the above bracket may be identified with what is known as the *Dickey bracket* on Noether currents [14] [8, (3.10)]. We generalize this to the case where ω is not assumed to be globally exact below in proposition 3.6.

Moreover, when a fixed potential $\Delta_{[v_1, v_2]}$ entering the definition of $J_{[v_1, v_2]}$ is picked from its equivalence class, then in terms of this the above bracket becomes

$$[(v_1, \Delta_{v_1}), (v_2, \Delta_{v_2})] = ([v_1, v_2], \Delta_{[v_1, v_2]}) + (0, \mathcal{L}_{v_1} \Delta_{v_2} - \mathcal{L}_{v_2} \Delta_{v_1} - \Delta_{[v_1, v_2]}) .$$

This may be identified with the bracket on Noether currents as displayed in [2, (13), (14)] for the case of super p -brane sigma-models. We turn to this example below in section 5.

Therefore, we have found traditional current algebras from a diagrammatic calculus of symmetries of higher connections.

Remark 2.5. This highlights two points which seem not to have been explicitly addressed in traditional literature:

1. *Higher groups*: when $p > 0$, the group of symmetries is a higher group (a higher group stack) of higher order symmetries-of-symmetries. Namely the homotopies η in the diagram (2.1) may themselves have homotopies-of-homotopies between them

$$\begin{array}{ccc}
 X & \xrightarrow{\phi} & X \\
 & \searrow \eta & \swarrow \eta \\
 & \nabla & \mathbf{B}^{p+1} \text{U}(1)_{\text{conn}}
 \end{array}$$

(Note: The diagram above shows a curved arrow η between the two ∇ arrows, indicating a homotopy between the two paths from X to $\mathbf{B}^{p+1} \text{U}(1)_{\text{conn}}$.)

corresponding to gauge-of-gauge transformations. Hence after Lie differentiation then, on top of the Lie bracket of conserved currents above, there are higher order gauge transformations between these currents. These may be most directly understood from the fact that the choice of Δ_v above is clearly only unique up to addition of exact terms, whose potentials in turn are themselves only unique up to exact terms, and so forth. As a result, we find not just a Lie algebra, but a dg-Lie algebra of currents, whose differential is the de Rham differential acting on higher order current forms.

2. *Refined cohomology and higher forms*: in full generality, the above discussion needs to be performed not just for globally defined θ , but for higher prequantizations $\bar{\theta}$ which are given by Čech-Deligne cocycles with curvature $(d+1)$ -form ω . This allows for the inclusion of geometric data via differential cohomology.

Incorporating these, the resulting dg-Lie algebra is the one given in [17, definition/proposition 4.2.1]:

Let X be a smooth manifold, $\omega \in \Omega_{\text{cl}}^{p+2}(X)$ a closed differential $(p+2)$ -form, with (X, ω) regarded as a pre- $(p+1)$ -plectic manifold. Let $\bar{\theta}$ be a higher pre-quantization of ω given by a Čech-Deligne cocycle with respect to a cover \mathcal{U} of X . Write $\text{Tot}^\bullet(\mathcal{U}, \Omega^\bullet)$ for the Čech-de Rham complex of the cover and notice that the Lie derivatives of $\bar{\theta}$ are naturally regarded as taking values in this complex. Then we have:

Definition 2.6. The *Poisson bracket dg-Lie algebra*

$$\mathfrak{Pois}_{\text{dg}}(X, \bar{\theta}) \in \text{dgLieAlg} \hookrightarrow L_\infty\text{Alg}$$

is the dg-Lie algebra whose underlying chain complex has components

$$\begin{aligned} \mathfrak{Pois}_{\text{dg}}(X, \bar{\theta})^0 &:= \{(v, \bar{\Delta}) \in \text{Vect}(X) \oplus \text{Tot}^p(\mathcal{U}, \Omega^\bullet) \mid \mathcal{L}_v \bar{\theta} = \mathbf{d}_{\text{Tot}} \bar{\Delta}\} , \\ \mathfrak{Pois}_{\text{dg}}(X, \bar{\theta})^{i \geq 1} &:= \text{Tot}^{p-i}(\mathcal{U}, \Omega^\bullet) , \end{aligned}$$

with differential \mathbf{d}_{Tot} , and whose non-vanishing Lie brackets are

$$\begin{aligned} [(v_1, \bar{\Delta}_1), (v_2, \bar{\Delta}_2)] &= ([v_1, v_2], \mathcal{L}_{v_1} \bar{\Delta}_2 - \mathcal{L}_{v_2} \bar{\Delta}_1) , \\ [(v, \bar{\Delta}), \bar{\eta}] &= -[\bar{\eta}, (v, \bar{\Delta})] = \mathcal{L}_v \bar{\eta} . \end{aligned}$$

It turns out that there is a very different looking but equivalent incarnation of this L_∞ -algebra, originally considered in [32]:

Definition 2.7 (Higher Poisson bracket of local observables). Given a pre- $(p+1)$ -plectic manifold (X, ω) ,

- (i) we say that the truncated de Rham complex ending in the Hamiltonian pairs of definition 2.3 is the *complex of local observables* of (X, ω) , denoted

$$\Omega_\omega^\bullet(X) := \left(C^\infty(X) \xrightarrow{\mathbf{d}} \Omega^1(X) \xrightarrow{\mathbf{d}} \dots \xrightarrow{\mathbf{d}} \Omega^{n-2}(X) \xrightarrow{(0, \mathbf{d})} \Omega_\omega^p(X) \right) ,$$

with $\Omega_\omega^p(X)$ in degree zero.

(ii) The *binary higher Poisson bracket* on local Hamiltonian observables is the linear map

$$\{-, -\} : \Omega_\omega^p(X) \otimes \Omega_\omega^p(X) \longrightarrow \Omega_\omega^p(X)$$

given by the formula

$$[(v_1, J_1), (v_2, J_1)] := [[v_1, v_2], \iota_{v_1 \wedge v_2} \omega] ; \tag{2.2}$$

(iii) and for $k \geq 3$ the *k-ary higher Poisson bracket* is the linear map

$$\{-, \dots, -\} : (\Omega_\omega^p(X))^{\otimes k} \longrightarrow \Omega^{p+2-k}(X)$$

given by the formula

$$[(v_1, J_1), \dots, (v_k, J_k)] := (-1)^{\lfloor \frac{k-1}{2} \rfloor} \iota_{v_1 \wedge \dots \wedge v_k} \omega .$$

(iv) The chain complex of local observables equipped with these linear maps for all k will be called the *higher Poisson bracket L_∞ algebra* of (X, ω) , denoted

$$\mathfrak{Pois}_\infty(X, \omega) := (\Omega_\omega^\bullet(X), \mathbf{d}, \{-, -\}, \{-, -, -\}, \dots) .$$

Remark 2.8. From [17, theorem 4.2.2], there is an equivalence of L_∞ -algebras

$$\mathfrak{Pois}_\infty(X, \omega) \xrightarrow{\simeq} \mathfrak{Pois}_{\text{dg}}(X, \omega)$$

between those of definition 2.6 and definition 2.7.

We would like to gain some further understanding of these L_∞ -algebras. The idea is that starting with \mathbb{R} , we can view it as a chain complex by placing it in various degrees with the remaining degrees being zero. Shifting the degrees up and down result in stack analogs of the classifying functor \mathbf{B} . When the resulting stack is flat, we denote this with a musical \flat .

Definition 2.9. For X a smooth manifold, denote

1. by $\mathbf{H}(X, \flat \mathbf{B}^p \mathbb{R})$ the abelian Lie $(p+1)$ -algebra given by the chain complex

$$\Omega^0(X) \xrightarrow{\mathbf{d}} \Omega^1(X) \xrightarrow{\mathbf{d}} \dots \xrightarrow{\mathbf{d}} \Omega^p(X) ,$$

with $\Omega^p(X)$ in degree 0;

2. by $\mathbf{BH}(X, \flat \mathbf{B}^p \mathbb{R})$ the abelian Lie $(p+1)$ -algebra given by the chain complex, including one more term,

$$\Omega^0(X) \xrightarrow{\mathbf{d}} \Omega^1(X) \xrightarrow{\mathbf{d}} \dots \xrightarrow{\mathbf{d}} \Omega^p(X) \xrightarrow{\mathbf{d}} \mathbf{d}\Omega^p(X) ,$$

with $\mathbf{d}\Omega^p(X)$ in degree zero.

We start with relating the latter to Hamiltonian vector fields.

Remark 2.10. From [17, proposition 3.2.3] there is the following result. For (X, ω) a pre- $(p + 1)$ -plectic manifold, the multilinear maps

$$\begin{aligned} \omega_{[1]} &: v \mapsto -\iota_v \omega; \\ \omega_{[2]} &: v_1 \wedge v_2 \mapsto \iota_{v_1 \wedge v_2} \omega; \\ &\vdots \\ \omega_{[p+2]} &: v_1 \wedge v_2 \wedge \cdots \wedge v_{p+2} \mapsto -(-1)^{\binom{p+2}{2}} \iota_{v_1 \wedge v_2 \wedge \cdots \wedge v_{p+2}} \omega \end{aligned}$$

define an L_∞ -morphism $\omega_{[\bullet]} : \text{Vect}_{\text{Ham}}(X) \rightarrow \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R})$. This is the higher Kirillov-Kostant-Souriau L_∞ -algebra $(p + 2)$ -cocycle on the Lie algebra of Hamiltonian vector fields with values in the abelian $(p + 2)$ -algebra $\mathbf{BH}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R})$ associated to the pre- $(p + 1)$ -plectic manifold (X, ω) .

The L_∞ -algebras in remark 2.8 are characterized as follows.

Remark 2.11. From [17, theorem 3.3.1], the Poisson bracket Lie $(p + 1)$ -algebra $\mathfrak{Pois}_\infty(X, \omega)$ is the extension of the Lie algebra of Hamiltonian vector fields, definition 2.2, by the abelian Lie $(p + 1)$ -algebra $\mathbf{H}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R})$ induced by the Kirillov-Kostant-Souriau L_∞ -algebra $(p + 2)$ -cocycle $\omega_{[\bullet]}$. Namely, there is a homotopy fiber sequence of L_∞ -algebras of the form

$$\begin{array}{ccc} \mathbf{H}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R}) & \longrightarrow & \mathfrak{Pois}_\infty(X, \omega) \\ & & \downarrow \\ & & \text{Vect}_{\text{Ham}}(X, \omega) \xrightarrow{\omega_{[\bullet]}} \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R}) . \end{array}$$

Here the vertical homomorphism comes from the projection $\Omega_\omega^p(X) \rightarrow \text{Vect}_{\text{Ham}}(X)$ (from Hamiltonian pairs to Hamiltonian vector fields, definition 2.3) which induces a surjective linear morphism of L_∞ -algebras.

3 Truncation to ordinary Lie algebras

We would like to consider appropriate truncations down to more classical structures. This is governed by controlling the ‘size’ of the corresponding chain complex. Given any homotopy-theoretic higher structure, hence a homotopy n -type, there its ‘truncation’ down to a 0-type, obtained by quotienting out all symmetries. This operation in general forgets much relevant structure, but the resulting object is sometimes more easily recognized and compared to more traditional concepts.

For the simple special case of chain complexes of vector spaces in non-negative degree (in the homological degree conventions), 0-truncation is the functor

$$\tau_{\leq 0} : \text{Ch}_\bullet \longrightarrow \text{Vect}$$

which sends a chain complex $(\cdots V_2 \xrightarrow{\partial_1} V_1 \xrightarrow{\partial_0} V_0)$ to its degree-0 homology group

$$\tau_{\leq 0} V_\bullet = V_0 / \text{im}(\partial_0) = H_0(V_\bullet) .$$

We now discuss this 0-truncation for the Lie $(p + 1)$ -algebras of conserved currents discussed in the previous section, section 2.

Lemma 3.1. *The 0-truncation functor $\tau_{\leq 0}$ on chain complexes induces a functor*

$$\tau_{\leq 0}: L_{\infty}\text{-alg}_{\geq 0} \rightarrow \text{Lie}$$

where $L_{\infty}\text{-alg}_{\geq 0}$ denotes the category of L_{∞} -algebras concentrated in nonnegative degrees, with L_{∞} -morphisms as morphisms, and Lie denotes the category of Lie algebras with Lie algebra morphisms. More explicitly, $\tau_{\leq 0}$ maps an L_{∞} -algebra $(\mathfrak{g}, \mathbf{d}, \{-, -\}, \{-, -, -\}, \dots)$ concentrated in nonpositive degrees to the Lie algebra $(H_0(\mathfrak{g}), \{-, -\})$ and an L_{∞} -morphism $f: \mathfrak{g} \rightarrow \mathfrak{h}$ to the Lie algebra morphism $H_0(f_1): H_0(\mathfrak{g}) \rightarrow H_0(\mathfrak{h})$, where f_1 is the linear component of f . Moreover, the natural morphism of chain complexes $\mathfrak{g} \rightarrow \tau_{\leq 0}\mathfrak{g}$ is a natural linear and surjective L_{∞} -morphism

$$(\mathfrak{g}, \mathbf{d}, \{-, -\}, \{-, -, -\}, \dots) \rightarrow (H_0(\mathfrak{g}), \{-, -\}). \quad (3.1)$$

Proof. Since the chain complex \mathfrak{g} is concentrated in nonnegative degrees, the chain complex $\tau_{\leq 0}\mathfrak{g}$ consists of the vector space $H_0(\mathfrak{g})$ concentrated in degree zero. All the statements in the lemma then follow straightforwardly from this.

Remark 3.2. Naturality of the linear L_{∞} -morphism (3.1) means that for any L_{∞} -morphism $f: \mathfrak{g} \rightarrow \mathfrak{h}$ between L_{∞} -algebras concentrated in nonnegative degree we have a commutative diagram of L_{∞} -morphisms

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{f} & \mathfrak{h} \\ \downarrow & & \downarrow \\ H_0(\mathfrak{g}) & \xrightarrow{H_0(f_1)} & H_0(\mathfrak{h}) \end{array}.$$

We now look at the 0-truncation of the corresponding Poisson algebras.

Definition 3.3. Let (X, ω) be a pre- $(p + 1)$ -plectic manifold. Write

$$\mathfrak{Pois}(X, \omega) := \tau_{\leq 0}\mathfrak{Pois}_{\infty}(X, \omega)$$

for the 0-truncation of $\mathfrak{Pois}_{\infty}(X, \omega) \simeq \mathfrak{Pois}_{\text{dg}}(X, \omega)$.

Note that using definition 2.6, definition 2.7, remark 2.8, and via lemma 3.1, this is the quotient $\mathfrak{Pois}_{\infty}(X, \omega) \rightarrow H_0\mathfrak{Pois}_{\infty}(X, \omega)$ by exact current p -forms. We now characterize this 0-truncation.

Proposition 3.4. *Let (X, ω) be a pre- $(p + 1)$ -plectic manifold. Then the Lie 1-algebra of currents, definition 3.3, is a central extension of the Hamiltonian vector fields by the abelian Lie algebra $H_{\text{dR}}^p(X)$ of de Rham p -forms. In other words, writing $\mathfrak{Pois}(X, \omega)$ for $H_0\mathfrak{Pois}_{\infty}(X, \omega)$, there is a short exact sequence of Lie algebras*

$$0 \rightarrow H_{\text{dR}}^p(X) \longrightarrow \mathfrak{Pois}(X, \omega) \longrightarrow \text{Vect}_{\text{Ham}}(X, \omega) \rightarrow 0,$$

and $H_{\text{dR}}^p(X)$ is central in $\mathfrak{Pois}(X, \omega)$.

Proof. From the short exact sequence of chain complexes given by proposition 2.11

$$0 \rightarrow \mathbf{H}(X, \mathfrak{b}\mathbf{B}^p\mathbb{R}) \rightarrow \mathfrak{Pois}_\infty(X, \omega) \rightarrow \text{Vect}_{\text{Ham}}(X, \omega) \rightarrow 0,$$

we get the long exact sequence in homology

$$0 \rightarrow H_{\text{dR}}^p(X) \rightarrow \mathfrak{Pois}(X, \omega) \rightarrow \text{Vect}_{\text{Ham}}(X, \omega) \rightarrow 0$$

and, by lemma 3.1, this is a short exact sequence of Lie algebras. The fact that $H_{\text{dR}}^p(X)$ is central in $\mathfrak{Pois}(X, \omega)$ is immediate from equation (2.2).

Remark 3.5. For the special case when X is the jet bundle of a field bundle, when the de Rham differentials appearing everywhere are constrained to be the corresponding horizontal differentials, and under the assumption that the cohomology of this horizontal de Rham complex is concentrated in degree 0, then another L_∞ -structure on the truncated de Rham complex appearing in definition 2.7 has been constructed in [7]. Inspection of the construction around theorem 7 there shows that this L_∞ -algebra is L_∞ -equivalent to its 0-truncation $\mathfrak{Pois}(X, \omega)$ which under these assumptions is the Dickey algebra [14]; see the comments above in section 2. Hence for all purposes of homotopy theory the algebra in [7] is this 0-truncation.

In order to further compare proposition 3.4 to existing literature, we consider now a choice of Hamiltonian current forms for each symmetry generator. The following statement is for definition 2.7 what remark 2.4 was for definition 2.6.

Proposition 3.6. *Under a choice of linear splitting $J: v \mapsto (v, J_v)$ of the natural projection $\Omega_\omega^p(X) \rightarrow \text{Vect}_{\text{Ham}}(X, \omega)$ from definition 2.2, the Lie bracket on de Rham cohomology classes of currents in $\mathfrak{Pois}(X, \omega)$, proposition 3.4, is isomorphic to*

$$\{v + [\alpha], w + [\beta]\} = [v, w] + \left[J_{[v, w]} - \frac{1}{2}(\mathcal{L}_v J_w - \mathcal{L}_w J_v) \right].$$

When moreover ω is globally exact via $\mathbf{d}\theta = \omega$, then in terms of the currents $\Delta_v := i_v\theta - J_v$ this reduces to

$$\{v + [\alpha], w + [\beta]\} = [v, w] + [\mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v - \Delta_{[v, w]}]$$

as in remark 2.4.

Proof. The splitting induces a linear isomorphism of the form

$$\begin{aligned} \text{Vect}_{\text{Ham}}(X, \omega) \oplus H_{\text{dR}}^p(X) &\xrightarrow{\sim} \mathfrak{Pois}(X, \omega) \\ v + [\alpha] &\mapsto (v, J_v - \alpha) \end{aligned}$$

and with this the Lie bracket on $\text{Vect}_{\text{Ham}}(X, \omega) \oplus H_{\text{dR}}^p(X)$ reads

$$\{v + [\alpha], w + [\beta]\} = [v, w] + [J_{[v, w]} - \iota_v \wedge w].$$

Since $\iota_v \omega = -\mathbf{d}J_v$, one has

$$\iota_v \wedge \omega = -\iota_v \mathbf{d}J_v = -\mathcal{L}_v J_v + \mathbf{d}\iota_v J_v$$

and

$$\iota_v \wedge \omega = -\iota_w \wedge \omega = \iota_v \mathbf{d}J_w = \mathcal{L}_v J_w - \mathbf{d}\iota_v J_w .$$

Hence, combining, we can write

$$\iota_v \wedge \omega = \frac{1}{2}(\mathcal{L}_v J_w - \mathcal{L}_w J_v) - \frac{1}{2}\mathbf{d}(\iota_v J_w - \iota_w J_v)$$

and so

$$\{v + [\alpha], w + [\beta]\} = [v, w] + \left[J_{[v,w]} - \frac{1}{2}(\mathcal{L}_v J_w - \mathcal{L}_w J_v) \right] .$$

If, moreover, a global potential θ for the pre- n -plectic form ω is given, i.e., if one has a p -form θ with $\mathbf{d}\theta = \omega$ then, writing $J_v = i_v \theta - \Delta_v$, one finds

$$\begin{aligned} J_{[v,w]} - \frac{1}{2}(\mathcal{L}_v J_w - \mathcal{L}_w J_v) &= -\Delta_{[v,w]} + \iota_{[v,w]}\theta - \frac{1}{2}(\mathcal{L}_v \iota_w \theta - \mathcal{L}_w \iota_v \theta) + \frac{1}{2}(\mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v) \\ &= -\Delta_{[v,w]} + \underbrace{\iota_{[v,w]}\theta - \frac{1}{2}(\iota_{[v,w]}\theta - \iota_{[w,v]}\theta)}_{=0} - \frac{1}{2}(\iota_w \mathcal{L}_v \theta - \iota_v \mathcal{L}_w \theta) \\ &\quad + \frac{1}{2}(\mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v) \\ &= -\Delta_{[v,w]} - \frac{1}{2}(\iota_w \mathbf{d}\Delta_v - \iota_v \mathbf{d}\Delta_w) + \frac{1}{2}(\mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v) \\ &= -\Delta_{[v,w]} + \mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v + \mathbf{d}\text{-exact terms} . \end{aligned}$$

Therefore, finally, we get the desired expression

$$\{v + [\alpha], w + [\beta]\} = [v, w] + [\mathcal{L}_v \Delta_w - \mathcal{L}_w \Delta_v - \Delta_{[v,w]}] .$$

4 Transgression to ordinary Lie algebras

We will now consider another process of getting ordinary Lie algebras starting from our higher algebras. While the 0-truncation which we considered in section 3 just quotients out and forgets higher gauge transformations to obtain a plain Lie 1-algebra of currents, there is another way to turn a higher current Lie $(p+1)$ -algebra into a Lie k -algebra for $k \leq p$ that retains more information: mathematically this is *transgression*, while in terms of physics this corresponds to the natural operation of integrating local currents over submanifolds of some codimension. A notable instance is to do so over Cauchy surfaces, in order to turn them into ordinary observables (i.e. functions on field configurations).

The conception of higher groups of currents as indicated in section 2, via automorphisms sliced over a higher differential moduli stack, lends itself to a clear picture of the process of transgression of higher currents as follows.

First observe that transgression of plain differential forms has the following neat sheaf-theoretic formulation. Write Ω^{p+1} for the sheaf of smooth differential $(p+1)$ -forms (on the

site of all smooth manifolds). Then, by the Yoneda lemma, a smooth differential $(p + 1)$ -form $\omega \in \Omega^{p+1}(X)$ on some smooth manifold X is equivalently a morphism in the category of sheaves of the form

$$\omega : X \longrightarrow \Omega^{p+1}.$$

Now, given a closed oriented smooth manifold Σ of dimension $k \leq p + 1$, write $[\Sigma, \Omega^{p+1}]$ for the mapping space, which is the sheaf that sends a test manifold to the set of sheaf morphisms $U \times \Sigma \rightarrow \Omega^{p+1}$, and hence to the set of smooth differential $(p + 1)$ -forms on $U \times \Sigma$. It is immediate then that ordinary fiber integration of differential forms along the projections $\Sigma \times U \rightarrow U$, for all possible U , is embodied in a single morphism of sheaves of the form

$$\int_{\Sigma} : [\Sigma, \Omega^{p+1}] \longrightarrow \Omega^{p+1-k}.$$

Now, forming the mapping sheaf out of Σ is a functorial process, and hence we obtain from the above ingredients the following

Proposition 4.1. *There a composite morphism of the form*

$$\int_{\Sigma} [\Sigma, \omega] : [\Sigma, X] \xrightarrow{[\Sigma, \omega]} [\Sigma, \Omega^{p+1}] \xrightarrow{\int_{\Sigma}} \Omega^{p+1-k},$$

representing a differential $(p + 1 - k)$ -form on the smooth mapping space $[\Sigma, X]$.

The point now is that, unwinding all the definitions, one finds

Corollary 4.2. *The differential $(p + 1 - k)$ -form $\int_{\Sigma} [\Sigma, \omega]$ is the transgression of ω to the above mapping space, namely is the form $\int_{\Sigma} \text{ev}_{\Sigma}^* \omega$, for*

$$\text{ev}_{\Sigma} : [\Sigma, X] \times \Sigma \longrightarrow X$$

the evaluation map.

While this is the traditional formulation of transgression, the above re-formulation via mapping sheaves has the advantage that it neatly generalizes from differential n -forms ω to their pre-quantization by $(p + 1)$ -connections ∇ , that we considered in section 2.

Recall that fiber integration of differential forms, viewed as an operation on all possible $\Sigma \times U$ as U varies, can be encapsulated in a single morphism of sheaves as above. Similarly, fiber integration of n -connections, hence fiber integration of cocycles in ordinary differential cohomology of degree $(p + 2)$, is equivalently encoded [18] in a morphism of smooth higher stacks of the form

$$\int_{\Sigma} : [\Sigma, \mathbf{B}^{p+1} \text{U}(1)_{\text{conn}}] \longrightarrow \mathbf{B}^{p+1-k} \text{U}(1)_{\text{conn}}.$$

With this in hand we immediately obtain

Proposition 4.3. (i) *There is a natural homomorphism from the smooth $(p + 1)$ -group of conserved currents as in section 2, to $(p + 1 - k)$ -group of their transgression to the mapping space out of Σ .*

- (ii) This is done simply by applying the ∞ -functor $[\Sigma, -]$ to the defining slice diagrams and then postcomposing the slicing with \int_X , as indicated here:

$$\left(\begin{array}{ccc} X & \xrightarrow{\phi} & X \\ & \searrow \eta \simeq & \swarrow \eta \\ & \nabla & \nabla \\ & \mathbf{B}^{p+1} \mathbf{U}(1)_{\text{conn}} & \end{array} \right) \mapsto \left(\begin{array}{ccc} [\Sigma, X] & \xrightarrow{[\Sigma, \phi]} & [\Sigma, X] \\ & \searrow [\Sigma, \eta] \simeq & \swarrow [\Sigma, \eta] \\ & [\Sigma, \nabla] & [\Sigma, \nabla] \\ & [\Sigma, \mathbf{B}^{p+1} \mathbf{U}(1)_{\text{conn}}] & \\ & \downarrow \int_{\Sigma} & \\ & \mathbf{B}^{p+1-k} \mathbf{U}(1)_{\text{conn}} & \end{array} \right).$$

Remark 4.4. The above operation now transgresses not only the $(p+1)$ -connection ∇ , which for us is the Lagrangian of the given field theory, but transgresses also all the currents which are embodied in the equivalences labeled η . Specifically, if ∇ happens to be given by a globally defined differential $(p+1)$ -form $\theta = \mathbf{L}$ then η is, infinitesimally, given by a current p -form J , and the above operation takes that to its transgression form $\int_{\Sigma} \text{ev}^* J$.

1. When $k = p$, then this transgressed form is a 0-form, hence a function, and has the interpretation of the actual observable induced by the current J .
2. Generally, for $k = p$ then on the right all higher homotopies vanish, due to the composition with \int_{Σ} which lands in the 1-stack $\mathbf{B} \mathbf{U}(1)_{\text{conn}}$, and hence in this case the truncation map factors through the 0-truncation of the group on the left, which we just discussed above. This way the 0-truncation of the current Lie $(p+1)$ -algebra maps into the generalized affine Lie algebra of currents, but the latter may in general be bigger.

Example 4.5. The archetypical example of this process is given by the case of the 3d WZW model on a compact Lie group G . In this case ∇ above is the WZW gerbe $\mathbf{L}_{\text{WZW}} : G \rightarrow \mathbf{B}^2 \mathbf{U}(1)_{\text{conn}}$ on the group, and its transgression $\int_{S^1} [S^1, \mathbf{L}_{\text{WZW}}] : [S^1, G] \rightarrow \mathbf{B} \mathbf{U}(1)_{\text{conn}}$ is a bundle with connection on the loop group. By the results of [16] the group of currents induced by this, as discussed above, is the “quantomorphism group” of this bundle regarded as a prequantum line bundle in the sense of Souriau. It is a famous fact [30] (see [37, around proposition 42]), that this is the Kac-Moody central extension of the loop group of G , whose Lie algebra is the corresponding affine Lie algebra of G . This is “the” current algebra as seen in most of the mathematics literature.

We see here that this Kac-Moody loop group is nothing but the transgression of the smooth 2-group of currents of the WZW model, which, by [16, 2.6.1], is what is called the String 2-group of G , and that the affine current Lie algebra is nothing but the transgression of the corresponding Lie 2-algebra, which is the *string* Lie 2-algebra [6].

Remark 4.6. Two variants of transgression are of relevance in physics:

1. When the target space is a product $X \times \Sigma$, there is a canonical morphism $X \rightarrow [\Sigma, X \times \Sigma]$. Transgression followed by precomposition with this inclusion

$$\left(X \times \Sigma \xrightarrow{\nabla} \mathbf{B}^{p+1} \mathbf{U}(1)_{\text{conn}} \right) \mapsto \left(X \rightarrow [\Sigma, X \times \Sigma] \xrightarrow{\int_{\Sigma} [\Sigma, \nabla]} \mathbf{B}^{p+1-\dim(\Sigma)} \mathbf{U}(1)_{\text{conn}} \right)$$

is *double dimensional reduction*.

2. Moreover, when Σ is equipped with a base point $s_0 : * \rightarrow \Sigma$, then there is also the induced restriction operation

$$\left(X \times \Sigma \xrightarrow{\nabla} \mathbf{B}^{p+1} U(1)_{\text{conn}} \right) \mapsto \left(X \rightarrow [\Sigma, X \times \Sigma] \xrightarrow{!s_0} X \times \Sigma \xrightarrow{\nabla} \mathbf{B}^{p+1} U(1)_{\text{conn}} \right).$$

Example 4.7. The product case occurs frequently in the M-theory circle bundle over 10-dimensional spacetime X . The combination of double dimensional reduction with restriction is, for instance, what takes the 11-dimensional supergravity C-field to the B-field and Ramonf-Ramond (RR) form in 10 dimensions. The above process allows us to do so at the level of stacks, where all the higher geometric and gauge structures are carried along. A stack perspective on this can be found in [19].

5 Application to brane charges

We now discuss how the above constructions lead to interesting applications in (supersymmetric) physical models. We note that all the previous discussion generalizes directly to the situation of superalgebra and supergeometry, where all manifolds are generalized to supermanifolds, differential forms to super-differential forms, and where Lie n -algebras are generalized to super Lie n -algebras (see [21] for details and pointers to the literature). This means that we may apply the previous results to supermanifolds X as they appear in higher supergravity theory, where they carry certain special super $(p + 2)$ -forms which serve as the curvatures of WZW terms for the super p -brane sigma-models of Green-Schwarz type, with target space X .

We consider this in detail. For N a real spin representation of $\text{Spin}(d - 1, 1)$, let $\mathbb{R}^{d-1,1|N}$ be the corresponding super-Minkowski spacetime, regarded as a super translation Lie group. The canonical left-invariant 1-form on $\mathbb{R}^{d-1,1|N}$ is traditionally denoted $E = (E^a, \psi^\alpha)_{a,\alpha}$, where a ranges over linear basis elements for \mathbb{R}^d and α runs over a basis for the linear representation space N . To make precise statements, let then (d, N, p) an item in the old brane scan, i.e. let $p \in \mathbb{N}$ be such that

$$\omega_{\text{WZW}} := \bar{\psi} \wedge \Gamma_{a_1 \dots a_p} \psi \wedge E^{a_1} \wedge \dots \wedge E^{a_p} \in \Omega_{\text{cl}}^{p+2}(\mathbb{R}^{d-1,1|N})$$

is a super Lie algebra cocycle of the given form (see [21] for review and details). This is then the WZW-curvature term for the Green-Schwarz sigma model for super p -branes propagating on $\mathbb{R}^{d-1,1|N}$.

For instance for $d = 10$, $p = 1$ and $N = (2, 0) = \mathbf{16} + \mathbf{16}$ there is such a cocycle, and it constitutes the curvature 3-form of the κ -symmetry WZW-term of the type IIB superstring on Minkowski background. More generally, let X be a more general supermanifold locally modeled on $\mathbb{R}^{d-1,1|N}$ and equipped with a super vielbein field (super soldering form) E and further relevant fields, including a super $(p + 2)$ -form ω . Then the equations of motion of d -dimensional N -supersymmetric supergravity constrain the vielbein to be such that the fermionic part ω_{WZW}^X of ω^X is the pullback of ω_{WZW} via E [9, 10].

Following terminology familiar from the theory of G_2 -manifolds, we may express this by saying that ω_{WZW}^X is a *definite form*, define on ω_{WZW} . We may regard the pair $(X, \omega_{\text{WZW}}^X)$ as

a pre- $(p+1)$ -plectic supermanifold according to definition 2.2. As such it induces the higher Poisson bracket super Lie $(p+1)$ -algebra $\mathfrak{Pois}(X, \omega_{\text{WZW}}^X)$ of definition 2.6, definition 2.7, and remark 2.8.

The fact that the equations of motion of supergravity force ω_{WZW}^X to be definite on ω_{WZW} means that isometries of X preserve ω_{WZW}^X , hence that their ‘super’ vector fields (the Killing vectors and Killing spinors) are $(p+1)$ -plectic vector fields in the sense of definition 2.2. It is not guaranteed that every isometry with respect to the given vielbein preserves θ up to gauge transformation; those that do are precisely the *Hamiltonian isometries* in the sense of definition 2.2. Write

$$\mathfrak{I}so\text{m}_{\text{Ham}}(X, \omega_{\text{WZW}}^X) \hookrightarrow \text{Vect}_{\text{Ham}}(X, \omega_{\text{WZW}}^X)$$

for the inclusion of these into all Hamiltonian vector fields

Definition 5.1. Write $\mathfrak{B}\mathfrak{P}\mathfrak{S}(X, \omega_{\text{WZW}}^X)$ for the restriction of the current Lie $(p+1)$ -algebra $\mathfrak{Pois}(X, \omega_{\text{WZW}}^X)$ (definition 2.6, definition 2.7, remark 2.8) of ω_{WZW}^X to isometries, i.e. for the super L_∞ -algebra in the homotopy pullback diagram

$$\begin{array}{ccc} \mathfrak{B}\mathfrak{P}\mathfrak{S}(X, \omega_{\text{WZW}}^X) & \longrightarrow & \mathfrak{Pois}(X, \omega_{\text{WZW}}^X) \\ \downarrow & & \downarrow \\ \mathfrak{I}so\text{m}(X, \omega_{\text{WZW}}^X) & \longrightarrow & \text{Vect}_{\text{Ham}}(X, \omega_{\text{WZW}}^X) . \end{array}$$

The following reproduces and generalizes the result in [2, p. 8].

Proposition 5.2. *The 0-truncation to a super-Lie algebra $\tau_0\mathfrak{B}\mathfrak{P}\mathfrak{S}(X, \omega)$ is the central extension of the supersymmetry algebra of X by charges of p -branes wrapping non-trivial cycles.*

Proof. This follows via remark 3.6 by proposition 3.4, which gives an extension

$$H_{\text{dR}}^p(X) \rightarrow \tau_0\mathfrak{B}\mathfrak{P}\mathfrak{S}(X, \omega_{\text{WZW}}^X) \rightarrow \mathfrak{I}so\text{m}(X, \omega_{\text{WZW}}^X) ,$$

classified by $\omega_{\text{WZW}}^X(-, -)$. The elements in $H_{\text{dR}}^p(X)$ are the p -brane charges.

Hence we have the following picture:

$$\left\{ \begin{array}{ccc} X & \xrightarrow{\phi} & X \\ & \searrow \eta \simeq & \swarrow \\ & \mathbf{B}^{p+1} & \mathbf{U}(1)_{\text{conn}} \\ & \swarrow \mathbf{L}_{\text{WZW}} & \searrow \mathbf{L}_{\text{WZW}} \end{array} \right\} , \quad \omega_{\text{WZW}}^X = F_{\mathbf{L}_{\text{WZW}}} \quad \left\{ \begin{array}{c} \xrightarrow{\text{Lie}} \\ \tau_0\mathfrak{Pois}(X, \omega_{\text{WZW}}^X) \\ \downarrow \\ \mathfrak{I}so\text{m}(X, \omega_{\text{WZW}}^X) \end{array} \right\} .$$

These are the supersymmetry extensions induced by a single brane species, as considered originally in [2]. But the ‘‘type II algebra’’ and the ‘‘M-theory algebra’’ [24, 41] are

supposed to arise from considering not just strings and membranes, but also the branes on which these may end, namely the D-branes and the M5-brane, respectively. Notably the M-theory supersymmetry algebra is given on the fermionic generators in traditional local component notation as [4, 41]

$$\{Q_\alpha, Q_\beta\} = (C\Gamma^M)_{\alpha\beta}P_M + (C\Gamma_{MN})_{\alpha\beta}Z_2^{MN} + (C\Gamma_{MNPQR})_{\alpha\beta}Z_5^{MNPQR}, \quad (5.1)$$

where C is the charge conjugation matrix and Γ are the Dirac matrices. The three terms on the right hand side correspond to the graviton momentum 1-form charge, the membrane 2-form charge and the fivebrane 5-form charge. Notice again that this traditional expression applies only locally, on patches of spacetime diffeomorphic to super-Minkowski spacetime.

Now the proper global analysis of [2, p. 8] and of proposition 5.2 only ever produces extensions by charges of a single brane species with no (higher) gauge fields on its worldvolume. But in [21, 22, 39] we explained that the WZW-type sigma models for super p_2 -branes with (higher) gauge fields on the their worldvolume and on which super p_1 -branes may end, are globally defined not on target superspacetime X itself, but on the total space \tilde{X} of a super p_1 -stack extension $\tilde{X} \rightarrow X$ of superspacetime, which itself is a differential refinement of the p_1 -gerbe \hat{X} that underlies the WZW term of the p_1 -branes. Moreover, in [16] we showed that the higher Heisenberg-Kostant-Souriau extensions of remark 2.11 generalizes to such higher stacky base spaces. Schematically this follows now by the following picture, the full details are in [38, 39]:

$$\left(\begin{array}{ccc} \hat{X} & \xrightarrow{\cong} & \hat{X} \\ \downarrow & & \downarrow \\ X & \xrightarrow{\cong} & X \\ \swarrow \mathbf{L}_{\text{WZW}}^{p_1} & & \swarrow \mathbf{L}_{\text{WZW}}^{p_1} \\ & \mathbf{B}^{p_1+1}U(1)_{\text{conn}} & \\ \downarrow & & \downarrow \\ & \mathbf{B}^{p_2+1}U(1)_{\text{conn}} & \end{array} \right) \xrightarrow{\cong} \begin{array}{c} H^{p_2}(\tilde{X}) \\ \downarrow \\ \tau_0\mathfrak{Pois}(\tilde{X}, \mathbf{L}_{\text{WZW}}^{p_2}) \\ \downarrow \\ \mathfrak{Isom}(\tilde{X}, \mathbf{L}_{\text{WZW}}^{p_2}) \end{array}$$

(All 2-cells on the left are filled by homotopies of higher stacks. We suppress them notationally just for convenience and readability.)

Here the differential refinement \tilde{X} of \hat{X} is what makes a sigma-model field $\Sigma_{p_2} \rightarrow \tilde{X}$ be a pair consisting of an ordinary map to target spacetime $\Sigma_{p_2} \rightarrow X$ together with a twisted p_1 -form gauge field on Σ_{p_2} . This is the global model for super p -branes with tensor multiplet higher gauge fields on their worldvolume.

While this is necessary for the full picture, the isometry group and the cohomology of \tilde{X} is hard to compute. There is however a canonical forgetful map $\tilde{X} \rightarrow \hat{X}$ to the geometric realization of this differential stack (regarded itself as a locally constant stack, see [38] for details). By combining results of [38] and [29], one finds that in the case at hand \tilde{X} is the homotopy type of the $K(\mathbb{Z}, p_1 + 1)$ -fiber bundle over spacetime X that is classified by the

integral class of the background field of the p_1 -brane. We now compute the cohomology of that geometric realization. In a full discussion one will have to pull the result of the following computation back along the above map to \tilde{X} , and kernel and cokernel of this pullback map potentially yield yet further corrections to the brane charges.

We specialize to the case $p_1 = 2$ and $p_2 = 5$ corresponding to M5-branes propagating in an M2-brane condensate [21]. First, consider the cohomology of the fiber $K(\mathbb{Z}, 3)$. At the integral level, this is known but has a complicated structure. We will instead consider the corresponding rational cohomology, which is much more accessible. Indeed, it directly follows from the Hurewicz theorem and the universal coefficient theorem that

$$H^k(K(\mathbb{Z}, 3); \mathbb{Q}) = \begin{cases} \mathbb{Q} & \text{for } k = 0, 3, \\ 0 & \text{otherwise.} \end{cases}$$

Given the homotopy fiber sequence

$$\begin{array}{ccc} K(\mathbb{Z}, 3) & \longrightarrow & \hat{X} \\ & & \downarrow \\ & & X \end{array}$$

the cohomology Serre spectral sequence takes the form

$$E_2^{p,q} = H^p(X); \quad H^q(K(\mathbb{Z}, 3)) \Rightarrow H^{p+q}(\hat{X}).$$

From the cohomology of the fiber determined above, we see that q has to be either 0 or 3 in order to contribute. The relevant differential $d_r : E_2^{p,q} \rightarrow E_2^{p+r,q-r+1}$ is then d_4 , which raises the cohomology degree by 4.

Hence the brane charge extension of the 11-dimensional superisometries is, rationally, by $H^5(\hat{X})$, and by the Serre spectral sequence this is the middle cohomology of

$$H^1(X) \xrightarrow{(0,d_4)} H^2(X) \oplus H^5(X) \xrightarrow{(d_4,0)} H^6(X), \tag{5.2}$$

where $d_4 = [G_4] \cup (-)$ is the cup product with the degree-4 class of the C-field. For torsion C-fields this vanishes rationally and hence one arrives at the conclusion that the M-theory super Lie algebra extension is by brane charges in $H^2(X) \oplus H^5(X)$, agreeing with the result of the argument in [40]. For non-torsion C-fields or else when considered not just rationally, then there are corrections to this statement by the kernel and cokernel of d_4 . Notice that these corrections are directly analogous to the correction by kernel and cokernel of a d_3 differential in an Atiyah-Hirzebruch spectral sequence, which appear when refining D-brane charges from ordinary cohomology to twisted K-theory [26, (3.2), (3.6)]. That 5-brane charges should be in a degree-4 twisted cohomology theory this way has been suggested earlier in [35, section 8] and has been discussed further in [22].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R. D'Auria and P. Fré, *Geometric Supergravity in $D = 11$ and Its Hidden Supergroup*, *Nucl. Phys. B* **201** (1982) 101 [Erratum *ibid.* **B 206** (1982) 496] [INSPIRE].
- [2] J.A. de Azcárraga, J.P. Gauntlett, J.M. Izquierdo and P.K. Townsend, *Topological Extensions of the Supersymmetry Algebra for Extended Objects*, *Phys. Rev. Lett.* **63** (1989) 2443 [INSPIRE].
- [3] J.A. de Azcárraga and J. Izquierdo, *Lie Groups, Lie Algebras, Cohomology and Some Applications in Physics*, Cambridge Monographs of Mathematical Physics, Cambridge University Press (1995).
- [4] D.V. Alekseevsky, V. Cortés, C. Devchand and A. Van Proeyen, *Polyvector super-Poincaré algebras*, *Commun. Math. Phys.* **253** (2004) 385 [hep-th/0311107] [INSPIRE].
- [5] M.F. Atiyah and G.B. Segal, *Twisted k -theory and cohomology*, in *Nankai Tracts in Mathematics: Volume 11*, World Scientific (2006) [math.KT/0510674] [INSPIRE].
- [6] J.C. Baez and C.L. Rogers, *Categorified Symplectic Geometry and the String Lie 2-Algebra*, *Homology Homotopy Appl.* **12** (2010) 221 [arXiv:0901.4721] [INSPIRE].
- [7] G. Barnich, R. Fulp, T. Lada and J. Stasheff, *The sh Lie structure of Poisson brackets in field theory*, *Commun. Math. Phys.* **191** (1998) 585 [hep-th/9702176] [INSPIRE].
- [8] G. Barnich and M. Henneaux, *Isomorphisms between the Batalin-Vilkovisky anti-bracket and the Poisson bracket*, *J. Math. Phys.* **37** (1996) 5273 [hep-th/9601124] [INSPIRE].
- [9] E. Bergshoeff, E. Sezgin and P.K. Townsend, *Superstring Actions in $D = 3, 4, 6, 10$ Curved Superspace*, *Phys. Lett. B* **169** (1986) 191 [INSPIRE].
- [10] E. Bergshoeff, E. Sezgin and P.K. Townsend, *Supermembranes and Eleven-Dimensional Supergravity*, *Phys. Lett. B* **189** (1987) 75 [INSPIRE]
<http://streaming.ictp.trieste.it/preprints/P/87/010.pdf>.
- [11] J.-L. Brylinski, *Loop spaces, Characteristic Classes and Geometric Quantization*, Birkhäuser (1993).
- [12] A.K. Callister and D.J. Smith, *Topological BPS charges in 10-dimensional and 11-dimensional supergravity*, *Phys. Rev. D* **78** (2008) 065042 [arXiv:0712.3235] [INSPIRE].
- [13] C. Chryssomalakos, J.A. de Azcárraga, J.M. Izquierdo and J.C. Pezéz Bueno, *The Geometry of branes and extended superspaces*, *Nucl. Phys. B* **567** (2000) 293 [hep-th/9904137] [INSPIRE].
- [14] L. Dickey, *Soliton equations and Hamiltonian systems*, Advanced Series in Mathematical Physics, volume 26, World Scientific (1991).
- [15] M. Ferraris, M. Palese and E. Winterroth, *Local variational problems and conservation laws*, *Differ. Geom. Appl.* **29** (2011) S80.
- [16] D. Fiorenza, C.L. Rogers and U. Schreiber, *Higher $U(1)$ -gerbe connections in geometric prequantization*, *Rev. Math. Phys.* **28** (2016) 1650012 [arXiv:1304.0236] [INSPIRE].
- [17] D. Fiorenza, C.L. Rogers and U. Schreiber, *L_∞ -algebras of local observables from higher prequantum bundles*, *Homology Homotopy Appl.* **16** (2014) 107 [arXiv:1304.6292].
- [18] D. Fiorenza, H. Sati and U. Schreiber, *Extended higher cup-product Chern-Simons theories*, *J. Geom. Phys.* **74** (2013) 130 [arXiv:1207.5449] [INSPIRE].

- [19] D. Fiorenza, H. Sati and U. Schreiber, *The E_8 Moduli 3-Stack of the C-Field in M-theory*, *Commun. Math. Phys.* **333** (2015) 117 [[arXiv:1202.2455](#)] [[INSPIRE](#)].
- [20] D. Fiorenza, H. Sati and U. Schreiber, *A higher stacky perspective on Chern-Simons theory*, in *Mathematical Aspects of Quantum Field Theories*, D. Calaque and T. Strobl eds., Springer International Publishing (2014) [[arXiv:1301.2580](#)] [[INSPIRE](#)].
- [21] D. Fiorenza, H. Sati and U. Schreiber, *Super Lie n -algebra extensions, higher WZW models and super p -branes with tensor multiplet fields*, *Int. J. Geom. Meth. Mod. Phys.* **12** (2015) 1550018 [[arXiv:1308.5264](#)] [[INSPIRE](#)].
- [22] D. Fiorenza, H. Sati and U. Schreiber, *The Wess-Zumino-Witten term of the M5-brane and differential cohomotopy*, *J. Math. Phys.* **56** (2015) 102301 [[arXiv:1506.07557](#)] [[INSPIRE](#)].
- [23] M. Henneaux and L. Mezincescu, *A σ -model Interpretation of Green-Schwarz Covariant Superstring Action*, *Phys. Lett. B* **152** (1985) 340 [[INSPIRE](#)].
- [24] C.M. Hull, *Gravitational duality, branes and charges*, *Nucl. Phys. B* **509** (1998) 216 [[hep-th/9705162](#)] [[INSPIRE](#)].
- [25] T. Lada and J. Stasheff, *Introduction to SH Lie algebras for physicists*, *Int. J. Theor. Phys.* **32** (1993) 1087 [[hep-th/9209099](#)] [[INSPIRE](#)].
- [26] J.M. Maldacena, G.W. Moore and N. Seiberg, *D-brane instantons and k -theory charges*, *JHEP* **11** (2001) 062 [[hep-th/0108100](#)] [[INSPIRE](#)].
- [27] M. Markl and S. Shnider, *Differential Operator Endomorphisms of an Euler-Lagrange Complex*, *Contemp. Math.* **231** (1999) MR1707343 [[math.DG/9808105](#)] [[INSPIRE](#)].
- [28] J. Mickelsson, *Current Algebras and Groups*, Springer (1989).
- [29] D. Pavlov, D. Berwick-Evans and P. Boavida de Brito, *Concordance theory for homotopy sheaves*, in preparation.
- [30] A. Pressley and G.B. Segal, *Loop groups*, Oxford University Press (1988).
- [31] P. Ritter and C. Sämann, *Lie 2-algebra models*, *JHEP* **04** (2014) 066 [[arXiv:1308.4892](#)] [[INSPIRE](#)].
- [32] C.L. Rogers, *L_∞ algebras from multisymplectic geometry*, *Lett. Math. Phys.* **100** (2012) 29 [[arXiv:1005.2230](#)] [[INSPIRE](#)].
- [33] H. Sati, *Duality-symmetry and the form-fields in M-theory*, *JHEP* **06** (2006) 062 [[hep-th/0509046](#)] [[INSPIRE](#)].
- [34] H. Sati, *An Approach to anomalies in M-theory via $KSpin$* , *J. Geom. Phys.* **58** (2008) 387 [[arXiv:0705.3484](#)] [[INSPIRE](#)].
- [35] H. Sati, *Geometric and topological structures related to M-branes*, *Proc. Symp. Pure Math.* **81** (2010) 181 [[arXiv:1001.5020](#)] [[INSPIRE](#)].
- [36] H. Sati, U. Schreiber and J. Stasheff, *L_∞ algebra connections and applications to String- and Chern-Simons n -transport*, in *Recent developments in QFT*, Birkhäuser (2009), pp. 303–424 [[arXiv:0801.3480](#)] [[INSPIRE](#)].
- [37] G.B. Segal, *Loop groups*, *Lect. Notes Math.* **1111** (1984) 155 <http://people.mpim-bonn.mpg.de/zagier/files/doi/10.1007/BFb0084581/chapter08.pdf>.
- [38] U. Schreiber, *Differential cohomology in a cohesive infinity-topos*, [arXiv:1310.7930](#) [[INSPIRE](#)].

- [39] U. Schreiber, *Structure theory for higher WZW terms*, in *Flavors of cohomology*, Pittsburgh, U.S.A., 3–5 June (2015)
<http://ncatlab.org/schreiber/show/Structure+Theory+for+Higher+WZW+Terms>.
- [40] D.P. Sorokin and P.K. Townsend, *M Theory superalgebra from the M five-brane*, *Phys. Lett. B* **412** (1997) 265 [[hep-th/9708003](#)] [[INSPIRE](#)].
- [41] P.K. Townsend, *p-brane democracy*, in *Particles, Strings and Cosmology*, J. Bagger, G. Domokos, A Falk and S. Kovesi-Domokos eds., World Scientific (1996), pp. 271–285 [[hep-th/9507048](#)] [[INSPIRE](#)].
- [42] I.S. Krasil'shchik et al., *Symmetries and conservation laws for differential equations of mathematical physics*, Translations of Mathematical Monographs, volume 182, AMS (1999).