Twining genera of (0,4) supersymmetric sigma models on K3

Sarah Harrison, Shamit Kachru and Natalie M. Paquette

*SITP, Department of Physics and Theory Group, SLAC, Stanford University
Stanford, CA 94305, U.S.A.*

E-mail: sarharr@stanford.edu, skachru@stanford.edu,
npaquette@stanford.edu

ABSTRACT: Conformal field theories with (0,4) worldsheet supersymmetry and K3 target can be used to compactify the $E_8 \times E_8$ heterotic string to six dimensions in a supersymmetric manner. The data specifying such a model includes an appropriate configuration of 24 gauge instantons in the $E_8 \times E_8$ gauge group to satisfy the constraints of anomaly cancellation. In this note, we compute twining genera — elliptic genera with appropriate insertions of discrete symmetry generators in the trace — for (0,4) theories with various instanton embeddings. We do this by constructing linear sigma models which flow to the desired conformal field theories, and using the techniques of localization. We present several examples of such twining genera which are consistent with a moonshine relating these (0,4) models to the finite simple sporadic group M_{24} .

KEYWORDS: Superstring Vacua, Conformal Field Models in String Theory, Sigma Models

ARXIV EPRINT: [1309.0510](https://arxiv.org/abs/1309.0510)

Contents

1	Introduction	1
2	Some simple (0,4) gauged linear sigma models	3
2.1	Basic multiplets and terms in the action	3
2.2	The class of models of interest	4
2.3	Specific examples of models and discrete symmetries	6
2.3.1	Model 1: a (24,0) model	6
2.3.2	Model 2: a (12,12) model	8
2.3.3	Model 3: a (14,10) model	8
2.3.4	Model 4: an (18,6) model	8
3	Computation of the twining genera	9
3.1	Residue formula for elliptic genus	9
3.1.1	K3 elliptic genus	10
3.2	Residue formula for twining genera	11
3.3	Examples	12
3.3.1	Model 1	12
3.3.2	Model 2	12
3.3.3	Model 3	13
3.3.4	Model 4	13
4	Discussion	13
A	Conventions	14
B	M_{24} character table and coefficients of twining genera	15

1 Introduction

To obtain a $(1, 0)$ supersymmetric compactification of the heterotic string to six dimensions, one should choose an internal CFT with $(0,4)$ supersymmetry and right-moving central charge $c_R = 6$. In the realm of geometry, such CFTs arise as non-linear sigma models with K3 target. In order to satisfy the Bianchi identity for the three-form field strength H of the heterotic string

$$dH = \text{Tr}(R \wedge R) - \text{Tr}(F \wedge F), \quad (1.1)$$

one should further embed 24 instantons into the $E_8 \times E_8$ gauge group. If one chooses bundles $V_{1,2}$ of rank $r_{1,2}$ in the two E_8 s (which should be stable and holomorphic, and have

vanishing first Chern class $c_1 = 0$, in the simplest case), then the left-moving fermions in the sigma model couple to the gauge connections on these bundles, and $c_L = 4 + r_1 + r_2$.

The explicit construction of such (0,4) CFTs is a difficult task, and computations of observables in such intricate theories are in general complicated to perform. Ideally, one would like to be able to compute the partition function of the internal conformal field theory. But more generally, one has to settle for obtaining coarser index information. One such compromise is given by the elliptic genus,

$$Z(\tau, z) = \text{Tr}(-1)^F y^{J_L} q^{L_0} \bar{q}^{\bar{L}_0}, \quad y = e^{2\pi i z}, \quad q = e^{2\pi i \tau}. \quad (1.2)$$

This is a graded trace over the Hilbert space of the left movers, containing further information about quantum numbers under a left-moving $U(1)$ current algebra whose generator is J_L .

In this paper, our focus will be on explicit examples of (0,4) models and their twining genera, which are close relatives of the elliptic genus. They can be defined as follows. Consider a (0,4) theory with discrete symmetry g . Then, one can modify (1.2) to

$$Z_g(\tau, z) = \text{Tr}(-1)^F g y^{J_L} q^{L_0} \bar{q}^{\bar{L}_0}, \quad (1.3)$$

that is, one can take the trace with an insertion of the action of g on the physical states.

We will construct (0,4) models as gauged linear sigma models with $K3$ target. The basic ideas involved in constructing such sigma models with Calabi-Yau target were developed in the beautiful paper [1], and the extension to (0,2) models was discussed in detail in [2]. As (0,4) models are a simpler subset of (0,2) models, our models will be simple examples of the constructions in [2].

We will compute the twining genera by using the techniques of localization. Localization was recently used to give a very explicit formula for the elliptic genus of linear sigma models with rank one gauge groups in [3], with an extension to higher ranks appearing in [4]; a small modification of that formula suffices to compute the twining genera (1.3). Earlier results on the elliptic genera of (0,2) gauged linear sigma models appeared in [5], which also anticipated (without derivation) aspects of the residue formula of [3].

While one justification for computing the observables (1.3) is that they contain valuable information about the spectrum of an interacting conformal field theory, we had a more specific motivation for undertaking this study. There is a Mathieu moonshine relating the (4,4) sigma models with $K3$ target to the Mathieu group M_{24} [6]. The key first piece of evidence for this moonshine was a decomposition of the coefficients of the expansion of the elliptic genus of $K3$ in $\mathcal{N} = 4$ characters, in terms of dimensions of representations of M_{24} . Given such a decomposition, one can make predictions for the twining genera (1.3) for the (4,4) theories, if one inserts any element of M_{24} . By finding explicit realizations of symmetries of $K3$ sigma models, and computing (1.3) explicitly, one can check whether these symmetries correspond to (conjugacy classes of) elements of the hypothetical M_{24} . Such checks were carried out in [7–10] with impressive results. The existence of a graded M_{24} module with the desired properties has since been proved at a rigorous level [11].

Possible extensions of this moonshine to theories with only half as much supersymmetry, including (0,4) heterotic string compactifications, were discussed in [12]. As a logical extension of that work, it is desirable to find explicit symmetries of (0,4) K3 conformal field theories and check if the twining genera (1.3) match with those of suitable M_{24} conjugacy classes. This note, as well as the companion [13] to [12] which studies twining genera of (0,4) supersymmetric K3 orbifold conformal field theories, will present examples where exactly such matching can be seen.

2 Some simple (0,4) gauged linear sigma models

2.1 Basic multiplets and terms in the action

We will write down (0,4) linear sigma models by working in (0,2) superspace and using vector bundles constructed as the cohomology of an exact sequence, as in [2]. The enhanced worldsheet supersymmetry is not manifest, but should be expected to emerge in the IR on general grounds when we construct models which have a large-radius interpretation as K3 sigma models.

The (0,2) multiplets we use are as follows (see [2, 14] for more discussion). (0,2) superspace has coordinates $(z, \bar{z}, \theta^+, \theta^-)$ (so $\pm$ here on the Grassman coordinates denotes U(1) charge, not chirality). The spinor superderivatives are

$$\bar{D}_{\pm} = \frac{\partial}{\partial \theta^{\pm}} + \theta^{\mp} \partial_{\bar{z}}. \quad (2.1)$$

Chiral superfields Φ satisfy

$$\bar{D}_+ \Phi = 0 \quad (2.2)$$

and have a component expansion

$$\Phi = \phi + \theta^- \psi + \theta^- \theta^+ \partial_{\bar{z}} \phi \quad (2.3)$$

with ψ a right-moving fermion. Fermi superfields Λ also satisfy $\bar{D}_+ \Lambda = 0$, but have component expansion

$$\Lambda = \lambda + \theta^- \ell + \theta^- \theta^+ \partial_{\bar{z}} \lambda \quad (2.4)$$

instead, with λ a left-moving fermion and ℓ an auxiliary field.

We will be considering (0,2) gauge theories with U(1) gauge group, so we also need to discuss the (0,2) gauge multiplet. It consists of a pair of superfields $V, \mathcal{A}$ whose expansion, in Wess-Zumino gauge, is given by

$$\begin{aligned} V &= \theta^- \theta^+ \bar{a} \\ \mathcal{A} &= a + \theta^+ \alpha - \theta^- \bar{\alpha} + \frac{1}{2} \theta^- \theta^+ D \end{aligned} \quad (2.5)$$

with $a, \bar{a}$ the left/right moving pieces of the gauge field, $\alpha, \bar{\alpha}$ left-moving gauginos, and D an auxiliary field. The field strength supermultiplets are

$$\begin{aligned} \mathcal{F} &= -\alpha + \theta^- (D + f) - \theta^- \theta^+ \partial_{\bar{z}} \alpha \\ \bar{\mathcal{F}} &= -\bar{\alpha} + \theta^+ (D - f) + \theta^- \theta^+ \partial_{\bar{z}} \alpha \end{aligned} \quad (2.6)$$

where

$$f = 2(\partial_z \bar{a} - \partial_{\bar{z}} a) . \quad (2.7)$$

The basic terms which appear in a supersymmetric action will be the following. A gauge invariant kinetic term for a charged chiral multiplet Φ with charge Q is

$$\begin{aligned} S_\Phi = & \int d^2z (\partial_z - Qa) \bar{\phi} (\partial_{\bar{z}} + Q\bar{a}) \phi + (\partial_{\bar{z}} - Q\bar{a}) \bar{\phi} (\partial_z + Qa) \phi \\ & + 2\bar{\psi} (\partial_z + Qa) \psi + Q(\bar{\alpha} \bar{\psi} \phi - \alpha \phi \bar{\psi}) - Q D \bar{\phi} \phi , \end{aligned} \quad (2.8)$$

while a gauge invariant kinetic term for a charged Fermi multiplet Λ of charge Q is

$$S_\Lambda = \int d^2z 2\bar{\lambda} (\partial_{\bar{z}} + Q\bar{a}) \lambda - \bar{\ell} \ell . \quad (2.9)$$

The gauge kinetic term is

$$\begin{aligned} S_{\text{gauge}} &= -\frac{1}{2e^2} \int d^2z d^2\theta \mathcal{F} \bar{\mathcal{F}} \\ &= \frac{1}{2e^2} \int d^2z (f^2 - D^2 + 2\alpha \partial_{\bar{z}} \bar{\alpha}) . \end{aligned} \quad (2.10)$$

The Fayet-Iliopoulos D-term and θ angle are

$$S_D = r \int d^2z D - i \frac{\theta}{2\pi} \int d^2z f \quad (2.11)$$

(where $t = \frac{\theta}{2\pi} + ir$ plays the role of a Kähler parameter in large radius geometric phases of the theories to come). The (0,2) superpotential takes the form

$$\begin{aligned} S_W &= \int d^2z d\theta^- \Lambda F(\Phi) + \text{h.c.} \\ &= \int d^2z (\ell F(\phi) - \lambda \frac{\partial F}{\partial \phi} \psi) + \text{h.c.} . \end{aligned} \quad (2.12)$$

Here, F needs to be chosen to be a homogeneous polynomial of the appropriate degree in the charged field Φ so that (2.12) is gauge invariant.

2.2 The class of models of interest

Our interest is to describe stable, holomorphic vector bundles V with $c_1(V) = 0$ and $c_2(V) = c_2(TM)$ over K3 surfaces M . A simple class of models which admits a gauged linear sigma model description is the following. We choose for M the Calabi-Yau hypersurface in the $W\mathbb{P}^3$ with weights w_i ($i = 1, \dots, 4$), described by the equation

$$W(\phi_i) = 0 \subset W\mathbb{P}_{w_1, \dots, w_4}^3 . \quad (2.13)$$

We define V as the cohomology of the exact sequence

$$0 \rightarrow V \rightarrow \oplus_a \mathcal{O}(n_a) \xrightarrow{\otimes F_a(\phi)} \mathcal{O}(m) \rightarrow 0 . \quad (2.14)$$

Field	Gauge charge
Φ^i	w_i
P	$-m$
Λ^a	n_a
Γ	$-d$

Table 1. Field content of our (0,2) sigma models

The conditions that $c_1(TM) = c_1(V) = 0$ and $c_2(TM) = c_2(V)$ are captured by the Diophantine equations

$$\begin{aligned} \sum_i w_i &= d, \quad \sum_a n_a = m, \\ m^2 - \sum_a n_a^2 &= d^2 - \sum_i w_i^2, \end{aligned} \quad (2.15)$$

with d being the degree of the defining polynomial $W(\phi)$ of the K3 surface; these equations follow simply from the adjunction formula for Chern classes. The second equation in (2.15) just imposes the requirement of worldsheet gauge anomaly cancellation for the abelian gauge field.

These theories can be represented as gauged linear sigma models in the following way. Let us consider the (0,2) supersymmetric abelian gauge theory with the matter content shown in table 1.

Φ^i, P are (0,2) chiral multiplets, while Λ^a and Γ are Fermi multiplets. For our (0,2) superpotential we choose

$$\int d^2z \, d\theta^- (\Gamma W(\Phi) + P \Lambda^a F_a(\Phi)) + \text{h.c.} \quad (2.16)$$

with W, F_a coinciding with the data in the definition of the K3 hypersurface and the bundle V above. One can verify, as in [2], that in the limit of large r , this theory flows to the sigma model governed by the geometric objects (2.13) and (2.14), with the scalars living on the hypersurface (2.13) while the left-moving fermions transform as sections of the bundle (2.14). Of course, as one varies the Fayet-Iliopoulos parameters in such a gauge theory, other interesting phases can arise (with Landau-Ginzburg orbifold phases being a prototypical such phase).

We will need to generalize this construction in a trivial way, in order to capture the geometry of two non-trivial bundles $V_{1,2}$ which we embed into the two E_8 s. The appropriate generalization is to introduce two chiral analogues of the P field $P^{1,2}$, with charges $m_{1,2}$, and two sets of Fermi multiplets Λ_1^a and Λ_2^α of charges n_a and q_α , with $a = 1, \dots, r_1 + 1$ and $\alpha = 1, \dots, r_2 + 1$. The superpotential is now

$$\int d^2z \, d\theta^- (\Gamma W(\phi) + P_1 \Lambda_1^a F_a(\Phi) + P_2 \Lambda_2^\alpha G_\alpha(\Phi)), \quad (2.17)$$

with F_a and G_α defining the bundles $V_{1,2}$ through exact sequences as in (2.14). The constraints on the Chern classes now become

$$\begin{aligned} m_1 &= \sum n_a, & m_2 &= \sum q_\alpha \\ d^2 - \sum w_i^2 &= (m_1^2 - \sum n_a^2) + (m_2^2 - \sum q_\alpha^2). \end{aligned} \quad (2.18)$$

Again, the second equation in (2.18) is required for gauge anomaly cancellation, and is interpreted in space-time as implementing the condition

$$c_2(TM) = c_2(V_1) + c_2(V_2), \quad (2.19)$$

which is required to satisfy the Bianchi identity (1.1).

Intuitively, the equation (2.19) means that in perturbative supersymmetric heterotic models on K3, one should choose non-negative integers $n^{(1)}, n^{(2)}$ with

$$n^{(1)} + n^{(2)} = 24, \quad (2.20)$$

and place $n^{(1)}$ and $n^{(2)}$ gauge instantons in the two E_8 s. Our goal in the next section will be to show that in a variety of examples constructed as above, reflecting distinct choices of $n_{1,2}$, one can find (0,4) sigma models with discrete symmetries g whose twining genera (1.3) are consistent with the properties expected from Mathieu moonshine for (0,4) models. This strengthens the case made in [12] that moonshine extends to a portion of the web of 4d $\mathcal{N} = 2$ (or 6d $\mathcal{N} = 1$) supersymmetric heterotic string theories, as well as their type II (or F-theory) Calabi-Yau duals.

2.3 Specific examples of models and discrete symmetries

We will focus on four classes of specific models with different values of $n^{(1)}$ and $n^{(2)}$, but it should be clear that many other models exist and could be fruitfully analyzed in this way. In each case, we just discuss some simple symmetries which arise for easy choices of the defining data; we are not exhaustive. We label the models by the instanton numbers $(n^{(1)}, n^{(2)})$ chosen in each. The four models we will study are:

2.3.1 Model 1: a (24,0) model

For our first example, we will study the theory with $d = 4$ and $w_i = 1, 1, 1, 1$. To begin with, we can choose the defining data of the target manifold to be

$$W(\Phi_i) = \sum_i \frac{1}{4} \Phi_i^4, \quad (2.21)$$

i.e. the Fermat point in the moduli space of this K3 hypersurface. The bundle is defined by choosing

$$V_1 : m = 4, \{n_a\} = \{1, 1, 1, 1\}, F_a(\phi) = \Phi_a^3. \quad (2.22)$$

For generic defining data, this model simply defines the (0,4) model obtained by deforming the tangent bundle of $K3$ away from the (4,4) supersymmetric locus, while extending the rank from $SU(2)$ to $SU(3)$ (by partially Higgsing the E_7 space-time gauge group with a **56**

of E_7). It has instanton numbers ($n^{(1)} = 24, n^{(2)} = 0$). The (4,4) theory was studied in more detail in [15]; we simply discuss this model here to provide a warm-up on more or less familiar territory.

We can study several simple symmetries in the Fermat K3. We will study three:

1. The $\mathbb{Z}_2$ symmetry which acts as

$$g : \Phi_{1,2} \rightarrow -\Phi_{1,2}, \quad \Lambda_{1,2} \rightarrow -\Lambda_{1,2} . \quad (2.23)$$

2. The S_3 symmetry which acts as a permutation of cycle shape (123) on $\Lambda_{1,2,3}$ and $\Phi_{1,2,3}$.
3. The $\mathbb{Z}_4$ symmetry

$$g : \Phi_{1,2} \rightarrow \pm i \Phi_{1,2}, \quad \Lambda_{1,2} \rightarrow \pm i \Lambda_{1,2} . \quad (2.24)$$

We can also obtain more elaborate symmetries by choosing slightly different data. For instance, if we choose a complex structure

$$W(\Phi_i) = \Phi_1^3 \Phi_2 + \Phi_2^3 \Phi_3 + \Phi_3^3 \Phi_4 + \Phi_4^3 \Phi_1 \quad (2.25)$$

then we can find a $\mathbb{Z}_5$ symmetry:

4. $\mathbb{Z}_5$ symmetry:

$$\begin{aligned} g : \quad \Phi_1 &\rightarrow \lambda \Phi_1, \quad \Phi_2 \rightarrow \lambda^2 \Phi_2, \quad \Phi_3 \rightarrow \lambda^4 \Phi_3, \quad \Phi_4 \rightarrow \lambda^3 \Phi_4, \quad \lambda \equiv e^{\frac{2\pi i}{5}}, \\ \Lambda_1 &\rightarrow \lambda \Lambda_1, \quad \Lambda_2 \rightarrow \lambda^2 \Lambda_2, \quad \Lambda_3 \rightarrow \lambda^4 \Lambda_3, \quad \Lambda_4 \rightarrow \lambda^3 \Lambda_4 . \end{aligned} \quad (2.26)$$

Defining data for the vector bundle which respects this symmetry could include e.g. $F_a(\Phi) = \frac{\partial W}{\partial \Phi_a}$ or suitable variants.

Another K3 which admits an interesting symmetry has the complex structure

$$W(\Phi_i) = \Phi_1^3 \Phi_2 + \Phi_2^3 \Phi_3 + \Phi_3^3 \Phi_1 + \Phi_4^4 . \quad (2.27)$$

This surface admits the $\mathbb{Z}_7$ symmetry:

5. $\mathbb{Z}_7$ symmetry:

$$\begin{aligned} g : \quad \Phi_1 &\rightarrow \lambda \Phi_1, \quad \Phi_2 \rightarrow \lambda^4 \Phi_2, \quad \Phi_3 \rightarrow \lambda^2 \Phi_3, \quad \lambda \equiv e^{\frac{2\pi i}{7}}, \\ \Lambda_1 &\rightarrow \lambda \Lambda_1, \quad \Lambda_2 \rightarrow \lambda^4 \Lambda_2, \quad \Lambda_3 \rightarrow \lambda^2 \Lambda_3. \end{aligned} \quad (2.28)$$

Again suitable defining data for the bundle could be $F_a(\Phi) = \frac{\partial W}{\partial \Phi_a}$ with other choices also possible.

2.3.2 Model 2: a (12,12) model

Again reverting to the Fermat quartic K3 (2.21), we choose now bundles $V_{1,2}$ each with $m_{1,2} = 3$ and $\{n_a\}, \{q_\alpha\} = \{1, 1, 1\}$. We consider the symmetries:

1. A $\mathbb{Z}_2$ with

$$g : \Lambda_{1,2,3} \rightarrow -\Lambda_{1,2,3}, \quad P_1 \rightarrow -P_1, \quad \Phi_{1,2} \rightarrow -\Phi_{1,2}. \quad (2.29)$$

Here the Λ s are those spanning V_1 , and one should choose data $F_a(\phi)$ which is consistent with the symmetry.

2. A $\mathbb{Z}_4$ with

$$g : \Lambda_{1,2} \rightarrow \pm i \Lambda_{1,2}, \quad \Phi_{1,2} \rightarrow \pm i \Phi_{1,2}. \quad (2.30)$$

Again, these fermions are from V_1 , and one should choose data $F_{1,2}(\phi)$ consistent with the symmetry.

2.3.3 Model 3: a (14,10) model

Now, we work on the K3 hypersurface embedded in $W\mathbb{P}_{1,1,1,3}^3$. For a defining equation, we choose

$$W(\Phi) = \Phi_1^6 + \Phi_2^6 + \Phi_3^6 + \Phi_4^2 \quad (2.31)$$

For bundles, we let V_1 be specified by $m_1 = 5, \{n_a\} = \{3, 1, 1\}$ and V_2 be specified by $m_2 = 4, \{q_\alpha\} = \{2, 1, 1\}$.

We consider two symmetries in this model:

1. A representative $\mathbb{Z}_2$ symmetry is, for instance,

$$g : \Lambda_{2,3} \rightarrow -\Lambda_{2,3}, \quad \Phi_{2,3} \rightarrow -\Phi_{2,3}, \quad (2.32)$$

with the Λ s being fermions involved in the construction of V_1 . Simple choices of the $F_a(\Phi)$ are consistent with such a symmetry.

2. We can consider a $\mathbb{Z}_3$ symmetry as follows:

$$g : \Phi_1 \rightarrow e^{\frac{2\pi i}{3}} \Phi_1, \quad \Phi_2 \rightarrow e^{\frac{4\pi i}{3}} \Phi_2 \quad (2.33)$$

with the two charge 1 fermions in V_1 , $\Lambda_{2,3}$, rotating as

$$g : \Lambda_2 \rightarrow e^{\frac{4\pi i}{3}} \Lambda_2, \quad \Lambda_3 \rightarrow e^{\frac{2\pi i}{3}} \Lambda_3. \quad (2.34)$$

There are simple choices of the $F_a(\Phi)$ that accomodate this symmetry.

2.3.4 Model 4: an (18,6) model

Finally, still working on the K3 hypersurface (2.31), we study the bundles V_1 with $m_1 = 5, \{n_a\} = \{2, 1, 1, 1\}$ and V_2 with $m_2 = 3, \{q_\alpha\} = \{1, 1, 1\}$. One $\mathbb{Z}_3$ symmetry arises in this model by permuting the fermions $\Lambda_{2,3,4}$ of charge 1 arising as part of V_1 ; the fermions $\Lambda_{4,5,6}$ arising as part of V_2 ; and the chiral fields $\Phi_{1,2,3}$, all with the permutation of cycle shape (123). Once again, simple choices of the bundle data $F(\Phi)$ are consistent with such a symmetry.

3 Computation of the twining genera

In this section, we compute the twining genera under the various model symmetries described in §2.3. We begin by outlining the general strategy and formulae that are relevant, and then simply present the results of applying these formulae to the various cases. Our work relies heavily on the elegant residue formula derived recently in [3].

3.1 Residue formula for elliptic genus

The elliptic genus was first discussed in [16–19]. Its application to string compactification was pioneered in [20], and it was first computed by localization in (2,2) supersymmetric Landau-Ginzburg models in [21] and for (0,2) models in [5]. It has recently been the focus of attention in, for instance, [3, 4, 22]

The formalism we discuss only assumes $\mathcal{N} = (0, 2)$ supersymmetry, though our application will be to (0,4) theories. Although in many discussions of the elliptic genus in theories with (2,2) supersymmetry the left-moving R-symmetry plays a crucial role, here there is no longer a left-moving R-charge. However, the models we consider will have an extra U(1) global current J_L , and we will grade by the quantum number under the associated charge in the elliptic genus. In the models described in the previous section, $J_L = 0$ for Γ and Φ_i , and for the $\Lambda_{a,\alpha}$, $J_L = -1$, whereas for the $P_{a,\alpha}$, $J_L = +1$.

We follow the discussion of (0,2) abelian gauge theory in [3]. Let us define u to be the holonomy of the U(1) gauge field around the cycles of the torus

$$u = \oint A_t dt - \tau \oint A_s ds \quad (3.1)$$

with t, s the temporal and spatial directions, and τ the modular parameters of the torus. The elliptic genus is given by the graded trace

$$Z(\tau, z) = \text{Tr}_{\text{RR}}(-1)^F y^{J_L} q^{H_L} \bar{q}^{H_R} . \quad (3.2)$$

Obtaining a formula for (3.2) via localization involves doing an integral over the Wilson lines u of the abelian gauge field.

This integral localizes to a sum of contour integrals around loci (in the moduli space of flat connections) where some of the fields become massless; we refer to these as singular points. Let us consider a general (0,2) U(1) gauge theory, with a number of gauge charged chiral and Fermi multiplets Φ_i and Λ_a , as well as one vector multiplet. Suppose that the charges of the chiral and Fermi multiplets under the gauge and U(1) global symmetry are $Q_{i,a}$ and $J_{i,a}$ respectively. Then, defining

$$x = e^{2\pi i u} , \quad (3.3)$$

the expression that has been obtained for the elliptic genus is [3]

$$Z(\tau, z) = -\eta(q)^2 \sum_{u_j \in \mathcal{M}^+_{u=u_j}} \oint du \prod_{\Phi_i} \frac{i\eta(q)}{\theta_1(q, y^{J_i} x^{Q_i})} \prod_{\Lambda_a} \frac{i\theta_1(q, y^{J_a} x^{Q_a})}{\eta(q)} , \quad (3.4)$$

where $\mathcal{M}^+$ is the relevant set of singular points.¹ These points are defined as the solutions to the equation

$$Q_i u + J_i z \equiv 0 \pmod{(\mathbb{Z} + \tau \mathbb{Z})}, \quad (3.5)$$

with *positive* Q_i . Equivalently one could sum over poles in the set $\mathcal{M}^-$ (including an overall change of sign, due to the reversed orientation of the contour), defined by solutions to the above equation for all *negative* Q_i .

One can roughly understand the origin of the formula (3.4) as follows. Each chiral, Fermi and vector multiplet makes a (multiplicative) contribution to the index at any fixed value of the Wilson lines u . For a (0,2) chiral multiplet with global U(1) charge J and flavor charge Q , the contribution is

$$Z_{\Phi, J, Q}^{(0,2)}(\tau, z, u) = \frac{i\eta(q)}{\theta_1(q, y^J x^Q)}. \quad (3.6)$$

That of a Fermi multiplet with global U(1) charge J is

$$Z_{\Lambda, J, Q}(\tau, z, u) = \frac{i\theta_1(q, y^J x^Q)}{\eta(q)}. \quad (3.7)$$

Finally, the contribution of a (0,2) vector multiplet is

$$Z_{\text{vector}}^{(0,2)}(\tau) = \eta(q)^2. \quad (3.8)$$

independent of u . The product of these expressions over all multiplets present in a given theory, integrated over the u -plane, can be reduced to the formula (3.4).

3.1.1 K3 elliptic genus

The standard results for the elliptic genus of K3 (or in the language of quantum field theory, for the $\mathcal{N} = (4, 4)$ sigma model with K3 target) is [20]

$$Z_{K3}(\tau, z) = 8 \sum_{i=2}^4 \left(\frac{\theta_i(q, y)}{\theta_i(q, 1)} \right)^2, \quad (3.9)$$

which has the expansion

$$Z_{K3} \sim \left(\frac{2}{y} + 20 + 2y \right) + \left(\frac{20}{y^2} - \frac{128}{y} + 216 - 128y + 20y^2 \right) q + \dots \quad (3.10)$$

For a (0, 4) model on K3 with rank r gauge bundle, the elliptic genus is given by

$$Z_{K3}^r = \left(\frac{\theta_1(q, y)}{i\eta(q)} \right)^{r-2} Z_{K3}, \quad (3.11)$$

as derived in [5]. It is easy to check that applying (3.4) to our models of §2.3 agrees with the result (3.11), with $r = r_1 + r_2$ the sum of the ranks of the bundles embedded in the two $E_{8\text{S}}$.

¹Our conventions for modular forms can be found in appendix A.

3.2 Residue formula for twining genera

Our real interest is to compute the elliptic genus with the insertion of a symmetry operator, g , into the path integral

$$Z_g^{(n_1, n_2)}(\tau, z) = \text{Tr}_{\text{RR}} g (-1)^F y^{J_L} q^{H_L} \bar{q}^{H_R} \quad (3.12)$$

for various particular (n_1, n_2) instanton embeddings. We can do this with a slight modification to the computation of the untwined elliptic genus.

Consider an operator g which acts on chiral and Fermi multiplets as

$$g\Phi_i = e^{2\pi i\alpha_i} \Phi_i, \quad g\Lambda_a = e^{2\pi i\beta_a} \Lambda_a, \quad (3.13)$$

and is a symmetry of the action. When inserting this operator into the path integral, it modifies the contribution due to the chiral and Fermi multiplets. The contribution of a $(0, 2)$ chiral multiplet Φ_i to the integrand in (3.4) becomes

$$\frac{i\eta(q) e^{\pi i\alpha_i}}{\theta_1(q, e^{2\pi i\alpha_i} y^{J_i} x^{Q_i})}, \quad (3.14)$$

while one obtains

$$\frac{i\theta_1(q, e^{2\pi i\beta_a} y^{J_a} x^{Q_a})}{e^{i\pi\beta_a} \eta(q)} \quad (3.15)$$

from the twined Fermi multiplet Λ_a . One then sums over the (now shifted) poles that previously contributed to the elliptic genus - the detailed locations of the poles in $\mathcal{M}^+$ on the u -plane, as well as their orders, can be modified depending on the g charges of the fields involved. Since the twining induces an additive shift in the pole location without changing the sign of the chiral gauge charge, no new poles contribute to $\mathcal{M}^+$.

Denote the elliptic genus of the $(4, 4)$ theory twined by a conjugacy class g of M_{24} by Z_g .² Then we expect the twined elliptic genus of an (n_1, n_2) model to decompose as

$$Z_g^{(n_1, n_2)} = \text{ch}(\text{SO}(2r - 4)) Z_g, \quad (3.16)$$

i.e. a product of twined $(4, 4)$ genera and twined $\text{SO}(2r - 4)$ characters.

In writing (3.16), we are making two important assumptions:

- 1) We assume that the M_{24} module which is relevant in the moonshine for $(0, 4)$ models with arbitrary instanton embeddings, has the same representations at each level as the one which arises in the $(4, 4)$ theory. Evidence for this was presented already in the new supersymmetric index computations of [12], which are valid for all instanton embeddings.
- 2) We are assuming that the factor of

$$\left(\frac{\theta_1(q, y)}{i\eta(q)} \right)^{r-2} \quad (3.17)$$

²The Z_g are discussed in detail in appendix B, where also the M_{24} character table and the first few coefficients in the q -expansion of the various Z_g are presented.

in the elliptic genus of a $(0, 4)$ theory with rank r bundle transforms as an element of the spinor minus conjugate spinor representation of $\mathrm{SO}(2r - 4)$. This is motivated by the results to appear in the companion paper about $(0, 4)$ orbifolds [13]. Heuristically, the $\mathrm{SO}(2r - 4)$ symmetry could appear manifestly in a field theory where one deformed the bundle $V_1 \oplus V_2$ to be an $\mathrm{SU}(2)$ bundle with instanton number $n^{(1)} + n^{(2)}$. As the elliptic genus is invariant under such smooth deformations, this may explain the appearance of such factors (related to further ‘hidden symmetries’) in the twining genera of $(0, 4)$ sigma models.

We now show that our results for the set of models discussed in §2.3 satisfy the assumption (3.16). We view this as a check of M_{24} moonshine for $(0, 4)$ theories with a variety of instanton embeddings.

3.3 Examples

3.3.1 Model 1

Here, we considered five symmetries in §2.3.1: a $\mathbb{Z}_2$ symmetry, a $\mathbb{Z}_3$ symmetry, a $\mathbb{Z}_4$ symmetry, a $\mathbb{Z}_5$ symmetry and a $\mathbb{Z}_7$ symmetry. The results for the twining genera are:

$$\begin{aligned} Z_{\mathbb{Z}_2} &= \frac{\theta_1(y)}{i\eta(q)} Z_{2A} , \\ Z_{\mathbb{Z}_3} &= \frac{\theta_1(y)}{i\eta(q)} Z_{3A} , \\ Z_{\mathbb{Z}_4} &= \frac{\theta_1(y)}{i\eta(q)} Z_{4B} , \\ Z_{\mathbb{Z}_5} &= \frac{\theta_1(y)}{i\eta(q)} Z_{5A} , \\ Z_{\mathbb{Z}_7} &= \frac{\theta_1(y)}{i\eta(q)} Z_{7A} . \end{aligned} \tag{3.18}$$

Here, Z_{2A} , Z_{3A} , Z_{4A} , Z_{5A} and Z_{7A} are the corresponding twining genera of the $(4, 4)$ elliptic genus with an insertion in those M_{24} conjugacy classes (see appendix B). The first argument of the theta function has been suppressed here and below.

3.3.2 Model 2

We considered two symmetries in §2.3.2: a $\mathbb{Z}_2$ symmetry and a $\mathbb{Z}_4$ symmetry. The results for the twining genera are:

$$\begin{aligned} Z_{\mathbb{Z}_2} &= \frac{\theta_1(y)^2}{(i\eta(q))^2} Z_{2A} , \\ Z_{\mathbb{Z}_4} &= \frac{\theta_1(iy)\theta_1(-iy)}{(i\eta(q))^2} Z_{4B} . \end{aligned} \tag{3.19}$$

3.3.3 Model 3

Here, we also considered two symmetries in §2.3.3 — a $\mathbb{Z}_2$ and a $\mathbb{Z}_3$. The results are:

$$\begin{aligned} Z_{\mathbb{Z}_2} &= \left(\frac{\theta_1(y)}{i\eta(q)} \right)^2 Z_{2A}, \\ Z_{\mathbb{Z}_3} &= \frac{\theta_1(e^{\frac{2\pi i}{3}} y) \theta_1(e^{\frac{4\pi i}{3}} y)}{(i\eta(q))^2} Z_{3B}. \end{aligned} \quad (3.20)$$

3.3.4 Model 4

We considered a $\mathbb{Z}_3$ symmetry in §2.3.4. The result is

$$Z_{\mathbb{Z}_3} = \frac{\theta_1(y) \theta_1(e^{\frac{2\pi i}{3}} y) \theta_1(e^{\frac{4\pi i}{3}} y)}{(i\eta(q))^3} Z_{3A}. \quad (3.21)$$

4 Discussion

In this note, we used the recently derived localization formula for the elliptic genus of $(0, 2)$ supersymmetric rank one two-dimensional gauge theories [3] to compute twining genera of $(0, 4)$ gauged linear sigma models with K3 target. We did this for a variety of discrete symmetries in $(0, 4)$ models with four different sets of instanton numbers $(n^{(1)}, n^{(2)})$.

In several cases, we found that the simple discrete symmetries give twining genera which are consistent with those of M_{24} elements of the same order, with the trace in the elliptic genus taken over the M_{24} module conjectured to exist in [6] and constructed in [11]. These direct computations are an analogue, for a conjectural $(0, 4)$ moonshine with various instanton numbers, of the twining calculations in [7–10]. Interestingly, the 3B conjugacy class of M_{24} , which does not descend from the classical symmetries of K3 surfaces (as they lie in M_{23} [23, 24]) and which has been elusive, appears here in one of the first cases we examined.

It should not be difficult to find linear sigma models which admit relatively elaborate discrete symmetries. The $\mathbb{Z}_5$ and $\mathbb{Z}_7$ examples of §2.3.1 were found by using a strategy developed in [25], and it seems quite plausible that one can write down examples which show twining in higher order M_{24} conjugacy classes in this way. It should also be instructional to go through the list of e.g. the ‘famous 95’ weighted projective K3 hypersurfaces of Reid [26], and see which of them admit interesting symmetries; this may lead to interesting new examples even in the $(4, 4)$ theory.

A major question which remains is the proper interpretation of the evidence presented here, as well as in [12, 13], for a moonshine relating heterotic $(0, 4)$ theories (and their type II Calabi-Yau duals) to M_{24} . The observations of [15] indicate that M_{24} does not play a canonical role as an embedding group for symmetries of $(4, 4)$ superconformal theories with K3 target. The symmetries available in $(0, 4)$ theories will of course only be richer; developing a classification would be very interesting. Failing a complete classification, a detailed study of particular families with large symmetry groups (extending the philosophy of [27, 28] from the $(4, 4)$ case) could also prove illuminating. It is even within the realm of possibility that some $(0, 4)$ superconformal theory, or perhaps a non-perturbative heterotic

vacuum with small instantons replacing the gauge bundles $V_{1,2}$, could manifest the full symmetry and ‘explain’ the appearance of M_{24} in the elliptic genus. But other interpretations of the moonshine, in terms of Rademacher sums arising naturally in AdS/CFT [29], or in terms of supersymmetric indices of NS5 branes [30], are also quite promising. Related directions to explore are discussed in [31–33].

Acknowledgments

We are grateful to Francesco Benini for helpful discussions about the results in [3], and to Tohru Eguchi and Kazuhiro Hikami for permission to use their tables of M_{24} characters and twining genera in appendix B. We are also very happy to thank Miranda Cheng, Xi Dong, John Duncan, Jeff Harvey, Daniel Whalen and Timm Wrase for extensive discussions about twining genera and moonshine in general. S.H. and S.K. enjoyed the hospitality of the Simons Center workshop on “Mock Modular Forms, Moonshine and String Theory” as this work was completed, and thank the participants for creating a very stimulating intellectual atmosphere. S.K. acknowledges the support of the NSF under grant PHY-0756174, the Department of Energy under contract DE-AC02-76SF00515, and the John Templeton Foundation. S.H. is supported by the John Templeton Foundation, and N.P. by a Stanford Humanities and Sciences Fellowship.

A Conventions

We use the following conventions for the Jacobi $\theta_i(q, y)$ functions

$$\theta_1(q, y) = i \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{(n-\frac{1}{2})^2}{2}} y^{n-\frac{1}{2}}, \quad (\text{A.1})$$

$$\theta_2(q, y) = \sum_{n=-\infty}^{\infty} q^{\frac{(n-\frac{1}{2})^2}{2}} y^{n-\frac{1}{2}}, \quad (\text{A.2})$$

$$\theta_3(q, y) = \sum_{n=-\infty}^{\infty} q^{\frac{n^2}{2}} y^n, \quad (\text{A.3})$$

$$\theta_4(q, y) = \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{n^2}{2}} y^n, \quad (\text{A.4})$$

where $q = e^{2\pi i \tau}$ and $y = e^{2\pi i z}$. Whenever the y -dependence is not specified, we have set $y = 1$, for example $\theta_i = \theta_i(q) = \theta_i(q, 1)$ and likewise for the other functions defined below.

These $\theta_i(q, y)$ functions have the following product expansion

$$\theta_1(q, y) = -iq^{1/8}y^{1/2} \prod_{n=1}^{\infty} (1 - q^n)(1 - yq^n)(1 - y^{-1}q^{n-1}), \quad (\text{A.5})$$

$$\theta_2(q, y) = q^{1/8}y^{1/2} \prod_{n=1}^{\infty} (1 - q^n)(1 + yq^n)(1 + y^{-1}q^{n-1}), \quad (\text{A.6})$$

$$\theta_3(q, y) = \prod_{n=1}^{\infty} (1 - q^n)(1 + yq^{n-\frac{1}{2}})(1 + y^{-1}q^{n-\frac{1}{2}}), \quad (\text{A.7})$$

$$\theta_4(q, y) = \prod_{n=1}^{\infty} (1 - q^n)(1 - yq^{n-\frac{1}{2}})(1 - y^{-1}q^{n-\frac{1}{2}}). \quad (\text{A.8})$$

We also use the Dedekind $\eta(q)$ function

$$\eta(q) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n). \quad (\text{A.9})$$

B M_{24} character table and coefficients of twining genera

In §3, we expressed the results for twining genera in various (0,4) models in terms of the Z_g which appear in the twined elliptic genus of the (4,4) K3 sigma model, for various M_{24} conjugacy classes g . In practice, to work out the q-expansions for the resulting forms, one needs the character table of M_{24} . It is reproduced in table 2 for completeness. The classes appearing before 12B in the top row can also be considered as conjugacy classes in M_{23} , while 12B and those appearing to its right are intrinsic elements of M_{24} with no precursor in M_{23} .

The q-expansions of the Z_g can be written as follows. For the elliptic genus of K3, one writes

$$Z_{K3}(z; \tau) = 20\text{ch}_{h=1/4, \ell=0}(z; \tau) - 2\text{ch}_{h=1/4, \ell=1/2}(z; \tau) + \sum_{n=1}^{\infty} A(n)\text{ch}_{h=n+1/4, \ell=1/2}(z; \tau) \quad (\text{B.1})$$

where $\text{ch}_{h, \ell}$ are characters of the $\mathcal{N} = 4$ superconformal algebra with a given conformal weight and isospin (whose explicit forms can be found in [34, 35]). The M_{24} module associated with this theory via Mathieu moonshine is a graded vector space

$$V = \oplus_{n=1}^{\infty} V(n) \quad (\text{B.2})$$

with $\dim(V(n)) = A(n)$. Then the twining genus Z_g can be written as

$$Z_g(z; \tau) = (\chi_g - 4)\text{ch}_{h=1/4, \ell=0}(z; \tau) - 2\text{ch}_{h=1/4, \ell=1/2}(z; \tau) + \sum_{i=1}^{\infty} A_g(n)\text{ch}_{h=n+1/4, \ell=1/2}(z; \tau), \quad (\text{B.3})$$

with

$$A_g(n) = \text{Tr}_{V(n)} g. \quad (\text{B.4})$$

In practice, one can find simple closed-form expressions for Z_g as discussed in detail in e.g. [7–10]. The first few terms in the q-expansions of the Z_g for various conjugacy classes are shown in table 3.

	1A	2A	3A	5A	4B	7A	7B	8A	6A	11A	15A	15B	14A	14B	23A	23B	12B	6B	4C	3B	2B	10A	21A	21B	4A	12A
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
23	7	5	3	3	2	2	1	1	1	0	0	0	0	0	0	0	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
252	28	9	2	4	0	0	0	1	-1	-1	-1	0	0	0	-1	-1	0	0	0	0	12	2	0	0	4	1
253	13	10	3	1	1	1	-1	-2	0	0	0	0	-1	-1	0	0	1	1	1	1	-11	-1	1	1	-3	0
1771	-21	16	1	-5	0	0	-1	0	0	1	1	0	0	0	0	0	-1	-1	-1	7	11	1	0	0	3	0
3520	64	10	0	0	-1	-1	0	-2	0	0	0	0	1	1	1	1	0	0	0	-8	0	0	-1	-1	0	0
45	-3	0	0	1	e_7^+	e_7^-	-1	0	1	0	0	0	$-e_7^+$	$-e_7^-$	-1	-1	1	-1	1	3	5	0	e_7^+	e_7^-	-3	0
45	-3	0	0	1	e_7^-	e_7^+	-1	0	1	0	0	0	$-e_7^-$	$-e_7^+$	-1	-1	1	-1	1	3	5	0	e_7^-	e_7^+	-3	0
990	-18	0	0	2	e_7^+	e_7^-	0	0	0	0	0	0	e_7^+	e_7^-	1	1	1	-1	-2	3	-10	0	e_7^+	e_7^-	6	0
990	-18	0	0	2	e_7^-	e_7^+	0	0	0	0	0	0	e_7^-	e_7^+	1	1	1	-1	-2	3	-10	0	e_7^-	e_7^+	6	0
1035	-21	0	0	3	$2e_7^+$	$2e_7^-$	-1	0	1	0	0	0	0	0	0	0	-1	1	-1	-3	-5	0	$-e_7^+$	$-e_7^-$	3	0
1035	-21	0	0	3	$2e_7^-$	$2e_7^+$	-1	0	1	0	0	0	0	0	0	0	-1	1	-1	-3	-5	0	$-e_7^-$	$-e_7^+$	3	0
1035'	27	0	0	-1	-1	-1	1	0	1	0	0	0	-1	-1	0	0	0	2	3	6	35	0	-1	-1	3	0
231	7	-3	1	-1	0	0	-1	1	0	e_{15}^+	e_{15}^-	0	0	1	1	1	0	0	3	0	-9	1	0	0	-1	-1
231	7	-3	1	-1	0	0	-1	1	0	e_{15}^-	e_{15}^+	0	0	1	1	1	0	0	3	0	-9	1	0	0	-1	-1
770	-14	5	0	-2	0	0	0	1	0	0	0	0	0	0	e_{23}^+	e_{23}^-	1	1	-2	-7	10	0	0	0	2	-1
770	-14	5	0	-2	0	0	0	1	0	0	0	0	0	0	e_{23}^-	e_{23}^+	1	1	-2	-7	10	0	0	0	2	-1
483	35	6	-2	3	0	0	-1	2	-1	1	1	0	0	0	0	0	0	0	3	0	-2	0	0	0	3	0
1265	49	5	0	1	-2	-2	1	1	0	0	0	0	0	0	0	0	0	0	-3	8	-15	0	1	1	-7	-1
2024	8	-1	-1	0	1	1	0	-1	0	-1	-1	1	1	0	0	0	0	0	0	8	24	-1	1	1	8	-1
2277	21	0	-3	1	2	2	-1	0	0	0	0	0	0	0	0	0	0	2	-3	6	-19	1	-1	-1	-3	0
3312	48	0	-3	0	1	1	0	0	1	0	0	0	-1	-1	0	0	0	-2	0	-6	16	1	1	1	0	0
5313	49	-15	3	-3	0	0	-1	1	0	0	0	0	0	0	0	0	0	0	-3	0	9	-1	0	0	1	1
5796	-28	-9	1	4	0	0	0	-1	-1	1	1	0	0	0	0	0	0	0	0	0	36	1	0	0	-4	-1
5544	-56	9	-1	0	0	0	0	1	0	-1	-1	0	0	1	1	1	0	0	0	0	24	-1	0	0	-8	1
10395	-21	0	0	-1	0	0	1	0	0	0	0	0	0	0	-1	-1	0	0	3	0	-45	0	0	0	3	0

Table 2. Character table for M_{24} .

n	1A	2A	3A	5A	4B	7A	8A	6A	11A	15A	14A	23A	12B	6B	4C	3B	2B	10A	21A	4A	12A
1	90	-6	0	0	2	-1	-2	0	2	0	1	-2	2	-2	2	6	10	0	-1	-6	0
2	462	14	-6	2	-2	0	-2	2	0	-1	0	2	0	0	6	0	-18	2	0	-2	-2
3	1540	-28	10	0	-4	0	0	2	0	0	0	-1	2	2	-4	-14	20	0	0	4	-2
4	4554	42	0	-6	2	4	-2	0	0	0	0	0	0	4	-6	12	-38	2	-2	-6	0
5	11592	-56	-18	2	8	0	0	-2	-2	2	0	0	0	0	0	0	72	2	0	-8	-2
6	27830	86	20	0	-2	-2	2	-4	0	0	2	0	0	0	6	-16	-90	0	-2	6	0
7	61686	-138	0	6	-10	2	-2	0	-2	0	2	0	-2	-2	-2	30	118	-2	2	6	0
8	131100	188	-30	0	4	-3	0	2	2	0	-1	0	0	0	-12	0	-180	0	0	-4	2
9	265650	-238	42	-10	10	0	-2	2	0	2	0	0	-2	6	10	-42	258	-2	0	-14	-2
10	521136	336	0	6	-8	0	-4	0	0	0	0	2	-2	2	16	42	-352	-2	0	0	0
11	988770	-478	-60	0	-14	6	2	-4	2	0	-2	0	0	0	-6	0	450	0	0	18	0
12	1830248	616	62	8	8	0	0	-2	2	2	0	0	2	-6	-16	-70	-600	0	0	-8	-2
13	3303630	-786	0	0	22	-6	2	0	0	0	-2	2	0	-4	6	84	830	0	0	-18	0
14	5844762	1050	-90	-18	-6	0	2	6	0	0	0	2	0	0	18	0	-1062	-2	0	10	-2
15	10139734	-1386	118	4	-26	-4	-2	6	0	-2	0	0	2	2	-10	-110	1334	4	2	22	-2
16	17301060	1764	0	0	12	0	0	0	-4	0	0	0	2	6	-28	126	-1740	0	0	-12	0
17	29051484	-2212	-156	14	28	0	-4	-4	0	-1	0	0	0	0	12	0	2268	-2	0	-36	0
18	48106430	2814	170	0	-18	8	-2	-6	-2	0	0	-2	2	-6	38	-166	-2850	0	2	14	2
19	78599556	-3612	0	-24	-36	0	0	0	2	0	0	0	-2	-6	-20	210	3540	0	0	36	0
20	126894174	4510	-228	14	14	-6	-2	4	0	2	2	0	0	0	-42	0	-4482	-2	0	-18	0
21	202537080	-5544	270	0	48	4	4	6	-2	0	0	0	-2	6	16	-282	5640	0	-2	-40	2
22	319927608	6936	0	18	-16	-7	4	0	0	0	-1	0	0	4	48	300	-6968	2	-1	24	0
23	500376870	-8666	-360	0	-58	0	-2	-8	4	0	0	2	0	0	-18	0	8550	0	0	54	0
24	775492564	10612	400	-36	28	0	0	-8	0	0	0	0	0	-8	-60	-392	-10556	4	0	-28	-4
25	1191453912	-12936	0	12	64	12	-4	0	0	0	0	0	2	-10	32	462	13064	4	0	-72	0
26	1815754710	15862	-510	0	-34	0	-6	10	0	0	0	-1	0	0	78	0	-15930	0	0	22	-2
27	2745870180	-19420	600	30	-76	-10	4	8	-2	0	-2	0	0	8	-36	-600	19268	-2	2	84	0
28	4122417420	23532	0	0	36	2	0	0	0	0	-2	0	0	12	-84	660	-23460	0	2	-36	0
29	6146311620	-28348	-762	-50	100	-6	4	-10	-2	-2	2	0	0	0	36	0	28548	-2	0	-92	-2
30	9104078592	34272	828	22	-40	0	4	-12	4	-2	0	0	0	-8	96	-840	-34352	-2	0	48	0
31	13401053820	-41412	0	0	-116	0	-4	0	0	0	0	-2	-2	-10	-44	966	41180	0	0	108	0
32	19609321554	49618	-1062	34	50	18	2	10	-2	-2	2	0	0	0	-126	0	-49518	2	0	-46	2
33	28530824630	-59178	1220	0	126	0	-6	12	0	0	0	2	-4	12	62	-1204	59430	0	2	-138	0
34	41286761478	70758	0	-72	-66	-10	-6	0	6	0	2	0	0	12	150	1332	-70890	0	2	54	0
35	59435554926	-84530	-1518	26	-154	6	2	-14	0	2	2	0	0	0	-66	0	84222	2	0	158	2
36	85137361430	100310	1670	0	70	-12	-2	-10	0	0	0	0	-2	-18	-170	-1666	-100170	0	0	-74	-2

Table 3. Coefficients in the q-expansion of Z_g for various conjugacy classes g .

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Witten, *Phases of $N = 2$ theories in two-dimensions*, *Nucl. Phys. B* **403** (1993) 159 [[hep-th/9301042](#)] [[INSPIRE](#)].
- [2] J. Distler and S. Kachru, *$(0,2)$ Landau-Ginzburg theory*, *Nucl. Phys. B* **413** (1994) 213 [[hep-th/9309110](#)] [[INSPIRE](#)].
- [3] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic genera of two-dimensional $N = 2$ gauge theories with rank-one gauge groups*, [arXiv:1305.0533](#) [[INSPIRE](#)].
- [4] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic genera of 2d $N = 2$ gauge theories*, [arXiv:1308.4896](#) [[INSPIRE](#)].
- [5] T. Kawai and K. Mohri, *Geometry of $(0,2)$ Landau-Ginzburg orbifolds*, *Nucl. Phys. B* **425** (1994) 191 [[hep-th/9402148](#)] [[INSPIRE](#)].
- [6] T. Eguchi, H. Ooguri and Y. Tachikawa, *Notes on the $K3$ Surface and the Mathieu group M_{24}* , *Exper. Math.* **20** (2011) 91 [[arXiv:1004.0956](#)] [[INSPIRE](#)].
- [7] M.C.N. Cheng, *$K3$ Surfaces, $N = 4$ Dyons and the Mathieu Group M_{24}* , *Commun. Num. Theor. Phys.* **4** (2010) 623 [[arXiv:1005.5415](#)] [[INSPIRE](#)].
- [8] M.R. Gaberdiel, S. Hohenegger and R. Volpato, *Mathieu twining characters for $K3$* , *JHEP* **09** (2010) 058 [[arXiv:1006.0221](#)] [[INSPIRE](#)].
- [9] M.R. Gaberdiel, S. Hohenegger and R. Volpato, *Mathieu Moonshine in the elliptic genus of $K3$* , *JHEP* **10** (2010) 062 [[arXiv:1008.3778](#)] [[INSPIRE](#)].
- [10] T. Eguchi and K. Hikami, *Note on Twisted Elliptic Genus of $K3$ Surface*, *Phys. Lett. B* **694** (2011) 446 [[arXiv:1008.4924](#)] [[INSPIRE](#)].
- [11] T. Gannon, *Much ado about Mathieu*, [arXiv:1211.5531](#) [[INSPIRE](#)].
- [12] M.C.N. Cheng, X. Dong, J. Duncan, J. Harvey, S. Kachru and T. Wrase, *Mathieu Moonshine and $N=2$ String Compactifications*, [arXiv:1306.4981](#) [[INSPIRE](#)].
- [13] D. Whalen et al, to appear.
- [14] J. Distler, *Notes on $(0,2)$ superconformal field theories*, [hep-th/9502012](#) [[INSPIRE](#)].
- [15] M.R. Gaberdiel, S. Hohenegger and R. Volpato, *Symmetries of $K3$ σ -models*, *Commun. Num. Theor. Phys.* **6** (2012) 1 [[arXiv:1106.4315](#)] [[INSPIRE](#)].
- [16] A.N. Schellekens and N.P. Warner, *Anomalies and Modular Invariance in String Theory*, *Phys. Lett. B* **177** (1986) 317 [[INSPIRE](#)].
- [17] A.N. Schellekens and N.P. Warner, *Anomaly Cancellation and Selfdual Lattices*, *Phys. Lett. B* **181** (1986) 339 [[INSPIRE](#)].
- [18] K. Pilch, A.N. Schellekens and N.P. Warner, *Path Integral Calculation of String Anomalies*, *Nucl. Phys. B* **287** (1987) 362 [[INSPIRE](#)].
- [19] E. Witten, *Elliptic Genera and Quantum Field Theory*, *Commun. Math. Phys.* **109** (1987) 525 [[INSPIRE](#)].
- [20] T. Eguchi, H. Ooguri, A. Taormina and S.-K. Yang, *Superconformal Algebras and String Compactification on Manifolds with $SU(N)$ Holonomy*, *Nucl. Phys. B* **315** (1989) 193 [[INSPIRE](#)].

- [21] E. Witten, *On the Landau-Ginzburg description of $N = 2$ minimal models*, *Int. J. Mod. Phys. A* **9** (1994) 4783 [[hep-th/9304026](#)] [[INSPIRE](#)].
- [22] A. Gadde and S. Gukov, *2d Index and Surface operators*, [arXiv:1305.0266](#) [[INSPIRE](#)].
- [23] S. Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, *Invent. Math.* **94** (1988) 183.
- [24] S. Kondo, *Niemeier lattices, Mathieu groups and finite groups of symplectic automorphisms of K3 surfaces*, *Duke Math. J.* **92** (1998) 598.
- [25] B.R. Greene, M.R. Plesser and S.S. Roan, *New constructions of mirror manifolds: Probing moduli space far from Fermat points*, *Mirror symmetry I*, S.T. Yau ed., pg. 347.
- [26] M. Reid, *Canonical 3-folds*, J. de Geometrie algebrique d'Angers and A. Beauville ed., Sijthoff and Noordhoff, Alphen aan den Rijn, 1980, pg. 273.
- [27] A. Taormina and K. Wendland, *The overarching finite symmetry group of Kummer surfaces in the Mathieu group M_{24}* , *JHEP* **08** (2013) 125 [[arXiv:1107.3834](#)] [[INSPIRE](#)].
- [28] A. Taormina and K. Wendland, *Symmetry-surfing the moduli space of Kummer K3s*, [arXiv:1303.2931](#) [[INSPIRE](#)].
- [29] M.C.N. Cheng and J.F.R. Duncan, *Rademacher Sums and Rademacher Series*, [arXiv:1210.3066](#) [[INSPIRE](#)].
- [30] J.A. Harvey and S. Murthy, *Moonshine in Fivebrane Spacetimes*, *JHEP* **01** (2014) 146 [[arXiv:1307.7717](#)] [[INSPIRE](#)].
- [31] S. Govindarajan, *Brewing Moonshine for Mathieu*, [arXiv:1012.5732](#) [[INSPIRE](#)].
- [32] S. Govindarajan, *Unravelling Mathieu Moonshine*, *Nucl. Phys. B* **864** (2012) 823 [[arXiv:1106.5715](#)] [[INSPIRE](#)].
- [33] Y.-H. He and J. McKay, *Eta Products, BPS States and K3 Surfaces*, *JHEP* **01** (2014) 113 [[arXiv:1308.5233](#)] [[INSPIRE](#)].
- [34] T. Eguchi and A. Taormina, *Unitary Representations of $N = 4$ Superconformal Algebra*, *Phys. Lett. B* **196** (1987) 75 [[INSPIRE](#)].
- [35] T. Eguchi and A. Taormina, *Character Formulas for the $N = 4$ Superconformal Algebra*, *Phys. Lett. B* **200** (1988) 315 [[INSPIRE](#)].