RECEIVED: January 19, 2015

REVISED: March 9, 2015

ACCEPTED: March 19, 2015

PUBLISHED: April 10, 2015

Flavour dependent gauged radiative neutrino mass model

Seungwon Baek,^a Hiroshi Okada^a and Kei Yagyu^b

^a*School of Physics, KIAS,
Seoul 130-722, Korea*

^b*School of Physics and Astronomy, University of Southampton,
Southampton, SO17 1BJ, U.K.*

E-mail: swbaek@kias.re.kr, hokada@kias.re.kr, K.Yagyu@soton.ac.uk

ABSTRACT: We propose a one-loop induced radiative neutrino mass model with anomaly free flavour dependent gauge symmetry: μ minus τ symmetry $U(1)_{\mu-\tau}$. A neutrino mass matrix satisfying current experimental data can be obtained by introducing a weak isospin singlet scalar boson that breaks $U(1)_{\mu-\tau}$ symmetry, an inert doublet scalar field, and three right-handed neutrinos in addition to the fields in the standard model. We find that a characteristic structure appears in the neutrino mass matrix: two-zero texture form which predicts three non-zero neutrino masses and three non-zero CP-phases from five well measured experimental inputs of two squared mass differences and three mixing angles. Furthermore, it is clarified that only the inverted mass hierarchy is allowed in our model. In a favored parameter set from the neutrino sector, the discrepancy in the muon anomalous magnetic moment between the experimental data and the standard model prediction can be explained by the additional neutral gauge boson loop contribution with mass of order 100 MeV and new gauge coupling of order 10^{-3} .

KEYWORDS: Beyond Standard Model, Neutrino Physics

ARXIV EPRINT: [1501.01530](https://arxiv.org/abs/1501.01530)

Contents

1	Introduction	1
2	The model	2
3	Lepton mass matrix	4
4	Muon anomalous magnetic moment and lepton flavour violation	8
5	Conclusions and discussions	9

1 Introduction

Radiative neutrino mass models are one of the most promising scenarios at TeV scale physics to explain tiny neutrino masses. The original model based on the idea of radiative generation of neutrino masses is known as the Zee model [1] proposed in early 80's, where neutrino masses are generated at the one-loop level. After the Zee model, the Zee-Babu model [2–7] has also been proposed, where neutrino masses are explained at the two-loop level. In 2000's, radiative neutrino mass models have been extended so as to include a dark matter (DM) candidate by introducing an unbroken symmetry such as a discrete Z_2 symmetry known as; e.g., the models by Krauss-Nasri-Trodden [8, 9] and by Ma [10–17]. After these models appeared, various kinds of extensions have been considered in the scenario based on the radiative neutrino mass generation such as models with the supersymmetry [18–21], the B-L symmetry [22–29], flavour symmetries [30–34] and the conformal symmetry [35, 36]. Furthermore, in refs. [37–39], a complex $SU(2)_L$ triplet scalar field is introduced, in which the collider phenomenology can be rich because of the existence of doubly-charged scalar bosons. Loop induced Dirac type neutrino masses have been proposed in refs. [40–44]. In models proposed in refs. [45–49], charged lepton masses are also introduced at quantum levels in addition to neutrino masses. In addition to the above mentioned models, a number of radiative neutrino mass models have been constructed [50–68] up to now, and they have been classified into some groups in refs. [69–74].

On the other hand, Abelian gauged $U(1)$ symmetries are well compatible with such radiative models. It has been known that there are four different anomaly free and flavour dependent types of $U(1)$ symmetries in the leptonic sector; namely, $L_e - L_\mu$, $L_e - L_\tau$, and $L_\mu - L_\tau$, where L_i denotes the lepton number with the flavour i . Especially in the case of $L_\mu - L_\tau$ [75–85], constraints on the mass of additional neutral gauge boson Z' and the new gauge coupling constant from the LEP experiment are very weak, because the Z' boson does not couple directly to the electron. We thus can consider a light Z' boson scenario, by which the discrepancy in the muon anomalous magnetic moment between current data

	Lepton Fields			Scalar Fields		
	$L_L^i = (\nu_L^i, e_L^i)^T$	e_R^i	N_R^i	Φ	η	S
$SU(2)_L$	2	1	1	2	2	1
$U(1)_Y$	$-1/2$	-1	0	$+1/2$	$+1/2$	0
Z_2	$+$	$+$	$-$	$+$	$-$	$+$

Table 1. The charge assignments of leptons and scalars under $SU(2)_L \times U(1)_Y$ and Z_2 symmetry. The index $i(= e, \mu, \tau)$ denotes the lepton flavour.

	(L_L^e, e_R, N_R^e)	$(L_L^\mu, \mu_R, N_R^\mu)$	$(L_L^\tau, \tau_R, N_R^\tau)$	S
$U(1)_{\mu-\tau}$	0	$+1$	-1	$+1$

Table 2. The charge assignments under the gauged $U(1)_{\mu-\tau}$ symmetry. Fields which are not displayed in this table are neutral under $U(1)_{\mu-\tau}$.

and the prediction in the standard model (SM) [75] can be explained with the mass of Z' to be $\mathcal{O}(100)$ MeV and the $U(1)_{\mu-\tau}$ gauge coupling to be $\mathcal{O}(10^{-3})$. The positron anomaly reported by AMS-02 [86] could be explained [77, 87]. Such a light Z' boson can be probed at the 14 TeV run of the LHC [84] through multi-lepton signals.

In our paper, we combine a radiative neutrino mass model at one-loop level and the gauged $U(1)_{\mu-\tau}$ symmetry to get neutrino masses, mixings, and dark matter candidates. We find that a predictive two-zero texture form of a neutrino mass matrix can be obtained corresponding to “Pattern C” in ref. [88]. In this texture, we only need five experimental inputs to determine all the neutrino parameters. We can choose the most accurately measured ones: two squared mass differences and three mixing angles. It turns out that only the inverted mass hierarchy is allowed in our texture. Non-vanishing one Dirac and two Majorana CP-phases, and non-zero three neutrino mass eigenvalues are predicted.

This paper is organized as follows. In section II, we define our model, and give mass formulae for scalar bosons. In section III, we calculate the mass matrices for the lepton sector; charged leptons, right-handed neutrinos and left-handed neutrinos. The detailed analysis for the two-zero texture form of neutrino mass matrix is also discussed. In section IV, we discuss new contributions to the muon $g-2$ and lepton flavour violation in our model. Conclusions and discussions are given in section V.

2 The model

We consider a model in the framework of the gauge symmetry of $SU(2)_L \times U(1)_Y \times U(1)_{\mu-\tau}$ with an unbroken discrete Z_2 symmetry. The particle content in our model is listed in table 1. The charge assignment for the $U(1)_{\mu-\tau}$ symmetry is separately shown in table 2.

Our model is an extension of the model proposed by Ma [10–16], where neutrino masses are generated at the one-loop level. In the Ma model, three right-handed neutrinos and an inert scalar doublet field are added to the standard model (SM). We introduce only one

additional $SU(2)_L$ singlet scalar field S with the even parity under Z_2 to the Ma model. The vacuum expectation value (VEV) of S breaks the $U(1)_{\mu-\tau}$ symmetry.

The mass terms for right-handed neutrinos N_R^i and the relevant Yukawa interactions are given by

$$\begin{aligned}
 -\mathcal{L}_Y = & \frac{1}{2}M_{ee}\overline{N_R^{e^c}}N_R^e + \frac{1}{2}M_{\mu\tau}(\overline{N_R^{\mu^c}}N_R^\tau + \overline{N_R^{\tau^c}}N_R^\mu) + \text{h.c.} \\
 & + y_e\overline{L_L^e}\Phi e_R + y_\mu\overline{L_L^\mu}\Phi\mu_R + y_\tau\overline{L_L^\tau}\Phi\tau_R + \text{h.c.} \\
 & + h_{e\mu}(\overline{N_R^{e^c}}N_R^\mu + \overline{N_R^{\mu^c}}N_R^e)S^* + h_{e\tau}(\overline{N_R^{e^c}}N_R^\tau + \overline{N_R^{\tau^c}}N_R^e)S + \text{h.c.} \\
 & + f_e\overline{L_L^e}(i\sigma_2)\eta^*N_R^e + f_\mu\overline{L_L^\mu}(i\sigma_2)\eta^*N_R^\mu + f_\tau\overline{L_L^\tau}(i\sigma_2)\eta^*N_R^\tau + \text{h.c.} \quad (2.1)
 \end{aligned}$$

The scalar sector of our model is composed of a singlet (S) and two doublets, one active (Φ) and one inert (η). The most general scalar potential is given by

$$\begin{aligned}
 \mathcal{V} = & \mu_\Phi^2|\Phi|^2 + \mu_\eta^2|\eta|^2 + \mu_S^2|S|^2 \\
 & + \frac{1}{2}\lambda_1|\Phi|^4 + \frac{1}{2}\lambda_2|\eta|^4 + \lambda_3|\Phi|^2|\eta|^2 + \lambda_4|\Phi^\dagger\eta|^2 + \frac{1}{2}\lambda_5[(\Phi^\dagger\eta)^2 + \text{h.c.}] \\
 & + \lambda_S|S|^4 + \lambda_{S\Phi}|S|^2|\Phi|^2 + \lambda_{S\eta}|S|^2|\eta|^2, \quad (2.2)
 \end{aligned}$$

where all the parameters can be taken to be real without any loss of generality. The scalar fields are parameterized by

$$\Phi = \begin{bmatrix} G^+ \\ \frac{1}{\sqrt{2}}(v + \varphi_H + iG^0) \end{bmatrix}, \quad \eta = \begin{bmatrix} \eta^+ \\ \frac{1}{\sqrt{2}}(\eta_H + i\eta_A) \end{bmatrix}, \quad S = \frac{1}{\sqrt{2}}(v_S + S_H + iG_S), \quad (2.3)$$

where v is the VEV related with the Fermi constant G_F by $v^2 = 1/(\sqrt{2}G_F)$, and v_S is the VEV of S which breaks the $U(1)_{\mu-\tau}$ symmetry. In eq. (2.3), $G^\pm$, G^0 and G_S are the Nambu-Goldstone bosons which are absorbed by the longitudinal component of the $W^\pm$, Z and an extra neutral gauge boson Z' associated with the $U(1)_{\mu-\tau}$ symmetry, respectively.

The tadpole conditions for φ_H and S_H are respectively given by

$$\begin{aligned}
 \left. \frac{\partial \mathcal{V}}{\partial \varphi_H} \right|_0 &= v \left(\mu_\Phi^2 + \frac{v^2}{2}\lambda_1 + \frac{v_S^2}{2}\lambda_{S\Phi} \right) = 0, \\
 \left. \frac{\partial \mathcal{V}}{\partial S_H} \right|_0 &= v_S \left(\mu_S^2 + \frac{v^2}{2}\lambda_{S\Phi} + v_S^2\lambda_S \right) = 0. \quad (2.4)
 \end{aligned}$$

Using the above two equations, we can eliminate μ_Φ^2 and μ_S^2 . There is no tadpole condition for η_H , because the VEV of inert doublet field η is zero due to the unbroken Z_2 symmetry.

The Z_2 -odd component scalar fields, $\eta^\pm$, η_A and η_H , do not mix with the other fields, and their squared masses are simply given by

$$m_{\eta^\pm}^2 = \mu_\eta^2 + \frac{v_S^2}{2}\lambda_{S\eta} + \frac{v^2}{2}\lambda_3, \quad (2.5)$$

$$m_{\eta_A}^2 = \mu_\eta^2 + \frac{v_S^2}{2}\lambda_{S\eta} + \frac{v^2}{2}(\lambda_3 + \lambda_4 - \lambda_5), \quad (2.6)$$

Figure 1. Feynman diagram for neutrino masses at the one-loop level. In the internal fermion line, N_k denotes the mass eigenstate of the right-handed neutrinos.

$$m_{\eta_H}^2 = \mu_\eta^2 + \frac{v_S^2}{2} \lambda_{S\eta} + \frac{v^2}{2} (\lambda_3 + \lambda_4 + \lambda_5). \quad (2.7)$$

For the Z_2 -even sector, two CP-even scalar states φ_H and S_H are mixed with each other. Their mass matrix, $\mathcal{M}_H^2$, in the basis of (φ_H, S_H) is given by

$$\mathcal{M}_H^2 = \begin{pmatrix} v^2 \lambda_1 & v v_S \lambda_{S\Phi} \\ v v_S \lambda_{S\Phi} & 2 v_S^2 \lambda_S \end{pmatrix}. \quad (2.8)$$

The mass eigenstates for the CP-even states are given by introducing the mixing angle α by

$$\begin{pmatrix} \varphi_H \\ S_H \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} h \\ H \end{pmatrix}. \quad (2.9)$$

In terms of the matrix element expressed in eq. (2.8), the mass eigenvalues are

$$m_h^2 = \cos^2 \alpha (\mathcal{M}_H^2)_{11} + \sin^2 \alpha (\mathcal{M}_H^2)_{22} + \sin 2\alpha (\mathcal{M}_H^2)_{12}, \quad (2.10)$$

$$m_H^2 = \sin^2 \alpha (\mathcal{M}_H^2)_{11} + \cos^2 \alpha (\mathcal{M}_H^2)_{22} - \sin 2\alpha (\mathcal{M}_H^2)_{12}, \quad (2.11)$$

and the mixing angle is

$$\tan 2\alpha = \frac{2(\mathcal{M}_H^2)_{12}}{(\mathcal{M}_H^2)_{11} - (\mathcal{M}_H^2)_{22}}. \quad (2.12)$$

We define h as the SM-like Higgs boson with the mass of 126 GeV. Thus, H corresponds to an additional singlet-like Higgs boson. Finally, if the conditions

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_S > 0, \quad (2.13)$$

$$\lambda_{S\Phi} + \frac{1}{\sqrt{2}} \sqrt{\lambda_1 \lambda_S} > 0, \quad \lambda_{S\eta} + \frac{1}{\sqrt{2}} \sqrt{\lambda_2 \lambda_S} > 0, \quad (2.14)$$

$$\lambda_3 + \frac{1}{2} \sqrt{\lambda_1 \lambda_S} + \min(0, \lambda_4 \pm \lambda_5) > 0. \quad (2.15)$$

are satisfied, the Higgs potential eq. (2.2) is bounded from below.

3 Lepton mass matrix

The mass matrices for the charged-leptons and right-handed neutrinos are defined as

$$\begin{aligned} -\mathcal{L}_{\text{mass}} = & (\bar{e}, \bar{\mu}, \bar{\tau}) \mathcal{M}_\ell (e, \mu, \tau)^T \\ & + \frac{1}{2} (\bar{N}_R^{e\bar{c}}, \bar{N}_R^{\mu\bar{c}}, \bar{N}_R^{\tau\bar{c}}) \mathcal{M}_N (N_R^e, N_R^\mu, N_R^\tau)^T + \text{h.c.}, \end{aligned} \quad (3.1)$$

where e , μ and τ are, respectively, $(e_L + e_R)$, $(\mu_L + \mu_R)$ and $(\tau_L + \tau_R)$. After the phase redefinition of the fields, e_R^i and N_R^i , the mass matrices can be written in the form

$$\mathcal{M}_\ell = \frac{v}{\sqrt{2}} \text{diag}(|y_e|, |y_\mu|, |y_\tau|), \quad \mathcal{M}_N = \begin{pmatrix} |M_{ee}| & \frac{v_S}{\sqrt{2}} |h_{e\mu}| & \frac{v_S}{\sqrt{2}} |h_{e\tau}| \\ \frac{v_S}{\sqrt{2}} |h_{e\mu}| & 0 & |M_{\mu\tau}| e^{i\theta_R} \\ \frac{v_S}{\sqrt{2}} |h_{e\tau}| & |M_{\mu\tau}| e^{i\theta_R} & 0 \end{pmatrix}, \quad (3.2)$$

where θ_R is the remaining unremovable phase. Notice here that the $U(1)_{\mu-\tau}$ symmetry predicts the diagonal form of the mass matrix for the charged leptons. The mass matrix $\mathcal{M}_N$ is diagonalized by introducing a unitary matrix V satisfying

$$V^T \mathcal{M}_N V = \mathcal{M}_N^{\text{diag}} \equiv \text{diag}(M_1, M_2, M_3). \quad (3.3)$$

The mass matrix for the left-handed Majorana neutrinos is then calculated to be

$$(\mathcal{M}_\nu)_{ij} = \frac{1}{32\pi^2} \sum_{k=1-3} (f_i V_{ik}) M_{N_k} (f_j V_{jk}) \left(\frac{m_{\eta_H}^2}{M_k^2 - m_{\eta_H}^2} \ln \frac{m_{\eta_H}^2}{M_k^2} - \frac{m_{\eta_A}^2}{M_k^2 - m_{\eta_A}^2} \ln \frac{m_{\eta_A}^2}{M_k^2} \right). \quad (3.4)$$

If we assume $m_0^2 \equiv (m_{\eta_H}^2 + m_{\eta_A}^2)/2 \gg M_k^2$, the neutrino mass matrix can be simplified to be

$$\begin{aligned} (\mathcal{M}_\nu)_{ij} &\simeq -\frac{1}{32\pi^2} \frac{\lambda_5 v^2}{m_0^2} \sum_{k=1-3} (f_i V_{ik}) M_k (f_j V_{jk}) \\ &= -\frac{1}{32\pi^2} \frac{\lambda_5 v^2}{m_0^2} \sum_{k=1-3} f_i (\mathcal{M}_N)_{ij} f_j. \end{aligned} \quad (3.5)$$

More explicitly, $\mathcal{M}_\nu$ can be written as

$$\mathcal{M}_\nu = \begin{pmatrix} f_e^2 M_{11} & f_e f_\mu M_{12} & f_e f_\tau M_{13} \\ f_e f_\mu M_{12} & 0 & f_\mu f_\tau M_{23} e^{i\theta_R} \\ f_e f_\tau M_{13} & f_\mu f_\tau M_{23} e^{i\theta_R} & 0 \end{pmatrix}, \quad (3.6)$$

where we reparametrized dimension-full real parameters M_{ij} defined as

$$M_{11} = M_{ee}, \quad M_{12} = \frac{v_S}{\sqrt{2}} h_{e\mu}, \quad M_{13} = \frac{v_S}{\sqrt{2}} h_{e\tau}, \quad M_{23} = M_{\mu\tau}, \quad (3.7)$$

in the unite of $-\lambda_5 v^2/(32\pi^2 m_0^2)$. The structure of matrix, eq. (3.6), implies that the $U(1)_{\mu-\tau}$ symmetry predicts the so-called two-zero texture form of the Majorana neutrino mass matrix. Fifteen patterns of the two-zero texture form have been discussed in ref. [88], and our form corresponds to one termed ‘‘Pattern C’’. Because of the two zero texture form, nine neutrino parameters, three mass eigenvalues, three mixing angles and three (one Dirac and two Majorana) CP-phases, are predicted from five input parameters. In the following, we’ll discuss how we can determine all the neutrino parameters by five experimental inputs.

First, we introduce the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix U_{PMNS} [89] to diagonalize the neutrino mass matrix:

$$\mathcal{M}_\nu = U_{\text{PMNS}} \text{diag}(m_1, m_2, m_3) U_{\text{PMNS}}^T, \quad (3.8)$$

where m_1 , m_2 and m_3 are the neutrino mass eigenvalues. The PMNS matrix is expressed as the product of two unitary matrices

$$U_{\text{PMNS}} = UP, \quad (3.9)$$

where

$$U \equiv \begin{bmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{bmatrix} \begin{bmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{bmatrix} \begin{bmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad P \equiv \text{diag}(e^{i\rho}, e^{i\sigma}, 1), \quad (3.10)$$

with $s_{ij} = \sin \theta_{ij}$ and $c_{ij} = \cos \theta_{ij}$. In eq. (3.10), δ is the Dirac phase, and ρ and σ are the Majorana phases. Using the matrix U , eq. (3.8) is rewritten by

$$\mathcal{M}_\nu = U \text{diag}(\tilde{m}_1, \tilde{m}_2, \tilde{m}_3) U^T, \quad (3.11)$$

where $\tilde{m}_3 = m_3 e^{2i\rho}$, $\tilde{m}_2 = m_2 e^{2i\sigma}$ and $\tilde{m}_1 = m_1$.

Second, we obtain the following two equations from the two-zero texture form

$$[U \text{diag}(\tilde{m}_1, \tilde{m}_2, \tilde{m}_3) U^T]_{22} = [U \text{diag}(\tilde{m}_1, \tilde{m}_2, \tilde{m}_3) U^T]_{33} = 0. \quad (3.12)$$

This gives [88]

$$\begin{aligned} \frac{\tilde{m}_1}{\tilde{m}_3} &= \frac{c_{12}c_{13}^2}{s_{13}} \frac{c_{12}(c_{23}^2 - s_{23}^2)e^{i\delta} - 2s_{12}s_{23}s_{23}c_{23}}{2s_{12}c_{12}s_{23}c_{23}(e^{2i\delta} + s_{13}^2) - s_{13}(c_{12}^2 - s_{12}^2)(c_{23}^2 - s_{23}^2)e^{i\delta}} e^{2i\delta}, \\ \frac{\tilde{m}_2}{\tilde{m}_3} &= -\frac{s_{12}c_{13}^2}{s_{13}} \frac{s_{12}(c_{23}^2 - s_{23}^2)e^{i\delta} - 2s_{12}s_{23}s_{23}c_{23}}{2s_{12}c_{12}s_{23}c_{23}(e^{2i\delta} + s_{13}^2) - s_{13}(c_{12}^2 - s_{12}^2)(c_{23}^2 - s_{23}^2)e^{i\delta}} e^{2i\delta}. \end{aligned} \quad (3.13)$$

The ratios of neutrino mass eigenvalues and the Majorana phases are obtained from eq. (3.13) as

$$R_{13} \equiv \frac{m_1}{m_3} = \left| \frac{\tilde{m}_1}{\tilde{m}_3} \right|, \quad R_{23} \equiv \frac{m_2}{m_3} = \left| \frac{\tilde{m}_2}{\tilde{m}_3} \right|, \quad \rho = \frac{1}{2} \arg \left[\frac{\tilde{m}_1}{\tilde{m}_3} \right], \quad \sigma = \frac{1}{2} \arg \left[\frac{\tilde{m}_2}{\tilde{m}_3} \right]. \quad (3.14)$$

Using $0 \leq \theta_{ij} < \pi/2$ ($ij = 12, 13$, and 23) and $\theta_{13} \ll 1$, we obtain approximate formulae for R_{13} and R_{23} as

$$\begin{aligned} R_{13} &\simeq \left[1 - \frac{2 \cot \theta_{12}}{\sin \theta_{13}} \cot 2\theta_{23} \cos \delta + \left(\frac{\cot \theta_{12}}{\sin \theta_{13}} \cot 2\theta_{23} \right)^2 \right]^{1/2}, \\ R_{23} &\simeq \left[1 + \frac{2 \tan \theta_{12}}{\sin \theta_{13}} \cot 2\theta_{23} \cos \delta + \left(\frac{\tan \theta_{12}}{\sin \theta_{13}} \cot 2\theta_{23} \right)^2 \right]^{1/2}. \end{aligned} \quad (3.15)$$

In order to guarantee $m_2 > m_1$ (i.e., $R_{23} > R_{13}$), we require $\cot 2\theta_{23} \cos \delta > 0$. In that case, we obtain $R_{23} > 1$, which shows that only the inverted mass hierarchy ($m_2 > m_1 > m_3$) is allowed in our model as already mentioned in ref. [88].

Finally, we define the ratio of two squared mass difference:

$$R_\nu \equiv \frac{\Delta m_{21}^2}{|\Delta m_{31}^2|} = \frac{m_2^2 - m_1^2}{|m_3^2 - m_1^2|}. \quad (3.16)$$

From eq. (3.14), it can be rewritten in the inverted mass hierarchy as

$$R_\nu = \frac{R_{23}^2 - R_{13}^2}{R_{13}^2 - 1} \simeq \frac{2}{\cos^2 \theta_{12}} \frac{\cot 2\theta_{12} \cot 2\theta_{23} - \sin \theta_{13} \cos \delta}{2 \sin \theta_{13} \cos \delta - \cot \theta_{12} \cot 2\theta_{23}}. \quad (3.17)$$

We can obtain three mass eigenvalues in terms of Δm_{21}^2 , R_{13} and R_{23} as

$$m_3 = \frac{\sqrt{\Delta m_{21}^2}}{\sqrt{R_{23}^2 - R_{13}^2}}, \quad m_1 = m_3 R_{31}, \quad m_2 = m_3 R_{23}. \quad (3.18)$$

Now, we are ready to determine all the neutrino parameters by using five experimental inputs. The best fit (3σ range) values in the inverted mass hierarchy are given as follows [90]:

$$\begin{aligned} s_{12}^2 &= 0.323 \text{ (0.278-0.375)}, \quad s_{23}^2 = 0.573 \text{ (0.403-0.640)}, \quad s_{13}^2 = 0.0229 \text{ (0.0193-0.0265)}, \\ \Delta m_{21}^2 &= 7.60 \text{ (7.11-8.18)} \times 10^{-5} \text{ eV}^2, \quad |\Delta m_{31}^2| = 2.38 \text{ (2.20-2.54)} \times 10^{-3} \text{ eV}^2, \end{aligned} \quad (3.19)$$

In the following, we present our predictions using the three sets of input parameters; namely, using the best fit values (BF), using the upper limit of the 3σ range ($+3\sigma$) and using the lower limit of the 3σ range (-3σ). From two squared mass differences, we can obtain the numerical value

$$R_\nu = 0.0319 \text{ (BF)}, \quad 0.0322 \text{ (+3}\sigma\text{)}, \quad 0.0323 \text{ (-3}\sigma\text{)}, \quad (3.20)$$

from eq. (3.16). We can see that the analytic formula of R_ν in eq. (3.17) is a function of δ . From eq. (3.17) and eq. (3.20), we obtain the Dirac phase

$$\delta = \pm 1.96 \text{ (BF)}, \quad \pm 2.07 \text{ (+3}\sigma\text{)}, \quad \pm 0.774 \text{ (-3}\sigma\text{)}. \quad (3.21)$$

All the negative (positive) solutions for δ are allowed (excluded) by the experimental data at 95% CL [90], so that we choose the negative solution. We then obtain the ratios as

$$\begin{aligned} \frac{\tilde{m}_1}{\tilde{m}_3} &= 1.39 \times e^{1.91i}, & \frac{\tilde{m}_2}{\tilde{m}_3} &= 1.40 \times e^{-2.68i} \text{ (BF)}, \\ \frac{\tilde{m}_1}{\tilde{m}_3} &= 2.03 \times e^{1.52i}, & \frac{\tilde{m}_2}{\tilde{m}_3} &= 2.06 \times e^{-2.51i} \text{ (+3}\sigma\text{)}, \\ \frac{\tilde{m}_1}{\tilde{m}_3} &= 1.73 \times e^{-1.19i}, & \frac{\tilde{m}_2}{\tilde{m}_3} &= 1.74 \times e^{2.78i} \text{ (-3}\sigma\text{)}, \end{aligned} \quad (3.22)$$

and the mass eigenvalues and Majorana phases from eqs. (3.14) and (3.18)

$$\begin{aligned} (m_1, m_2, m_3) \text{ [eV]} &= 0.0583, 0.0589, 0.0420, \quad (\rho, \sigma) = (0.956, -1.34) \text{ (BF)}, \\ (m_1, m_2, m_3) \text{ [eV]} &= 0.0533, 0.0540, 0.0262, \quad (\rho, \sigma) = (0.759, -1.25) \text{ (+3}\sigma\text{)}, \\ (m_1, m_2, m_3) \text{ [eV]} &= 0.0585, 0.0591, 0.0339, \quad (\rho, \sigma) = (-0.596, 1.39) \text{ (-3}\sigma\text{)}. \end{aligned} \quad (3.23)$$

Using eq. (3.8), we can get the neutrino mass matrix M_ν

$$\mathcal{M}_\nu \simeq \begin{pmatrix} 0.0408 & -0.0186 & -0.0378 \\ -0.0186 & 0 & -0.0420 - 0.00631i \\ -0.0378 & -0.0420 - 0.00631i & 0 \end{pmatrix} \text{ eV} \text{ (BF)},$$

$$\begin{aligned}\mathcal{M}_\nu &\simeq \begin{pmatrix} 0.0249 & -0.0228 & -0.0410 \\ -0.0228 & 0 & -0.0270 - 0.00352i \\ -0.0410 & -0.0270 - 0.00352i & 0 \end{pmatrix} \text{ eV} \quad (+3\sigma), \\ \mathcal{M}_\nu &\simeq \begin{pmatrix} 0.0321 & 0.0399 & 0.0271 \\ 0.0399 & 0 & -0.0344 + 0.00252i \\ 0.0271 & -0.0344 + 0.00252i & 0 \end{pmatrix} \text{ eV} \quad (-3\sigma),\end{aligned}\quad (3.24)$$

where we performed a phase redefinition so that the phase appears in the $(2, 3)$ -component as in eq. (3.6). We thus determine our model parameters by comparing each element of the above matrix with corresponding one given in eq. (3.6). We note in passing that the lightest right-handed neutrino can be a DM candidate in our scenario discussed in this section. The phenomenology of fermionic DM is quite similar to that in the Ma's model [10–16], and its detailed discussions have been presented in ref. [17].

4 Muon anomalous magnetic moment and lepton flavour violation

The muon anomalous magnetic moment, so-called the muon $g-2$, has been measured at Brookhaven National Laboratory. The current average of the experimental results is given by [91]

$$a_\mu^{\text{exp}} = 11659208.0(6.3) \times 10^{-10}. \quad (4.1)$$

It has been known that there is a discrepancy from the SM prediction by 3.2σ [92] to 4.1σ [93]:

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (29.0 \pm 9.0 \text{ to } 33.5 \pm 8.2) \times 10^{-10}. \quad (4.2)$$

In our model, the dominant contribution to the muon $g-2$ is obtained through the one loop diagram where the muon and the extra neutral gauge boson Z' of the $U(1)_{\mu-\tau}$ symmetry are running in the loop. The resulting form is given by

$$\Delta a_\mu(Z') = \frac{g_{Z'}^2}{8\pi^2} \int_0^1 dx \frac{2rx(1-x)^2}{r(1-x)^2 + x}, \quad (4.3)$$

where $g_{Z'}$ and $m_{Z'}$ are the $U(1)_{\mu-\tau}$ gauge coupling constant, the mass of Z' , respectively, and $r \equiv (m_\mu/m_{Z'})^2$.

On the other hand, the parameter space on $m_{Z'}$ and $g_{Z'}$ has been severely constrained by the neutrino trident production process [95, 96] observed in neutrino beam experiments at the CHARMII [97] and at the CCFR [98], whose measured cross section well agrees with the SM prediction. For example, $g_{Z'} \gtrsim 0.1$, $g_{Z'} \gtrsim 0.02$, $g_{Z'} \gtrsim 0.002$ and $g_{Z'} \gtrsim 0.001$ have been excluded with 95% CL in the cases of $m_{Z'} = 100, 10, 1$ and 0.1 GeV , respectively [95, 96]. However, we note that the muon ($g-2$) in our model is not constrained by the dark photon search experiment at BaBar because Z' does not couple to the electron in our case [99, 100].

By taking into account the constraint from the neutrino trident production, the discrepancy in the muon $g - 2$ can be compensated to be less than 2σ by $m_{Z'} \simeq 200 \text{ MeV}$ with $g_{Z'} \simeq 10^{-3}$ ¹

The lepton flavor violation also arises through the $\eta^\pm$ loop in our model. The most stringent constraint is imposed by the MEG experiment: $\mathcal{B}(\mu \rightarrow e\gamma) < 5.7 \times 10^{-13}$ [94]. The branching fraction is written by

$$\mathcal{B}(\mu \rightarrow e\gamma) \simeq (900 \text{ GeV}^2)^2 \times \left| \sum_{i=1-3} \frac{f_e f_\mu}{2m_{\eta^\pm}^2} V_{1i} V_{2i}^* G \left(\frac{M_i^2}{m_{\eta^\pm}^2} \right) \right|^2. \quad (4.4)$$

If we take $\sum_{i=1-3} f_e f_\mu V_{1i} V_{2i}^* \lesssim \mathcal{O}(10^{-3})$ with $m_{\eta^\pm} = \mathcal{O}(1) \text{ TeV}$, we can avoid this constraint. Therefore, the anomaly in the muon $g - 2$ can be well explained in the favored parameter region suggested from neutrino data and lepton flavour violation data.

We note that our Z' boson does not couple to the SM quarks, because it appears from the $U(1)_{\mu-\tau}$ gauge symmetry; i.e., it has a *quark phobic* nature. Therefore, any constraints for the Z' boson using hadron events such as dijet searches cannot be applied.

5 Conclusions and discussions

We have constructed a one-loop induced radiative neutrino mass model in the gauge symmetry $SU(2)_L \times U(1)_Y \times U(1)_{\mu-\tau}$ with the unbroken discrete Z_2 symmetry. In our model, three right-handed neutrinos are introduced in addition to the SM, and the scalar sector is composed of two isospin doublets, one inert and one active, and a $U(1)_{\mu-\tau}$ charged singlet.

We have shown that the $U(1)_{\mu-\tau}$ symmetry predicts a characteristic structure of the lepton mass matrices. First, the mass matrix of charged leptons is diagonal in the interaction basis. Second, the mass matrix of left-handed neutrinos is in the two-zero texture form if inert scalar bosons are much heavier than the right-handed neutrinos. The two-zero texture form of the neutrino mass matrix has been intensively studied in ref. [88], and our model provides a texture with vanishing (2,2) and (3,3) elements, corresponding to “Pattern C” in [88]. In this pattern, only the inverted mass hierarchy is allowed. And we only need five input experimental data to fix the neutrino mass matrix. We can choose the most accurately measured ones: two squared mass differences and three mixing angles. Using the best fit values of five observables, we obtained non-zero Dirac and Majorana CP-phases, and non-zero three neutrino mass eigenvalues.

We showed that the Z' -loop contribution to the muon $g - 2$ can explain the discrepancy between the current experimental data and the SM prediction if the Z' mass is of $\mathcal{O}(100) \text{ MeV}$ and the $U(1)_{\mu-\tau}$ gauge coupling of $\mathcal{O}(10^{-3})$, which has not been excluded by the neutrino trident production process. The constraint from lepton flavour violation such as $\mu \rightarrow e\gamma$ can be avoided in the parameter space favored by the neutrino data and the muon $g - 2$.

¹In addition to the Z' loop contribution, there is a negative contribution to the muon $g - 2$ from the $\eta^\pm$ and N_i ($i=1-3$) loop diagram. However, it can be neglected due to the assumption $M_k^2 \ll m_{\eta^\pm}^2$ that provides two-zero texture in the neutrino sector.

Finally, we would like to briefly discuss the collider phenomenology of our model. One of the important features is that the SM-like Higgs boson h which was discovered at the LHC [101, 102] can mix with the $U(1)_{\mu-\tau}$ Higgs boson H via the mixing angle α . As a consequence, the coupling constants of h with the SM gauge bosons hVV ($V = W, Z$) and fermions $hf\bar{f}$ can be universally deviated from those of the SM predictions by the factor $\cos\alpha$. Since the pattern of the deviation in the h couplings strongly depends on the structure of the Higgs sector as discussed in ref. [103], we can indirectly probe the model by looking at the deviation even if we cannot discover new particles such as H . In future collider experiments such as the LHC Run-II, the high luminosity LHC and the International Linear Collider (ILC), the Higgs boson couplings are expected to be measured quite accurately, especially they can be measured. In particular at the ILC with the collision energy of 500 GeV and the integrated luminosity of 500 fb $^{-1}$, the hVV and $hf\bar{f}$ ($f = b, \tau$ and t) couplings can be measured with about 0.4% and $\mathcal{O}(1)\%$ [104, 105], respectively. Therefore, we can test our model by the comparison between the precisely measured Higgs boson coupling and the theory predictions.

Acknowledgments

The authors would like to thank W. Altmannshofer, T. Araki, J. Heeck and M. Kohda for a useful comment on the neutrino trident production related to the Z' contribution to the muon $g - 2$. This work was supported in part by National Research Foundation of Korea (NRF) Research Grant 2012R1A2A1A01006053 (SB). KY was supported by JSPS postdoctoral fellowships for research abroad.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Zee, *A theory of lepton number violation, neutrino Majorana mass and oscillation*, *Phys. Lett. B* **93** (1980) 389 [Erratum *ibid.* **B 95** (1980) 461] [[INSPIRE](#)].
- [2] A. Zee, *Quantum numbers of Majorana neutrino masses*, *Nucl. Phys. B* **264** (1986) 99 [[INSPIRE](#)].
- [3] K.S. Babu, *Model of ‘calculable’ Majorana neutrino masses*, *Phys. Lett. B* **203** (1988) 132 [[INSPIRE](#)].
- [4] S. Baek, P. Ko, H. Okada and E. Senaha, *Can Zee-Babu model implemented with scalar dark matter explain both Fermi/LAT 130 GeV γ -ray excess and neutrino physics?*, *JHEP* **09** (2014) 153 [[arXiv:1209.1685](#)] [[INSPIRE](#)].
- [5] D. Schmidt, T. Schwetz and H. Zhang, *Status of the Zee-Babu model for neutrino mass and possible tests at a like-sign linear collider*, *Nucl. Phys. B* **885** (2014) 524 [[arXiv:1402.2251](#)] [[INSPIRE](#)].
- [6] H. Okada, T. Toma and K. Yagyu, *Inert extension of the Zee-Babu model*, *Phys. Rev. D* **90** (2014) 095005 [[arXiv:1408.0961](#)] [[INSPIRE](#)].

- [7] S. Baek, *3.5 keV X-ray line signal from dark matter decay in local $U(1)_{B-L}$ extension of Zee-Babu model*, [arXiv:1410.1992](#) [[INSPIRE](#)].
- [8] L.M. Krauss, S. Nasri and M. Trodden, *A model for neutrino masses and dark matter*, *Phys. Rev. D* **67** (2003) 085002 [[hep-ph/0210389](#)] [[INSPIRE](#)].
- [9] A. Ahriche, S. Nasri and R. Soualah, *Radiative neutrino mass model at the e^-e^+ linear collider*, *Phys. Rev. D* **89** (2014) 095010 [[arXiv:1403.5694](#)] [[INSPIRE](#)].
- [10] E. Ma, *Verifiable radiative seesaw mechanism of neutrino mass and dark matter*, *Phys. Rev. D* **73** (2006) 077301 [[hep-ph/0601225](#)] [[INSPIRE](#)].
- [11] T. Hambye, K. Kannike, E. Ma and M. Raidal, *Emanations of dark matter: muon anomalous magnetic moment, radiative neutrino mass and novel leptogenesis at the TeV scale*, *Phys. Rev. D* **75** (2007) 095003 [[hep-ph/0609228](#)] [[INSPIRE](#)].
- [12] R. Bouchand and A. Merle, *Running of radiative neutrino masses: the scotogenic model*, *JHEP* **07** (2012) 084 [[arXiv:1205.0008](#)] [[INSPIRE](#)].
- [13] E. Ma, *Radiative scaling neutrino mass and warm dark matter*, *Phys. Lett. B* **717** (2012) 235 [[arXiv:1206.1812](#)] [[INSPIRE](#)].
- [14] D. Hehn and A. Ibarra, *A radiative model with a naturally mild neutrino mass hierarchy*, *Phys. Lett. B* **718** (2013) 988 [[arXiv:1208.3162](#)] [[INSPIRE](#)].
- [15] E. Ma, *Vanishing Higgs one-loop quadratic divergence in the scotogenic model and beyond*, *Phys. Lett. B* **732** (2014) 167 [[arXiv:1401.3284](#)] [[INSPIRE](#)].
- [16] S. Fraser, E. Ma and O. Popov, *Scotogenic inverse seesaw model of neutrino mass*, *Phys. Lett. B* **737** (2014) 280 [[arXiv:1408.4785](#)] [[INSPIRE](#)].
- [17] D. Schmidt, T. Schwetz and T. Toma, *Direct detection of leptophilic dark matter in a model with radiative neutrino masses*, *Phys. Rev. D* **85** (2012) 073009 [[arXiv:1201.0906](#)] [[INSPIRE](#)].
- [18] J. March-Russell, C. McCabe and M. McCullough, *Neutrino-flavoured sneutrino dark matter*, *JHEP* **03** (2010) 108 [[arXiv:0911.4489](#)] [[INSPIRE](#)].
- [19] M. Aoki, S. Kanemura, T. Shindou and K. Yagyu, *An R -parity conserving radiative neutrino mass model without right-handed neutrinos*, *JHEP* **07** (2010) 084 [Erratum *ibid.* **11** (2010) 049] [[arXiv:1005.5159](#)] [[INSPIRE](#)].
- [20] M. Aoki, J. Kubo, T. Okawa and H. Takano, *Impact of inert Higgsino dark matter*, *Phys. Lett. B* **707** (2012) 107 [[arXiv:1110.5403](#)] [[INSPIRE](#)].
- [21] S. Kanemura, N. Machida and T. Shindou, *Radiative neutrino mass, dark matter and electroweak baryogenesis from the supersymmetric gauge theory with confinement*, *Phys. Lett. B* **738** (2014) 178 [[arXiv:1405.5834](#)] [[INSPIRE](#)].
- [22] S. Kanemura, O. Seto and T. Shimomura, *Masses of dark matter and neutrino from TeV scale spontaneous $U(1)_{B-L}$ breaking*, *Phys. Rev. D* **84** (2011) 016004 [[arXiv:1101.5713](#)] [[INSPIRE](#)].
- [23] M. Lindner, D. Schmidt and T. Schwetz, *Dark matter and neutrino masses from global $U(1)_{B-L}$ symmetry breaking*, *Phys. Lett. B* **705** (2011) 324 [[arXiv:1105.4626](#)] [[INSPIRE](#)].
- [24] S. Kanemura, T. Nabeshima and H. Sugiyama, *TeV-scale seesaw with loop-induced Dirac mass term and dark matter from $U(1)_{B-L}$ gauge symmetry breaking*, *Phys. Rev. D* **85** (2012) 033004 [[arXiv:1111.0599](#)] [[INSPIRE](#)].

- [25] H. Okada and T. Toma, *Fermionic dark matter in radiative inverse seesaw model with $U(1)_{B-L}$* , *Phys. Rev. D* **86** (2012) 033011 [[arXiv:1207.0864](#)] [[INSPIRE](#)].
- [26] Y. Kajiyama, H. Okada and T. Toma, *Light dark matter candidate in $B-L$ gauged radiative inverse seesaw*, *Eur. Phys. J. C* **73** (2013) 2381 [[arXiv:1210.2305](#)] [[INSPIRE](#)].
- [27] H. Okada, *Dark matters in gauged $B-3L_i$ model*, [arXiv:1212.0492](#) [[INSPIRE](#)].
- [28] S. Baek, H. Okada and T. Toma, *Two loop neutrino model and dark matter particles with global $B-L$ symmetry*, *JCAP* **06** (2014) 027 [[arXiv:1312.3761](#)] [[INSPIRE](#)].
- [29] S. Kanemura, T. Matsui and H. Sugiyama, *Neutrino mass and dark matter from gauged $U(1)_{B-L}$ breaking*, *Phys. Rev. D* **90** (2014) 013001 [[arXiv:1405.1935](#)] [[INSPIRE](#)].
- [30] Y.H. Ahn and H. Okada, *Non-zero θ_{13} linking to dark matter from non-Abelian discrete flavor model in radiative seesaw*, *Phys. Rev. D* **85** (2012) 073010 [[arXiv:1201.4436](#)] [[INSPIRE](#)].
- [31] E. Ma, A. Natale and A. Rashed, *Scotogenic A_4 neutrino model for nonzero θ_{13} and large δ_{CP}* , *Int. J. Mod. Phys. A* **27** (2012) 1250134 [[arXiv:1206.1570](#)] [[INSPIRE](#)].
- [32] Y. Kajiyama, H. Okada and K. Yagyu, *T_7 flavor model in three loop seesaw and Higgs phenomenology*, *JHEP* **10** (2013) 196 [[arXiv:1307.0480](#)] [[INSPIRE](#)].
- [33] A.E. Carcamo Hernandez, I. de Medeiros Varzielas, S.G. Kovalenko, H. Päs and I. Schmidt, *Lepton masses and mixings in an A_4 multi-Higgs model with a radiative seesaw mechanism*, *Phys. Rev. D* **88** (2013) 076014 [[arXiv:1307.6499](#)] [[INSPIRE](#)].
- [34] E. Ma and A. Natale, *Scotogenic Z_2 or $U(1)_D$ model of neutrino mass with $\Delta(27)$ symmetry*, *Phys. Lett. B* **734** (2014) 403 [[arXiv:1403.6772](#)] [[INSPIRE](#)].
- [35] M. Lindner, S. Schmidt and J. Smirnov, *Neutrino masses and conformal electro-weak symmetry breaking*, *JHEP* **10** (2014) 177 [[arXiv:1405.6204](#)] [[INSPIRE](#)].
- [36] H. Okada and Y. Orikasa, *Classically conformal radiative neutrino model with gauged $B-L$ symmetry*, [arXiv:1412.3616](#) [[INSPIRE](#)].
- [37] S. Kanemura and H. Sugiyama, *Dark matter and a suppression mechanism for neutrino masses in the Higgs triplet model*, *Phys. Rev. D* **86** (2012) 073006 [[arXiv:1202.5231](#)] [[INSPIRE](#)].
- [38] Y. Kajiyama, H. Okada and K. Yagyu, *Two loop radiative seesaw model with inert triplet scalar field*, *Nucl. Phys. B* **874** (2013) 198 [[arXiv:1303.3463](#)] [[INSPIRE](#)].
- [39] A. Ahriche, C.-S. Chen, K.L. McDonald and S. Nasri, *Three-loop model of neutrino mass with dark matter*, *Phys. Rev. D* **90** (2014) 015024 [[arXiv:1404.2696](#)] [[INSPIRE](#)].
- [40] P.-H. Gu and U. Sarkar, *Radiative neutrino mass, dark matter and leptogenesis*, *Phys. Rev. D* **77** (2008) 105031 [[arXiv:0712.2933](#)] [[INSPIRE](#)].
- [41] S. Kanemura, T. Nabeshima and H. Sugiyama, *Neutrino masses from loop-induced Dirac Yukawa couplings*, *Phys. Lett. B* **703** (2011) 66 [[arXiv:1106.2480](#)] [[INSPIRE](#)].
- [42] Y. Farzan and E. Ma, *Dirac neutrino mass generation from dark matter*, *Phys. Rev. D* **86** (2012) 033007 [[arXiv:1204.4890](#)] [[INSPIRE](#)].
- [43] S. Kanemura, T. Matsui and H. Sugiyama, *Loop suppression of Dirac neutrino mass in the neutrinophilic two Higgs doublet model*, *Phys. Lett. B* **727** (2013) 151 [[arXiv:1305.4521](#)] [[INSPIRE](#)].

- [44] H. Okada, *Two loop induced Dirac neutrino model and dark matters with global $U(1)'$ symmetry*, [arXiv:1404.0280](#) [[INSPIRE](#)].
- [45] E. Ma, *Radiative origin of all quark and lepton masses through dark matter with flavor symmetry*, *Phys. Rev. Lett.* **112** (2014) 091801 [[arXiv:1311.3213](#)] [[INSPIRE](#)].
- [46] H. Okada and K. Yagyu, *Radiative generation of lepton masses*, *Phys. Rev. D* **89** (2014) 053008 [[arXiv:1311.4360](#)] [[INSPIRE](#)].
- [47] S. Baek, H. Okada and T. Toma, *Radiative lepton model and dark matter with global $U(1)'$ symmetry*, *Phys. Lett. B* **732** (2014) 85 [[arXiv:1401.6921](#)] [[INSPIRE](#)].
- [48] H. Okada and K. Yagyu, *Radiative generation of lepton masses with the $U(1)'$ gauge symmetry*, *Phys. Rev. D* **90** (2014) 035019 [[arXiv:1405.2368](#)] [[INSPIRE](#)].
- [49] E. Ma, *Syndetic model of fundamental interactions*, *Phys. Lett. B* **741** (2015) 202 [[arXiv:1411.6679](#)] [[INSPIRE](#)].
- [50] M. Aoki, S. Kanemura and O. Seto, *Neutrino mass, dark matter and baryon asymmetry via TeV-scale physics without fine-tuning*, *Phys. Rev. Lett.* **102** (2009) 051805 [[arXiv:0807.0361](#)] [[INSPIRE](#)].
- [51] M. Aoki, S. Kanemura and O. Seto, *A model of TeV scale physics for neutrino mass, dark matter and baryon asymmetry and its phenomenology*, *Phys. Rev. D* **80** (2009) 033007 [[arXiv:0904.3829](#)] [[INSPIRE](#)].
- [52] M. Aoki, S. Kanemura and K. Yagyu, *Triviality and vacuum stability bounds in the three-loop neutrino mass model*, *Phys. Rev. D* **83** (2011) 075016 [[arXiv:1102.3412](#)] [[INSPIRE](#)].
- [53] E. Ma, *Pathways to naturally small neutrino masses*, *Phys. Rev. Lett.* **81** (1998) 1171 [[hep-ph/9805219](#)] [[INSPIRE](#)].
- [54] A. Pilaftsis, *Radiatively induced neutrino masses and large Higgs neutrino couplings in the standard model with Majorana fields*, *Z. Phys. C* **55** (1992) 275 [[hep-ph/9901206](#)] [[INSPIRE](#)].
- [55] P.-H. Gu and U. Sarkar, *Radiative seesaw in left-right symmetric model*, *Phys. Rev. D* **78** (2008) 073012 [[arXiv:0807.0270](#)] [[INSPIRE](#)].
- [56] P. Fileviez Perez and M.B. Wise, *On the origin of neutrino masses*, *Phys. Rev. D* **80** (2009) 053006 [[arXiv:0906.2950](#)] [[INSPIRE](#)].
- [57] M. Aoki, S. Kanemura and K. Yagyu, *Doubly-charged scalar bosons from the doublet*, *Phys. Lett. B* **702** (2011) 355 [*Erratum ibid.* **706** (2012) 495] [[arXiv:1105.2075](#)] [[INSPIRE](#)].
- [58] K. Kumericki, I. Picek and B. Radovic, *Critique of fermionic $R\nu$ MDM and its scalar variants*, *JHEP* **07** (2012) 039 [[arXiv:1204.6597](#)] [[INSPIRE](#)].
- [59] K. Kumericki, I. Picek and B. Radovic, *TeV-scale seesaw with quintuplet fermions*, *Phys. Rev. D* **86** (2012) 013006 [[arXiv:1204.6599](#)] [[INSPIRE](#)].
- [60] P.S.B. Dev and A. Pilaftsis, *Minimal radiative neutrino mass mechanism for inverse seesaw models*, *Phys. Rev. D* **86** (2012) 113001 [[arXiv:1209.4051](#)] [[INSPIRE](#)].
- [61] M. Gustafsson, J.M. No and M.A. Rivera, *Predictive model for radiatively induced neutrino masses and mixings with dark matter*, *Phys. Rev. Lett.* **110** (2013) 211802 [[arXiv:1212.4806](#)] [[INSPIRE](#)].

- [62] M. Aoki, J. Kubo and H. Takano, *Two-loop radiative seesaw mechanism with multicomponent dark matter explaining the possible γ excess in the Higgs boson decay and at the Fermi LAT*, *Phys. Rev. D* **87** (2013) 116001 [[arXiv:1302.3936](#)] [[INSPIRE](#)].
- [63] Y. Kajiyama, H. Okada and T. Toma, *Multicomponent dark matter particles in a two-loop neutrino model*, *Phys. Rev. D* **88** (2013) 015029 [[arXiv:1303.7356](#)] [[INSPIRE](#)].
- [64] B. Dasgupta, E. Ma and K. Tsumura, *Weakly interacting massive particle dark matter and radiative neutrino mass from Peccei-Quinn symmetry*, *Phys. Rev. D* **89** (2014) 041702 [[arXiv:1308.4138](#)] [[INSPIRE](#)].
- [65] K.L. McDonald, *Probing exotic fermions from a seesaw/radiative model at the LHC*, *JHEP* **11** (2013) 131 [[arXiv:1310.0609](#)] [[INSPIRE](#)].
- [66] M. Aoki and T. Toma, *Impact of semi-annihilation of Z_3 symmetric dark matter with radiative neutrino masses*, *JCAP* **09** (2014) 016 [[arXiv:1405.5870](#)] [[INSPIRE](#)].
- [67] H. Okada and Y. Orikasa, *X-ray line in radiative neutrino model with global U(1) symmetry*, *Phys. Rev. D* **90** (2014) 075023 [[arXiv:1407.2543](#)] [[INSPIRE](#)].
- [68] H. Hatanaka, K. Nishiwaki, H. Okada and Y. Orikasa, *A three-loop neutrino model with global U(1) symmetry*, *Nucl. Phys. B* **894** (2015) 268 [[arXiv:1412.8664](#)] [[INSPIRE](#)].
- [69] F. Bonnet, M. Hirsch, T. Ota and W. Winter, *Systematic study of the $D = 5$ Weinberg operator at one-loop order*, *JHEP* **07** (2012) 153 [[arXiv:1204.5862](#)] [[INSPIRE](#)].
- [70] A. Ahriche, K.L. McDonald and S. Nasri, *A model of radiative neutrino mass: with or without dark matter*, *JHEP* **10** (2014) 167 [[arXiv:1404.5917](#)] [[INSPIRE](#)].
- [71] C.-S. Chen, K.L. McDonald and S. Nasri, *A class of three-loop models with neutrino mass and dark matter*, *Phys. Lett. B* **734** (2014) 388 [[arXiv:1404.6033](#)] [[INSPIRE](#)].
- [72] D. Aristizabal Sierra, A. Degee, L. Dorame and M. Hirsch, *Systematic classification of two-loop realizations of the Weinberg operator*, *JHEP* **03** (2015) 040 [[arXiv:1411.7038](#)] [[INSPIRE](#)].
- [73] S.S.C. Law and K.L. McDonald, *The simplest models of radiative neutrino mass*, *Int. J. Mod. Phys. A* **29** (2014) 1450064 [[arXiv:1303.6384](#)] [[INSPIRE](#)].
- [74] E. Ma, *Derivation of dark matter parity from lepton parity*, [arXiv:1502.02200](#) [[INSPIRE](#)].
- [75] S. Baek, N.G. Deshpande, X.G. He and P. Ko, *Muon anomalous $g - 2$ and gauged $L_\mu - L_\tau$ models*, *Phys. Rev. D* **64** (2001) 055006 [[hep-ph/0104141](#)] [[INSPIRE](#)].
- [76] E.J. Chun and K. Turzyski, *Quasi-degenerate neutrinos and leptogenesis from $L_\mu - L_\tau$* , *Phys. Rev. D* **76** (2007) 053008 [[hep-ph/0703070](#)] [[INSPIRE](#)].
- [77] S. Baek and P. Ko, *Phenomenology of $U(1)_{L_\mu - L_\tau}$ charged dark matter at PAMELA and colliders*, *JCAP* **10** (2009) 011 [[arXiv:0811.1646](#)] [[INSPIRE](#)].
- [78] P.J. Fox and E. Poppitz, *Leptophilic dark matter*, *Phys. Rev. D* **79** (2009) 083528 [[arXiv:0811.0399](#)] [[INSPIRE](#)].
- [79] D. Feldman, Z. Liu, P. Nath and G. Peim, *Multicomponent dark matter in supersymmetric hidden sector extensions*, *Phys. Rev. D* **81** (2010) 095017 [[arXiv:1004.0649](#)] [[INSPIRE](#)].
- [80] J. Heeck and W. Rodejohann, *Gauged $L_\mu - L_\tau$ symmetry at the electroweak scale*, *Phys. Rev. D* **84** (2011) 075007 [[arXiv:1107.5238](#)] [[INSPIRE](#)].

- [81] X.-G. He, G.C. Joshi, H. Lew and R.R. Volkas, *Simplest Z' model*, *Phys. Rev. D* **44** (1991) 2118 [[INSPIRE](#)].
- [82] R. Foot, X.G. He, H. Lew and R.R. Volkas, *Model for a light Z' boson*, *Phys. Rev. D* **50** (1994) 4571 [[hep-ph/9401250](#)] [[INSPIRE](#)].
- [83] S. Baek and P. Ko, *Phenomenology of $U(1)_{L_\mu-L_\tau}$ charged dark matter at PAMELA and colliders*, *JCAP* **10** (2009) 011 [[arXiv:0811.1646](#)] [[INSPIRE](#)].
- [84] K. Harigaya, T. Igari, M.M. Nojiri, M. Takeuchi and K. Tobe, *Muon $g-2$ and LHC phenomenology in the $L_\mu-L_\tau$ gauge symmetric model*, *JHEP* **03** (2014) 105 [[arXiv:1311.0870](#)] [[INSPIRE](#)].
- [85] T. Araki et al., *Cosmic neutrino spectrum and the muon anomalous magnetic moment in the gauged $L_\mu-L_\tau$ model*, *Phys. Rev. D* **91** (2015) 037301 [[arXiv:1409.4180](#)] [[INSPIRE](#)].
- [86] AMS collaboration, M. Aguilar et al., *First result from the Alpha Magnetic Spectrometer on the International Space Station: precision measurement of the positron fraction in primary cosmic rays of 0.5-350 GeV*, *Phys. Rev. Lett.* **110** (2013) 141102 [[INSPIRE](#)].
- [87] Q.-H. Cao, C.-R. Chen and T. Gong, *Leptophilic dark matter and AMS-02 cosmic-ray positron flux*, [arXiv:1409.7317](#) [[INSPIRE](#)].
- [88] H. Fritzsch, Z.-Z. Xing and S. Zhou, *Two-zero textures of the Majorana neutrino mass matrix and current experimental tests*, *JHEP* **09** (2011) 083 [[arXiv:1108.4534](#)] [[INSPIRE](#)].
- [89] Z. Maki, M. Nakagawa and S. Sakata, *Remarks on the unified model of elementary particles*, *Prog. Theor. Phys.* **28** (1962) 870 [[INSPIRE](#)].
- [90] D.V. Forero, M. Tortola and J.W.F. Valle, *Neutrino oscillations refitted*, *Phys. Rev. D* **90** (2014) 093006 [[arXiv:1405.7540](#)] [[INSPIRE](#)].
- [91] MUON G-2 collaboration, G.W. Bennett et al., *Final report of the muon E821 anomalous magnetic moment measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)] [[INSPIRE](#)].
- [92] F. Jegerlehner and A. Nyffeler, *The muon $g-2$* , *Phys. Rept.* **477** (2009) 1 [[arXiv:0902.3360](#)] [[INSPIRE](#)].
- [93] M. Benayoun, P. David, L. DelBuono and F. Jegerlehner, *Upgraded breaking of the HLS model: a full solution to the $\tau^-e^+e^-$ and ϕ decay issues and its consequences on $g-2$ VMD estimates*, *Eur. Phys. J. C* **72** (2012) 1848 [[arXiv:1106.1315](#)] [[INSPIRE](#)].
- [94] MEG collaboration, J. Adam et al., *New constraint on the existence of the $\mu^+ \rightarrow e^+\gamma$ decay*, *Phys. Rev. Lett.* **110** (2013) 201801 [[arXiv:1303.0754](#)] [[INSPIRE](#)].
- [95] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Neutrino trident production: a powerful probe of new physics with neutrino beams*, *Phys. Rev. Lett.* **113** (2014) 091801 [[arXiv:1406.2332](#)] [[INSPIRE](#)].
- [96] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Quark flavor transitions in $L_\mu-L_\tau$ models*, *Phys. Rev. D* **89** (2014) 095033 [[arXiv:1403.1269](#)] [[INSPIRE](#)].
- [97] CHARM-II collaboration, D. Geiregat et al., *First observation of neutrino trident production*, *Phys. Lett. B* **245** (1990) 271 [[INSPIRE](#)].
- [98] CCFR collaboration, S.R. Mishra et al., *Neutrino tridents and WZ interference*, *Phys. Rev. Lett.* **66** (1991) 3117 [[INSPIRE](#)].

- [99] BABAR collaboration, J.P. Lees et al., *Search for a dark photon in e^+e^- collisions at BaBar*, *Phys. Rev. Lett.* **113** (2014) 201801 [[arXiv:1406.2980](#)] [[INSPIRE](#)].
- [100] S. Baek, P. Ko and W.-I. Park, *Local Z_2 scalar dark matter model confronting galactic GeV-scale γ -ray and muon ($g-2$)*, [arXiv:1407.6588](#) [[INSPIRE](#)].
- [101] ATLAS collaboration, *Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [102] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].
- [103] S. Kanemura, K. Tsumura, K. Yagyu and H. Yokoya, *Fingerprinting nonminimal Higgs sectors*, *Phys. Rev. D* **90** (2014) 075001 [[arXiv:1406.3294](#)] [[INSPIRE](#)].
- [104] D.M. Asner et al., *ILC Higgs white paper*, [arXiv:1310.0763](#) [[INSPIRE](#)].
- [105] S. Dawson et al., *Working group report: Higgs boson*, [arXiv:1310.8361](#) [[INSPIRE](#)].