$\mathcal{N} = 2$ central charge bounds from $2d$ chiral algebras

Madalena Lemos^a and Pedro Liendo^b

^a*DESY Hamburg, Theory Group,
Notkestrasse 85, D-22607 Hamburg, Germany*

^b*IMIP, Humboldt-Universität zu Berlin, IRIS Adlershof,
Zum Großen Windkanal 6, 12489 Berlin, Germany*

E-mail: madalena.lemos@desy.de, pliendo@physik.hu-berlin.de

ABSTRACT: We study protected correlation functions in $\mathcal{N} = 2$ SCFT whose description is captured by a two-dimensional chiral algebra. Our analysis implies a new analytic bound for the c -anomaly as a function of the flavor central charge k , valid for any theory with a flavor symmetry. Combining our result with older bounds in the literature puts strong constraints on the parameter space of $\mathcal{N} = 2$ theories. In particular, it singles out a special set of models whose value of c is uniquely fixed once k is given. This set includes the canonical rank one $\mathcal{N} = 2$ SCFTs given by Kodaira's classification.

KEYWORDS: Conformal and W Symmetry, Extended Supersymmetry

ARXIV EPRINT: [1511.07449](https://arxiv.org/abs/1511.07449)

Contents

1	Introduction and summary	1
2	A new analytic bound	3
2.1	Schur correlators in $4d$	4
2.2	Twisted correlators in $2d$	6
3	Consequences for the $\mathcal{N} = 2$ landscape	8

1 Introduction and summary

Every four-dimensional $\mathcal{N} = 2$ superconformal field theory (SCFT) contains a protected subsector isomorphic to a two-dimensional chiral algebra [1]. This observation has led to interesting insights into the structure of $\mathcal{N} = 2$ superconformal dynamics, and it will be our main tool in this work. The construction starts by considering a certain class of operators that belong to the cohomology of a nilpotent supercharge. These operators are restricted to lie on a plane, and their correlation functions are meromorphic functions of the coordinates on the plane. Two-dimensional chiral algebras are very rigid structures, allowing to fix an infinite amount of protected data from the parent four-dimensional theory. To name a few, it allows to fix selected correlation functions, predict Higgs branch relations, and put constraints on central charges [2–4]. There has also been a very fruitful interplay between these chiral algebras and the superconformal index [5–10].

A similar construction was considered in six dimensions, allowing the computation of certain exact three-point functions [11]; and in three dimensions [12], where the cohomological argument implies a subset of operators is described by a one-dimensional topological theory.

In this note we will be interested in central charge bounds, which can be obtained thanks to a subtle interplay between two-dimensional and four-dimensional quantities. As will be reviewed below, the existence of bounds is closely connected to the presence of null states in the chiral algebra.

Our assumptions will be minimal: an interacting theory with a unique stress tensor, and a flavor current generating a global flavor symmetry. These two conserved currents sit in shortened multiplets of the $\mathcal{N} = 2$ superconformal algebra, that contain an element captured by the chiral algebra. These multiplets have been studied individually from the chiral perspective in [1] and [4], leading to new unitarity bounds.¹ Here we consider the logical next step and study the most natural mixed system of correlators: all possible combinations between flavor and stress-tensor multiplets. This analysis will lead to a new

¹Similarly, a unitarity bound for $\mathcal{N} = 4$ SCFTs was obtained in [13].

G	H_1	H_2	D_4	E_6	E_7	E_8
c_{4d}	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{7}{6}$	$\frac{13}{6}$	$\frac{19}{6}$	$\frac{31}{6}$
k_{4d}	$\frac{8}{3}$	3	4	6	8	12

Table 1. Properties of the canonical rank one SCFTs associated to maximal mass deformations of the Kodaira singularities with a flavor symmetry [14–19]. The flavor symmetry of the theories is given by G with $H_i \rightarrow A_i$. They can also be obtained as the low energy theory living on a single $D3$ -brane probing an F -theory singularity of type G [15, 16, 20–23].

analytic bound that when complemented with older results severely reduces the parameter space of $\mathcal{N} = 2$ theories. This bound constrains, for each flavor group, the allowed values of the flavor central charge k_{4d} and the c_{4d} -anomaly coefficient:

$$k_{4d} \left(-180c_{4d}^2 + 66c_{4d} + 3\dim_G \right) + 60c_{4d}^2 h^\vee - 22c_{4d} h^\vee \leq 0, \quad (1.1)$$

where the group information enters through the dual Coxeter number $h^\vee$ and the dimension of the algebra $\dim_G$.

This bound carves out the low c_{4d} portion of the (c_{4d}, k_{4d}) plane, explaining why there are no known theories in this region of the $\mathcal{N} = 2$ landscape. This new bound does not only give a general constraint valid for any theory, but it will also allow us to learn more about a specific set of models that sit at special corners of the allowed parameter space.

We present them in table 1 along with their values of c_{4d} and k_{4d} , and a short description on ways in which they can be obtained. These models all share the property that their Higgs branch is given by the one-instanton moduli space of G , with their Higgs branch chiral rings finitely generated by the moment maps subject to a relation defining the “Joseph ideal” (see e.g., [24]). In fact, in [1], the Higgs branch relation was formulated as a saturation of a unitarity bound, and it was shown that only a restricted number of flavor symmetries can satisfy this constraint: $G = A_1, A_2, D_4, E_6, E_7, E_8, G_2, F_4$. Moreover, for this to happen the values of c_{4d} and k_{4d} are fixed to specific values (the only exception being A_1 ²). These values matched precisely the ones given in table 1, and predicted two more cases which do not correspond to known theories: G_2 and F_4 . This was an elegant way of rederiving table 1, and one can now ask the interesting question of how unique these theories are. Namely, if one considers a generic theory assuming nothing about its Higgs branch, and we fix k_{4d} to one of the values given in table 1, what can we say about the value of c_{4d} ? The results of [1] allow for a range of c_{4d} values for each of the k_{4d} in the table, and the question of uniqueness remained unanswered.

The new analytic bound obtained here (1.1), in combination with the old bounds of [1], will succeed in singling out the canonical set of table 1 with minimal physical input. We will see that these theories live at the intersection of the old and new bound: given k_{4d} from table 1, the old bound acts as an upper bound on the central charge c_{4d} , while the

²For generic G the relations obtained imply two constraints that fix both c_{4d} and k_{4d} , while for A_1 there is only one constraint that fixes c_{4d} as a function of k_{4d} .

new becomes a lower bound on c_{4d} , allowing for a unique value. The Higgs branch relation then follows as a consequence!

The new bound (1.1) also has implications for the numerical bootstrap program for $\mathcal{N} = 2$ theories initiated in [25, 26]. It indicates that the set of correlators studied in this paper is a promising way to bootstrap the canonical set of table 1, along with the mysterious G_2 and F_4 cases. While D_4 is the familiar $\mathfrak{su}(2)$ superconformal QCD, the remaining ones have no known Lagrangian description, and the numerical bootstrap seems to be the best approach in order to access their non-protected sector. The bootstrap program for $\mathcal{N} = 2$ theories can be thought of as a two-step process. The first one consists in fixing the part of the correlators captured by the chiral algebra, and obtaining from them an infinite number of OPE coefficients. The second step involves the numerical bootstrap of the complete system of correlators, constraining the CFT data not captured by the chiral algebra. For the second step it is necessary to write the superconformal block decomposition of these correlators. While the blocks for the superconformal primary of the flavor current multiplet four-point function have been obtained in the literature [27–29], the remaining ones are still unknown. Still, the results of this paper fulfil the first requirement, and provide the starting point toward the full-blown numerical bootstrap for this system of correlators.

2 A new analytic bound

Let us start by reviewing the $2d$ chiral algebra description of $4d$ $\mathcal{N} = 2$ SCFTs. We refer the reader to [1] for more details. The construction relies on the existence of an $\mathfrak{su}(1,1|2)$ subalgebra of the full superconformal algebra with an $\mathfrak{sl}(2) \times \mathfrak{su}(2)_R$ factor, and a nilpotent supercharge $\mathbb{Q}$. It can be shown that the diagonal subalgebra of $\mathfrak{sl}(2) \times \mathfrak{su}(2)_R$, denoted by $\mathfrak{sl}(2)$, is $\mathbb{Q}$ -exact.

We restrict local operators to a plane with coordinates $(z, \bar{z})$, of which $\mathfrak{sl}(2) \times \mathfrak{sl}(2)$ is the global conformal algebra. If an operator at the origin is in the cohomology of $\mathbb{Q}$, we can use the twisted $\mathfrak{sl}(2)$ to generate anti-holomorphic translations such that it remains in the cohomology. Moreover, the anti-holomorphic dependence of the twisted-translated operator will be $\mathbb{Q}$ -exact, and it is then an easy exercise to prove that it will cancel from correlation functions. What remains is just a meromorphic correlator:

$$\langle \mathcal{O}(z_1, \bar{z}_1) \mathcal{O}(z_2, \bar{z}_2) \mathcal{O}(z_3, \bar{z}_3) \mathcal{O}(z_4, \bar{z}_4) \rangle = f(z_1, z_2, z_3, z_4). \quad (2.1)$$

The cohomology of $\mathbb{Q}$ is non-empty and its cohomology classes are in one-to-one correspondence with certain short multiplets of the $\mathcal{N} = 2$ superconformal algebra. We call the operators described by the $2d$ chiral algebra “Schur operators”, because they sit in multiplets that contribute to the Schur limit of the superconformal index [30–32]. In table 2 we list all the multiplets that contain Schur operators and their $4d$ and $2d$ quantum numbers. The two multiplets of interest to us are

- $\hat{\mathcal{C}}_{0(0,0)}$, conserved semi-short multiplet that contains the stress tensor and the conserved $\mathfrak{su}(2)_R$ and $\mathfrak{u}(1)_r$ currents.

Multiplet	$(\Delta, j, \bar{j}, R, r)$	$\mathcal{O}_{\text{Schur}}$	h	r
$\hat{\mathcal{B}}_R$	$(2R, 0, 0, 0, R, 0)$	$\Psi^{11\dots 1}$	R	0
$\mathcal{D}_{R(0,\bar{j})}$	$(2R + \bar{j} + 1, 0, \bar{j}, R, \bar{j} + 1)$	$\bar{\mathcal{Q}}_+^1 \Psi_{+\dots+}^{11\dots 1}$	$R + \bar{j} + 1$	$\bar{j} + \frac{1}{2}$
$\bar{\mathcal{D}}_{R(j,0)}$	$(2R + j + 1, j, 0, R, -j - 1)$	$\mathcal{Q}_+^1 \Psi_{+\dots+}^{11\dots 1}$	$R + j + 1$	$-j - \frac{1}{2}$
$\hat{\mathcal{C}}_{R(j,\bar{j})}$	$(2R + j + \bar{j} + 2, j, \bar{j}, R, 0)$	$\mathcal{Q}_+^1 \bar{\mathcal{Q}}_+^1 \Psi_{+\dots+ \bar{+}\dots\bar{+}}^{11\dots 1}$	$R + j + \bar{j} + 2$	$\bar{j} - j$

Table 2. Short multiplets that contain Schur operators using the notation of [33]. The second column gives the associated Dynkin labels: Δ corresponds to dilatations, $(j, \bar{j})$ label Lorentz representations, and (R, r) are the $\mathfrak{su}(2)_R \times \mathfrak{u}(1)_r$ quantum numbers. The third column indicates where inside the multiplet the Schur operator sits, where $(+, \bar{+}, 1)$ are the highest-weight values for $(j, \bar{j}, R)$. The last two columns give the $2d$ holomorphic dimension and r -charge in terms of $(j, \bar{j}, R)$.

- $\hat{\mathcal{B}}_1$, conserved short multiplet that contains the flavor current, and therefore transforms in the adjoint of the flavor group. Its highest weight is the moment map operator: a scalar of dimension $\Delta = 2$ that transforms in the triplet of $\mathfrak{su}(2)_R$.

We will be interested in the four-point functions of the Schur operators of these multiplets, as they are the ones described by the chiral algebra. For the $\hat{\mathcal{C}}_{0(0,0)}$ multiplet this corresponds to the $\mathfrak{su}(2)_R$ current: J_{++}^{IJ} , and we denote the associated $2d$ operator by T . For the $\hat{\mathcal{B}}_1$ multiplet the Schur operator is the moment map and we denote the corresponding $2d$ operator by J^a , with a an adjoint index. As shown in [1], the $4d$ OPEs imply the following $2d$ OPEs:

$$\begin{aligned}
 T(z)T(0) &\sim -\frac{6c_{4d}}{z^4} + 2\frac{T(0)}{z^2} + \frac{\partial T(0)}{z} + \dots, \\
 J^a(z)J^b(0) &\sim -\frac{k_{4d}}{2}\frac{\delta^{ab}}{z^2} + if^{abc}\frac{J^c(0)}{z} + \dots.
 \end{aligned}
 \tag{2.2}$$

The global $\mathfrak{sl}(2)$ enhances to full Virasoro and $T(z)$ becomes the $2d$ stress tensor. Similarly the global flavor symmetry gives rise to the holomorphic current $J^a(z)$ of an affine Kac-Moody algebra at level k_{2d} . The $2d$ central charges c_{2d} and k_{2d} are related to their four-dimensional counterparts by

$$c_{2d} = -12c_{4d}, \quad k_{2d} = -\frac{1}{2}k_{4d}.
 \tag{2.3}$$

2.1 Schur correlators in $4d$

In this work we are only interested in correlators of Schur operators because they are captured by the $2d$ chiral algebra, and not in the full content of the $\hat{\mathcal{C}}_{0,(0,0)}$ and $\hat{\mathcal{B}}_1$ multiplets. In particular we will consider the following mixed system of correlators:

$$\langle \hat{\mathcal{B}}_1 \hat{\mathcal{B}}_1 \hat{\mathcal{B}}_1 \hat{\mathcal{B}}_1 \rangle, \quad \langle \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{0,(0,0)} \rangle, \quad \langle \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{B}}_1 \hat{\mathcal{B}}_1 \rangle.
 \tag{2.4}$$

The OPE selection rules relevant for this set of correlators are,

$$\begin{aligned}\hat{\mathcal{B}}_1 \times \hat{\mathcal{B}}_1 &\sim \mathbf{1} + \hat{\mathcal{B}}_1 + \hat{\mathcal{B}}_2 + \hat{\mathcal{C}}_{0(j,j)} + \hat{\mathcal{C}}_{1(j,j)}, \\ \hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{C}}_{0,(0,0)} &\sim \mathbf{1} + \hat{\mathcal{C}}_{0(j,j)} + \hat{\mathcal{C}}_{1(j,j)}, \\ \hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{B}}_1 &\sim \hat{\mathcal{B}}_1 + \hat{\mathcal{B}}_2 + \hat{\mathcal{C}}_{0(j,j)} + \hat{\mathcal{C}}_{1(j,j)},\end{aligned}\tag{2.5}$$

where we are only listing the exchange of Schur operators.³ By an abuse of notation we denote the Schur operator coming from a given multiplet by the name of the corresponding multiplet. For the last OPE we only used $\mathfrak{su}(2)_R$ and $\mathfrak{u}(1)_r$ selection rules, which are necessary but not sufficient conditions for multiplets to appear. The multiplets $\hat{\mathcal{C}}_{0(j,j)}$ with $j > 0$ contain conserved currents of spin larger than two, and therefore they are absent in an interacting theory [36, 37]. In the last OPE all operators must transform in the adjoint of the flavor group, and since we are interested in interacting theories the $\hat{\mathcal{C}}_{0(0,0)}$ multiplet will be absent as well, as we expect a unique stress tensor in the singlet representation.

A subtle point that plays a key role in our analysis, is that the selection rules cannot distinguish between several copies of the same multiplet appearing in (2.5). In some cases when there is extra physical information one can specify their number. For example, we know there is a single stress-tensor multiplet $\hat{\mathcal{C}}_{0,(0,0)}$, and that in the $\hat{\mathcal{B}}_1 \times \hat{\mathcal{B}}_1$ OPE only the same $\hat{\mathcal{B}}_1$ appears. Generically, however, in order to distinguish between copies one would need a careful analysis of all possible correlators of the theory. Here we will be content with the mixed system presented in (2.4). We can split the operators between two groups: the ones appearing in the $\hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{C}}_{0,(0,0)}$ OPE, and the ones appearing in the $\hat{\mathcal{B}}_1 \times \hat{\mathcal{B}}_1$ OPE. When considering the first two four-point functions, only operators in either one of the groups are exchanged, but when considering the last four-point function operators in the intersection of the two groups are exchanged. We will be particularly interested in the $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}$ multiplet, because this is the one that will give the strongest bound.

An orthonormal basis: without loss of generality, let us pick an orthonormal basis of $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}$ such that only one multiplet appears in the $\hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{C}}_{0,(0,0)}$ OPE, and denote this multiplet by $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(1)}$, then,

$$\hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{C}}_{0,(0,0)} \sim \dots + \hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(1)} + \dots\tag{2.6}$$

This is possible as long as $\langle \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{0,(0,0)} \hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})} \rangle \neq 0$, which is always the case for $c > \frac{11}{30}$ [4]. Using this basis, the second group of operators appearing in the $\hat{\mathcal{B}}_1 \hat{\mathcal{B}}_1$ OPE contain $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(1)}$, plus extra multiplets which we collectively denote by $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(2)}$:

$$\hat{\mathcal{B}}_1 \times \hat{\mathcal{B}}_1 \sim \dots + \hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(1)} + \hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(2)} + \dots\tag{2.7}$$

Note that if we consider additional four-point functions we would be able to distinguish the different multiplets contained in $\hat{\mathcal{C}}_{1(\frac{1}{2},\frac{1}{2})}^{(2)}$. From our point of view however, we can treat

³See [4, 34] for the full OPE selection rules in the $\hat{\mathcal{B}}_1 \times \hat{\mathcal{B}}_1$ and $\hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{C}}_{0,(0,0)}$ channels. The $\hat{\mathcal{C}}_{0,(0,0)} \times \hat{\mathcal{B}}_1$ channel will appear in [35].

G	$h^\vee$	$\dim_G$	G	$h^\vee$	$\dim_G$
$\mathfrak{su}(N)$	N	$N^2 - 1$	F_4	9	52
$\mathfrak{so}(N)$	$N - 2$	$\frac{N(N-1)}{2}$	E_6	12	78
$\mathfrak{usp}(2N)$	$N + 1$	$N(2N + 1)$	E_7	18	133
G_2	4	14	E_8	30	248

Table 3. Dual Coxeter number and dimension of the adjoint representation for the simple Lie groups.

this collection of multiplets as a single entity, since orthogonality implies it contributes to the block expansion as a sum of squares of OPE coefficients, which is also a non-negative number. Each of the $\hat{\mathcal{C}}_{1(\frac{1}{2}, \frac{1}{2})}^{(2)}$ multiplets gives rise to a two-dimensional chiral operator, contributing to the twisted four-point functions as quasi-primary operators with holomorphic dimension $h = 4$ (see table 2).

2.2 Twisted correlators in 2d

Let us now consider the two-dimensional system of correlators:

$$\langle T(z_1)T(z_2)T(z_3)T(z_4) \rangle = \frac{1}{z_{12}^4 z_{34}^4} G_{TTTT}(z), \quad (2.8)$$

$$\langle T(z_1)T(z_2)J^a(z_3)J^b(z_4) \rangle = \frac{1}{z_{12}^4 z_{34}^2} G_{TTJJ}^{ab}(z), \quad (2.9)$$

$$\langle J^a(z_1)J^b(z_2)J^c(z_3)J^d(z_4) \rangle = \frac{1}{z_{12}^2 z_{34}^2} G_{JJJJ}^{abcd}(z). \quad (2.10)$$

We defined the usual cross-ratio $z = \frac{z_{12}z_{34}}{z_{13}z_{24}}$. These correlators can be fixed using crossing symmetry or 2d Ward identities (see e.g., [38]):

$$G_{TTTT}(z) = 1 + z^4 + \frac{z^4}{(1-z)^4} + \frac{8}{c_{2d}} \left(z^2 + z^3 + \frac{z^4}{(1-z)^2} + \frac{z^4}{1-z} \right), \quad (2.11)$$

$$G_{TTJJ}^{ab}(z) = \delta^{ab} \frac{c_{2d}(1-z)^2 + 2z^2(2-2z+z^2)}{c_{2d}(1-z)^2}, \quad (2.12)$$

$$G_{JJJJ}^{abcd}(z) = \delta^{ab}\delta^{cd} + z^2\delta^{ac}\delta^{bd} + \frac{z^2}{(1-z)^2}\delta^{ad}\delta^{cb} - \frac{z}{k_{2d}}f^{ace}f^{bde} - \frac{z}{k_{2d}(z-1)}f^{ade}f^{bce}. \quad (2.13)$$

For the last four-point function (2.13) we are only interested in operators being exchanged in the singlet channel, because these are the ones that contribute to the mixed correlator (2.12). As such, we project $G_{JJJJ}^{abcd}(z)$ onto the singlet and from here on consider only

$$G_{JJJJ}^{abcd}(z) \Big|_{\text{singlet}} = \delta^{ab}\delta^{cd} \left(1 + \frac{z^2}{\dim_G} \left(\frac{2h^\vee}{k_{2d}(1-z)} + \frac{1}{(z-1)^2} + 1 \right) \right), \quad (2.14)$$

where $\dim_G$ and $h^\vee$ are the dimension of the algebra and the dual Coxeter number, and are given in table 3 for convenience. Each correlator is described by the 2d chiral algebra

and we can therefore expand them using $\mathfrak{sl}(2)$ conformal blocks:

$$G(z) = 1 + \sum_{h=2}^{\infty} a^{(h)} g_h(z), \quad g_h(z) = z \left(-\frac{z}{2}\right)^{h-1} {}_2F_1(h, h; 2h; z), \quad h \text{ even}. \quad (2.15)$$

The flavor structure is now trivial so we omit it from the above expansion. The explicit expressions for the $a^{(h)}$ coefficients are

$$\begin{aligned} a_{TTTT}^{(h)} &= -\frac{2^{h-2}(h-3)(h-2)^2(h-1)\frac{h-4}{2}! \left(\frac{h}{2}+1\right)!! \frac{h}{2}!!(h+1)!!}{9(h-2)!!(2h-3)!!} \\ &\quad - \frac{2^{\frac{h}{2}+3}(h^2-h-1)(h-1)!!\Gamma\left(\frac{h}{2}\right)}{c_{2d}(2h-3)!!}, \\ a_{TTJJ}^{(h)} &= -\frac{4(h-1)h!}{c_{2d}(2h-3)!!}, \\ a_{JJJJ}^{(h)} &= -\frac{2^{h-2}(h-1)((h-2)!)^2}{\dim_G(2(h-2)+1)!} \left(2(h-1)h + \frac{4h^\vee}{k_{2d}}\right). \end{aligned}$$

Each $\mathfrak{sl}(2)$ block receives contributions from four-dimensional multiplets containing Schur operators, and the interplay between $2d$ and $4d$ perspectives will allow us to obtain bounds. Let us then interpret the coefficients of the $\mathfrak{sl}(2)$ block decomposition in terms of the $2d$ OPE coefficients. From the exchange of $h=4$ operators in the $2d$ correlators we find

$$a[T, T, \mathcal{O}_{h=4}^{(1)}] = a_{TTTT}^{(h=4)} = \frac{88}{15c_{4d}} - 16, \quad (2.16)$$

$$a[J, J, \mathcal{O}_{h=4}^{(1)}] + a[J, J, \mathcal{O}_{h=4}^{(2)}] = a_{JJJJ}^{(h=4)} = \frac{16(h^\vee - 3k_{4d})}{5\dim_G k_{4d}}, \quad (2.17)$$

$$a[T, T, \mathcal{O}_{h=4}^{(1)}]a[J, J, \mathcal{O}_{h=4}^{(1)}] = \left(a_{TTJJ}^{(h=4)}\right)^2 = \frac{64}{25c_{4d}^2}, \quad (2.18)$$

which is a system of equations we can solve for $a[J, J, \mathcal{O}_{h=4}^{(1)}]$ and $a[J, J, \mathcal{O}_{h=4}^{(2)}]$. These two objects determine the OPE coefficients $\lambda^2[\hat{\mathcal{B}}_1, \hat{\mathcal{B}}_1, \hat{\mathcal{C}}_{1(\frac{1}{2}, \frac{1}{2})}^{(1)}]$ and $\lambda^2[\hat{\mathcal{B}}_1, \hat{\mathcal{B}}_1, \hat{\mathcal{C}}_{1(\frac{1}{2}, \frac{1}{2})}^{(2)}]$ respectively. For that it is necessary to work out the exact relation between the $4d$ and the $2d$ OPE coefficients, which corresponds to a theory-independent numerical factor. This was done in [1, 25] for the $\hat{\mathcal{B}}_1$ correlator (using the superconformal blocks of [27–29]). Defining λ^2 as in (3.22) of [25], we can now translate the unitarity constraints in four dimensions to constraints on $a[J, J, \mathcal{O}_{h=4}^{(1)}]$ and $a[J, J, \mathcal{O}_{h=4}^{(2)}]$. Four-dimensional unitarity requires the λ^2 to be positive, and from

$$-\frac{1}{12}a[J, J, \mathcal{O}_{h=4}^{(2)}] = \lambda^2[\hat{\mathcal{B}}_1, \hat{\mathcal{B}}_1, \hat{\mathcal{C}}_{1(\frac{1}{2}, \frac{1}{2})}^{(2)}] \geq 0, \quad (2.19)$$

we get a nontrivial constraint on the allowed values of c_{4d} and k_{4d} :

$$k_{4d}(-180c_{4d}^2 + 66c_{4d} + 3\dim_G) + 60c_{4d}^2 h^\vee - 22c_{4d} h^\vee \leq 0. \quad (2.20)$$

Note that the only assumptions required to obtain this bound were the existence of a stress tensor, a flavor symmetry G , and that the theory is interacting. It provides a universal

lower bound for the central charge c_{4d} as a function of k_{4d} valid for any theory with a flavor symmetry.

Before we proceed to a more careful analysis of (2.20), let us comment that the same logic can be applied to higher-spin $\hat{\mathcal{C}}_{1(j,j)}$ operators, but the associated bounds are weaker. One could also ask whether different bounds could arise from analyzing different OPE channels or more general correlators. Decomposing (2.9) in $\mathfrak{sl}(2)$ blocks by taking the double OPE $(T(z_1)J^a(z_2))$ and $(J^b(z_3)T(z_4))$ does lead to a k_{4d} -independent bound on c_{4d} , but it turns out to be weaker than the $c_{4d} \geq \frac{11}{30}$ bound of [4]. Similarly, from an analysis of the $\langle T(z_1)J^a(z_2)J^b(z_3)J^c(z_4) \rangle$ correlator only weaker bounds on c_{4d} and k_{4d} are obtained. This exhausts all four-point functions of the minimal set given by T and J .

One can also enlarge the minimal set by adding extra multiplets. For example, any theory in which the two-dimensional stress tensor is not given by the Sugawara construction will have a $\hat{\mathcal{B}}_2$ multiplet associated to the difference $c_{\text{Sug}}T - cT_{\text{Sug}}$. Unfortunately, a preliminary analysis seems to suggest more physical input is needed in order to fix OPE coefficients from correlators involving $\hat{\mathcal{B}}_2$. More precisely, there are two different types of $4d$ multiplets that give rise to $2d$ operators of the same holomorphic dimension, and which we cannot disentangle without extra input.

3 Consequences for the $\mathcal{N} = 2$ landscape

In order to have a more comprehensive view of the landscape of $\mathcal{N} = 2$ theories, we will supplement our new bound (2.20) with the analytic bounds obtained in [1]. These were also obtained using the chiral algebra construction, but by looking only at the four-point function of affine Kac-Moody currents, instead of the full mixed system considered here. Among the most interesting bounds found in [1], is the one associated to the vanishing of a $\hat{\mathcal{B}}_2$ multiplet in the singlet channel:

$$\frac{1}{k_{4d}} \leq \frac{12c_{4d} + \dim_G}{24c_{4d}h^\vee}. \quad (3.1)$$

Bounds of the form $k_{4d} \geq n(G)$ where $n(G)$ is a real number that only depends on the flavor symmetry were also obtained. With the exception of $G = \mathfrak{su}(2)$, all of these bounds arise from the vanishing of a $\hat{\mathcal{B}}_2$ multiplet in a particular representation appearing in $\text{Sym}^2(\mathbf{adj})$. We refer the reader to the original paper for the complete list. Here we will be mostly concerned with (3.1) and (2.20).

Saturation of the analytic bounds implies relations among the different parameters appearing in the inequality. An interesting question is whether it is possible to saturate (3.1) and (2.20) simultaneously, in the range of k_{4d} allowed by the bounds of the type $k_{4d} \geq n(G)$. It turns out this can happen for only a handful of flavor symmetries with specific values for the central charges c_{4d} and k_{4d} . We list all the solutions in table 4. It is worth pointing out that in this list the value of k_{4d} saturates the bound of [1] for these groups (with the exception of the $\mathfrak{su}(2)$ case), namely $k_{4d} \geq \frac{h^\vee}{3} + 2$. Remarkably, all the rank one theories of table 1 are contained in this list. In addition, there are two solutions corresponding to flavor symmetries $\mathfrak{g}_2$ and $\mathfrak{f}_4$, which had already made an appearance in [1], but whose associated four-dimensional SCFTs are not known to exist. In fact, this collection of algebras

G	$\mathfrak{su}(2)$	$\mathfrak{su}(3)$	$\mathfrak{g}_2$	$\mathfrak{so}(8)$	$\mathfrak{f}_4$	$\mathfrak{e}_6$	$\mathfrak{e}_7$	$\mathfrak{e}_8$
c_{4d}	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{5}{6}$	$\frac{7}{6}$	$\frac{5}{3}$	$\frac{13}{6}$	$\frac{19}{6}$	$\frac{31}{6}$
k_{4d}	$\frac{8}{3}$	3	$\frac{10}{3}$	4	5	6	8	12

Table 4. Simultaneous saturation of the new bound (2.20) and the bound (3.1) for the allowed range of k_{4d} .

and central charges was also obtained in [1], although for a given k_{4d} , the bounds there still allowed for smaller c_{4d} . The pairs (c_{4d}, k_{4d}) were only fixed after imposing that the Higgs branch of the theory corresponded to the one-instanton moduli space for G instantons, i.e., by imposing Higgs branch relations that define the Joseph ideal (see e.g., [24]). This is known to hold for the aforementioned rank one theories. These relations imply the saturation of the bounds in all channels appearing in $\text{Sym}^2(\mathbf{adj}) - (2\mathbf{adj})$, where $(2\mathbf{adj})$ denotes the representation whose Dynkin labels are twice that of the adjoint. This requires G to be given by the list in table 4 and, with the exception of $G = \mathfrak{su}(2)$, the values of c_{4d} and k_{4d} are fixed by the requirement that both (3.1) and the $k_{4d} \geq n(G)$ bounds be saturated.

From the two-dimensional perspective saturation of these bounds implies the presence of a null state in the chiral algebra. For (3.1) this has as a consequence that the stress tensor of the $2d$ theory is given by the Sugawara construction. Indeed, it is easy to check that solving for c_{2d} gives the standard formula for the Sugawara central charge. This led [1] to conjecture that the chiral algebras associated to these theories are affine Kac-Moody current algebras at level $k_{2d} = -\frac{k_{4d}}{2}$, where k_{4d} is given in table 4.⁴ On the other hand, saturation of (2.20) signals the presence of a null state at level $h = 4$. This state can be interpreted as a difference between two holomorphic operators, and its absence implies the $\mathfrak{sl}(2)$ primary containing $:TT:$ can be written in terms of the J current.

Let us now analyze our new bound in the (c_{4d}, k_{4d}) plane to have a clearer picture of how the $\mathcal{N} = 2$ theory space looks. In figure 1 we plot in blue the analytic bound (2.20), and in red the analytic bounds of [1] ((3.1) and $k \geq n(G)$) for the simplest case of $G = \mathfrak{su}(2)$. We have added a sample of known theories with flavor group either equal to $\mathfrak{su}(2)$, or containing an $\mathfrak{su}(2)$ factor. The value of $k_{4d} = n(G = \mathfrak{su}(2)) = \frac{2}{3}$ is ruled out by (3.1), and now the smallest allowed k_{4d} is $\frac{8}{3}$, for which there is a unique allowed value of c_{4d} — that of the H_1 theory. Note that the free hypermultiplet, which enjoys an $\mathfrak{su}(2)$ flavor symmetry, is ruled out by the analytic bounds. This is consistent since both bounds were obtained imposing the absence of higher-spin currents. We have also superimposed the numerical bound obtained in [25]. It is clear that our analytic bound is significantly stronger.⁵ The numerical bound exhibited a puzzling minimum that would have corresponded to an $\mathcal{N} = 2$ theory with no conventional Coulomb branch, this now cannot be the case since that region is disallowed. Finally, we note that the central charges of $\mathcal{N} = 4$ SYM, which in $\mathcal{N} = 2$ language have an $\mathfrak{su}(2)$ flavor symmetry, have low values when compared to the $\mathcal{N} = 2$

⁴This was also studied in connection with the superconformal index in [6–8].

⁵This numerical $\mathfrak{su}(2)$ bound had already been improved by the $c \geq \frac{11}{30}$ result of [4], which applied to any local unitary SCFT, regardless of flavor symmetry.

Figure 1. Constraints on the (c_{4d}, k_{4d}) plane for $\mathfrak{su}(2)$ flavor symmetries arising as a combination of the analytic bound (2.20) (blue), the analytic bounds of [1] (red) and the numerical bound of [25] (gray). We have marked the position of some known $\mathcal{N} = 2$ theories which contain an $\mathfrak{su}(2)$ flavor symmetry, namely the free hyper multiplet, $\mathcal{N} = 4$ $\mathfrak{su}(N)$ SYM, and the theories obtained by N $D3$ -branes probing an F-theory singularity of type H_1 (starting with the rank $N = 1$ of table 1), and H_0 (for $N \geq 2$ when it has an $\mathfrak{su}(2)$ flavor symmetry) [19].

families. Any improvement of our central charge bound will at best be saturated by $\mathcal{N} = 4$, unless we restrict ourselves to pure $\mathcal{N} = 2$ theories. This can be accomplished by imposing the absence of certain $\mathcal{D}$ -type multiplets that contain the extra supercurrents of $\mathcal{N} = 4$.

As a second example we turn to $G = \mathfrak{e}_6$ in figure 2a. As in the $\mathfrak{su}(2)$ case there is an intersection between the analytic bounds (2.20) (shown in blue) and (3.1) (shown in red), but now the intersection point also coincides with the minimal allowed value of k_{4d} (also in red). This plot is a representative of what happens for all the remaining groups in table 4, in which the values of (c_{4d}, k_{4d}) are selected by a triple intersection of bounds, $\mathfrak{su}(2)$ being the only exception in which the k_{4d} bound is a bit weaker. The $\mathfrak{e}_6$ plot can also be complemented with the numerical bounds of [25]. Contrary to the $\mathfrak{su}(2)$ case, the numerical bounds allow to carve out a larger portion of the (c_{4d}, k_{4d}) plane. This is good news, the analytic bound is strong for low values of k_{4d} , but for larger values the numerical bound does better. The strongest results are achieved by complementing analytic and numerical techniques. In this plot we also marked the position of a few theories with flavor group equal to, or containing $\mathfrak{e}_6$ to give an idea of how the $\mathfrak{e}_6$ landscape looks.

Finally, as a representative of all the other flavor groups for which there is no intersection of (2.20) and (3.1), we show the case of $\mathfrak{su}(4)$ in figure 2b, where the first bound is shown in blue and the second in red. The gap on the right side means that for the smallest possible k_{4d} there is a range of allowed values for the central charge. Indeed, it is not hard to find a theory which lies inside this range. The rightmost dot in the figure represents $\mathfrak{su}(2)$ superconformal QCD, which has an enhanced $\mathfrak{so}(8)$ symmetry and corresponds to the $\mathfrak{so}(8)$ entry of table 4. Here we are considering it by embedding an $\mathfrak{su}(4)$ factor inside the $\mathfrak{so}(8)$. For larger rank theories, either $\mathfrak{su}(N)$, $\mathfrak{so}(N)$, $\mathfrak{usp}(N)$ the gap becomes larger

Figure 2. Constraints on the (c_{4d}, k_{4d}) plane for two different flavor symmetries arising as a combination of the analytic bound (2.20) (blue) and the analytic bounds of [1] (red). For the case of $\mathfrak{e}_6$ the numerical bound of [25] is also shown in gray. We also marked the position of several theories with at least $\mathfrak{e}_6$ and $\mathfrak{su}(4)$ flavor symmetry, namely the rank N version of some of the theories given in table 1 [19], the $T_{N \geq 4}$ family of theories of [39] which have an $\mathfrak{su}(N)^3$ flavor symmetry, and the $R_{0, N \geq 3}$ of [40] which have an $\mathfrak{su}(2) \times \mathfrak{su}(2N)$ flavor symmetry (for $N = 3$ the flavor symmetry enhances and it corresponds to the rank one $\mathfrak{e}_6$).

and the bounds are less constraining. As N increases, the upper and lower bounds get more separated, with the difference between them growing like N^2 .

To conclude, let us discuss how one could improve on the results presented here. Looking at the landscape plots, it is natural to ask whether a bound of the form $\frac{c_{4d}}{k_{4d}} \geq \dots$ could ever be found from the bootstrap. Holographic arguments suggest that such a bound should exist, at least for theories with a weakly-coupled gravitational description. On the dual AdS, the central charges c_{4d} and k_{4d} are related to the strength of the gravitational and gauge couplings, and the statement that $\frac{c_{4d}}{k_{4d}}$ is bounded from below would follow from the statement that *gravity is the weakest force* [41].⁶ The fact that the numerical $\mathfrak{e}_6$ bound improves on the analytic bound for large k , seems to hint that this information is indeed coded in the $\mathcal{N} = 2$ crossing symmetry equations. We hope to come back to this problem in the future.

Acknowledgments

We have greatly benefited from discussions with L. Rastelli and V. Schomerus. The research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union’s Seventh Framework Programme FP7/2007-2013/ under REA Grant Agreement No 317089 (GATIS). P. L. is supported by SFB 647 “Raum-Zeit-Materie. Analytische und Geometrische Strukturen”.

⁶See for example [42] for an AdS discussion of this statement and its implications for CFTs.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli and B.C. van Rees, *Infinite chiral symmetry in four dimensions*, *Commun. Math. Phys.* **336** (2015) 1359 [[arXiv:1312.5344](#)] [[INSPIRE](#)].
- [2] C. Beem, W. Peelaers, L. Rastelli and B.C. van Rees, *Chiral algebras of class S*, *JHEP* **05** (2015) 020 [[arXiv:1408.6522](#)] [[INSPIRE](#)].
- [3] M. Lemos and W. Peelaers, *Chiral algebras for trinion theories*, *JHEP* **02** (2015) 113 [[arXiv:1411.3252](#)] [[INSPIRE](#)].
- [4] P. Liendo, I.A. Ramírez and J. Seo, *Stress-tensor OPE in $N = 2$ superconformal theories*, *JHEP* **02** (2016) 019 [[arXiv:1509.00033](#)] [[INSPIRE](#)].
- [5] O. Chacaltana, J. Distler and A. Trimm, *A family of 4D $N = 2$ interacting SCFTs from the twisted A_{2N} series*, [arXiv:1412.8129](#) [[INSPIRE](#)].
- [6] M. Buican and T. Nishinaka, *On the superconformal index of Argyres-Douglas theories*, *J. Phys. A* **49** (2016) 015401 [[arXiv:1505.05884](#)] [[INSPIRE](#)].
- [7] C. Cordova and S.-H. Shao, *Schur indices, BPS particles and Argyres-Douglas theories*, *JHEP* **01** (2016) 040 [[arXiv:1506.00265](#)] [[INSPIRE](#)].
- [8] M. Buican and T. Nishinaka, *Argyres-Douglas theories, the Macdonald index and an RG inequality*, *JHEP* **02** (2016) 159 [[arXiv:1509.05402](#)] [[INSPIRE](#)].
- [9] J. Song, *Superconformal indices of generalized Argyres-Douglas theories from 2d TQFT*, *JHEP* **02** (2016) 045 [[arXiv:1509.06730](#)] [[INSPIRE](#)].
- [10] S. Cecotti, J. Song, C. Vafa and W. Yan, *Superconformal index, BPS monodromy and chiral algebras*, [arXiv:1511.01516](#) [[INSPIRE](#)].
- [11] C. Beem, L. Rastelli and B.C. van Rees, *W symmetry in six dimensions*, *JHEP* **05** (2015) 017 [[arXiv:1404.1079](#)] [[INSPIRE](#)].
- [12] S.M. Chester, J. Lee, S.S. Pufu and R. Yacoby, *Exact correlators of BPS operators from the 3d superconformal bootstrap*, *JHEP* **03** (2015) 130 [[arXiv:1412.0334](#)] [[INSPIRE](#)].
- [13] C. Beem, L. Rastelli and B.C. van Rees, *The $N = 4$ superconformal bootstrap*, *Phys. Rev. Lett.* **111** (2013) 071601 [[arXiv:1304.1803](#)] [[INSPIRE](#)].
- [14] P.C. Argyres, M.R. Plesser, N. Seiberg and E. Witten, *New $N = 2$ superconformal field theories in four-dimensions*, *Nucl. Phys. B* **461** (1996) 71 [[hep-th/9511154](#)] [[INSPIRE](#)].
- [15] J.A. Minahan and D. Nemeschansky, *An $N = 2$ superconformal fixed point with E_6 global symmetry*, *Nucl. Phys. B* **482** (1996) 142 [[hep-th/9608047](#)] [[INSPIRE](#)].
- [16] J.A. Minahan and D. Nemeschansky, *Superconformal fixed points with E_n global symmetry*, *Nucl. Phys. B* **489** (1997) 24 [[hep-th/9610076](#)] [[INSPIRE](#)].
- [17] Y.-K.E. Cheung, O.J. Ganor and M. Krogh, *Correlators of the global symmetry currents of 4D and 6D superconformal theories*, *Nucl. Phys. B* **523** (1998) 171 [[hep-th/9710053](#)] [[INSPIRE](#)].

- [18] P.C. Argyres and N. Seiberg, *S-duality in $N = 2$ supersymmetric gauge theories*, *JHEP* **12** (2007) 088 [[arXiv:0711.0054](#)] [[INSPIRE](#)].
- [19] O. Aharony and Y. Tachikawa, *A holographic computation of the central charges of $D = 4$, $N = 2$ SCFTs*, *JHEP* **01** (2008) 037 [[arXiv:0711.4532](#)] [[INSPIRE](#)].
- [20] A. Sen, *F-theory and orientifolds*, *Nucl. Phys. B* **475** (1996) 562 [[hep-th/9605150](#)] [[INSPIRE](#)].
- [21] T. Banks, M.R. Douglas and N. Seiberg, *Probing F-theory with branes*, *Phys. Lett. B* **387** (1996) 278 [[hep-th/9605199](#)] [[INSPIRE](#)].
- [22] K. Dasgupta and S. Mukhi, *F theory at constant coupling*, *Phys. Lett. B* **385** (1996) 125 [[hep-th/9606044](#)] [[INSPIRE](#)].
- [23] O. Aharony, A. Fayyazuddin and J.M. Maldacena, *The large- N limit of $N = 2$, $N = 1$ field theories from three-branes in F-theory*, *JHEP* **07** (1998) 013 [[hep-th/9806159](#)] [[INSPIRE](#)].
- [24] D. Gaiotto, A. Neitzke and Y. Tachikawa, *Argyres-Seiberg duality and the Higgs branch*, *Commun. Math. Phys.* **294** (2010) 389 [[arXiv:0810.4541](#)] [[INSPIRE](#)].
- [25] C. Beem, M. Lemos, P. Liendo, L. Rastelli and B.C. van Rees, *The $N = 2$ superconformal bootstrap*, [arXiv:1412.7541](#) [[INSPIRE](#)].
- [26] M. Lemos and P. Liendo, *Bootstrapping $N = 2$ chiral correlators*, *JHEP* **01** (2016) 025 [[arXiv:1510.03866](#)] [[INSPIRE](#)].
- [27] F.A. Dolan and H. Osborn, *Superconformal symmetry, correlation functions and the operator product expansion*, *Nucl. Phys. B* **629** (2002) 3 [[hep-th/0112251](#)] [[INSPIRE](#)].
- [28] M. Nirschl and H. Osborn, *Superconformal Ward identities and their solution*, *Nucl. Phys. B* **711** (2005) 409 [[hep-th/0407060](#)] [[INSPIRE](#)].
- [29] F.A. Dolan, L. Gallot and E. Sokatchev, *On four-point functions of 1/2-BPS operators in general dimensions*, *JHEP* **09** (2004) 056 [[hep-th/0405180](#)] [[INSPIRE](#)].
- [30] J. Kinney, J.M. Maldacena, S. Minwalla and S. Raju, *An index for 4 dimensional super conformal theories*, *Commun. Math. Phys.* **275** (2007) 209 [[hep-th/0510251](#)] [[INSPIRE](#)].
- [31] A. Gadde, L. Rastelli, S.S. Razamat and W. Yan, *The 4d superconformal index from q -deformed 2d Yang-Mills*, *Phys. Rev. Lett.* **106** (2011) 241602 [[arXiv:1104.3850](#)] [[INSPIRE](#)].
- [32] A. Gadde, L. Rastelli, S.S. Razamat and W. Yan, *Gauge theories and Macdonald polynomials*, *Commun. Math. Phys.* **319** (2013) 147 [[arXiv:1110.3740](#)] [[INSPIRE](#)].
- [33] F.A. Dolan and H. Osborn, *On short and semi-short representations for four-dimensional superconformal symmetry*, *Annals Phys.* **307** (2003) 41 [[hep-th/0209056](#)] [[INSPIRE](#)].
- [34] G. Arutyunov, B. Eden and E. Sokatchev, *On nonrenormalization and OPE in superconformal field theories*, *Nucl. Phys. B* **619** (2001) 359 [[hep-th/0105254](#)] [[INSPIRE](#)].
- [35] I.A. Ramírez, *Mixed OPEs in $N = 2$ superconformal theories*, [arXiv:1602.07269](#) [[INSPIRE](#)].
- [36] J. Maldacena and A. Zhiboedov, *Constraining conformal field theories with a higher spin symmetry*, *J. Phys. A* **46** (2013) 214011 [[arXiv:1112.1016](#)] [[INSPIRE](#)].
- [37] V. Alba and K. Diab, *Constraining conformal field theories with a higher spin symmetry in $D = 4$* , [arXiv:1307.8092](#) [[INSPIRE](#)].
- [38] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nucl. Phys. B* **241** (1984) 333 [[INSPIRE](#)].

- [39] D. Gaiotto, *$N = 2$ dualities*, *JHEP* **08** (2012) 034 [[arXiv:0904.2715](#)] [[INSPIRE](#)].
- [40] O. Chacaltana and J. Distler, *Tinkertoys for Gaiotto duality*, *JHEP* **11** (2010) 099 [[arXiv:1008.5203](#)] [[INSPIRE](#)].
- [41] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The string landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060 [[hep-th/0601001](#)] [[INSPIRE](#)].
- [42] Y. Nakayama and Y. Nomura, *Weak gravity conjecture in the AdS/CFT correspondence*, *Phys. Rev. D* **92** (2015) 126006 [[arXiv:1509.01647](#)] [[INSPIRE](#)].