RECEIVED: December 31, 2010

REVISED: June 1, 2011

ACCEPTED: June 2, 2011

PUBLISHED: July 21, 2011

Heavy octets and Tevatron signals with three or four b jets

Yang Bai^a and Bogdan A. Dobrescu^b

^a*SLAC National Accelerator Laboratory,
2575 Sand Hill Road, Menlo Park, CA 94025, U.S.A.*

^b*Theoretical Physics Department, Fermilab,
Batavia, IL 60510, U.S.A.*

E-mail: yangbai@slac.stanford.edu, bdob@fnal.gov

ABSTRACT: Hypothetical color-octet particles of spin 0, pair-produced at hadron colliders through their QCD coupling, may lead to final states involving three or four b jets. The scalar octets may also be resonantly produced in pairs via a spin-1 octet, as we show within a renormalizable coloron model. We analyze kinematic distributions of the $3b$ final state that differentiate between these two production mechanisms, as well as between scalar octets and supersymmetric Higgs bosons. A scalar octet of mass in the 140–150 GeV range can explain the nonstandard shape of the b -jet transverse energy distributions reported by the CDF Collaboration, especially when the coloron mass is slightly above twice the scalar mass. The dominant decay mode of the scalar octet in that case is into a pair of gluons, so that the production of a pair of dijet resonances is large in this model, of about 40 pb at the Tevatron. Even when a W boson is radiated from the initial state, the inclusive cross section for producing a dijet resonance near the scalar octet mass remains sizable, around 0.15 pb.

KEYWORDS: Beyond Standard Model, Hadronic Colliders, Technicolor and Composite Models

ARXIV EPRINT: [1012.5814](https://arxiv.org/abs/1012.5814)

Contents

1	Pairs of resonances	1
2	Spin-0, weak-singlet, color-octet particle	3
2.1	Renormalizable interactions	3
2.2	Decays of the scalar octet	3
2.3	CP violation in B_s mixing from scalar octet exchange	7
3	Tevatron phenomenology of scalar octets	8
3.1	$3b$ versus $4b$ signals	8
3.2	Properties of the $3b$ signal	10
3.3	Effects of bound states	11
3.4	Supersymmetric Higgs bosons	13
4	Renormalizable coloron model	14
4.1	Interactions and masses	15
4.2	ReCoM and the CDF excess in the $3b$ final state	17
4.3	Dijet resonances plus a W boson and other signals	20
5	Conclusions	22

1 Pairs of resonances

Hadron colliders, such as the Tevatron and the LHC, allow the production at a high rate of new particles carrying QCD color. The subsequent decays of such particles often involve only QCD jets, which are hard to separate from the background. However, when the jets originate from b -quark decays, the backgrounds are substantially reduced. Among the hypothetical particles leading to signatures involving several b jets are color-octet bosons. The simplest of those is the weak-singlet particle of spin 0, generically referred to here as the G_H scalar, which is present in various theories including Technicolor (“octet technipion” [1]), the 6D standard model (the Kaluza-Klein modes of the gluon polarized along extra dimensions [2]), vectorlike confinement [3, 4], weakly-interacting metastable pion models [5], or certain supersymmetric models (“sgluons” [6]). The G_H scalar can be produced in pairs through its QCD couplings to gluons, and may decay through higher-dimensional operators into a pair of heavy quarks. The signature is four b jets, forming two $b\bar{b}$ resonances of same mass [3, 4, 7–9].

Here we point out that an alternative way of searching for scalar octets is to require only three jets to pass the basic cuts and to be b tagged. We show that in this case the shapes of certain kinematic distributions may allow the separation of the signal from

the background, and the differentiation between various extensions of the standard model. For example, compared to the Minimal Supersymmetric Standard Model (MSSM) at large $\tan\beta$, where a Higgs boson is produced in association with a b quark and then decays to a $b\bar{b}$ pair, the octet pair production leads to a peak at a larger invariant mass of the leading and third jets.

Even more dramatic deformations of the distributions occur when the pair of G_H scalars is produced through an s -channel resonance. We study these effects within a renormalizable theory that includes an extension of the QCD gauge group, $SU(3) \times SU(3)$, as proposed in [10–13]. The scalar sector responsible for spontaneously breaking this symmetry down to $SU(3)_c$ includes a G_H scalar as well as two gauge singlet real scalar fields. We refer to this as the Renormalizable Coloron Model (ReCoM). The heavy gauge boson arising from the extended symmetry, labelled by G'_μ and referred to as the coloron (or more generally gluon-prime), is a color-octet particle of spin 1.

We show that the coloron couples to a pair of G_H scalars so that s -channel production of G'_μ leads to a $4b$ signal. This is an example of nested resonances:¹ two pairs of b jets form each a resonance and the combination of the two pairs also forms a resonance. These multiple resonant features allow an efficient rejection of the background. However, for that it is necessary to have four b -tagged jets, which requires large data sets. If only three b -tagged jets are required to pass the cuts, then the signal is larger and a detailed analysis could separate it from the background within smaller data sets.

The CDF Collaboration has searched for a resonance in the invariant mass distribution of the two leading b jets, produced in association with a third b jet. Preliminary results [14] suggest the existence of a resonance with mass of about 140 GeV, that could be attributed to a fluctuation of the standard model background at the 6% confidence level. A more puzzling feature of these CDF results is that the measured transverse energy (E_T) distributions of the b jets have shapes that differ notably from the standard model predictions [15]. We estimate that the shapes of the E_T distributions of the two leading b jets may be due to a fluctuation of the standard model only at around 1% confidence level. It should be emphasized though that the preliminary CDF results do not include information regarding the systematic uncertainties associated with these distributions. We demonstrate that these shapes are nicely explained within the ReCoM when the G_H mass is around 140 GeV and the G'_μ mass is slightly above the $G_H G_H$ threshold.

We describe the interactions and decays of the spin-0 octet in section 2. Then we analyze the Tevatron phenomenology of QCD-produced G_H pairs in section 3. There we also discuss the boundstate effects due to gluon exchange between the two G_H produced, and we compare the kinematic distributions of the multi- b -jet final state due to the scalar octet with those due to supersymmetric Higgs bosons. In section 4 we derive the ReCoM predictions, compare them with the CDF data for several $3b$ kinematic distributions, and discuss implications for other final states, such as those arising from associated production of a coloron and a W boson. We summarize strategies to distinguish different models with multi b jets in section 5.

¹This term was first used in an early, unpublished version of ref. [7].

2 Spin-0, weak-singlet, color-octet particle

The theory considered in this section is the standard model plus only one particle, G_H , which is a real field of spin 0, transforming as an octet under the QCD gauge group $SU(3)_c$ and as a singlet under the electroweak gauge group $SU(2)_W \times U(1)_Y$. These gauge charges imply that G_H is electrically neutral, and does not have any renormalizable interactions with the quarks and leptons (note that $G_H \bar{b}_L b_R$ does not arise from an $SU(2)_W$ invariant Yukawa interaction; the opposite is true in the case of weak-doublet scalar octets [16–18]). We refer to G_H as a scalar octet, independently of whether it is a composite (as in technicolor) or elementary particle.

2.1 Renormalizable interactions

The renormalizable couplings of G_H to gluons are fixed by $SU(3)_c$ gauge invariance:

$$\frac{g_s^2}{2} f^{abc} f^{ade} G_\mu^b G^{\mu d} G_H^c G_H^e + g_s f^{abc} G_\mu^a G_H^b \partial^\mu G_H^c, \quad (2.1)$$

where f^{abc} is the anti-symmetric $SU(3)_c$ tensor, g_s is the QCD coupling, and G_μ is the gluon field. The only other renormalizable couplings of G_H are to the Higgs doublet (H) and to itself:

$$\frac{\lambda_{HG}}{2} G_H^a G_H^a H^\dagger H + \frac{\lambda_G}{8} (G_H^a G_H^a)^2 + \mu_G d_{abc} G_H^a G_H^b G_H^c, \quad (2.2)$$

where λ_{HG} and $\lambda_G > 0$ are dimensionless parameters, μ_G is a parameter of mass dimension one, and d_{abc} is the totally-symmetric $SU(3)_c$ tensor.

The first term in eq. (2.2) contributes to the mass squared of G_H after electroweak symmetry breaking. We take the sum ($M_{G_H}^2$) of this contribution and the mass squared from the Lagrangian to be positive. As a result, G_H does not have a vacuum expectation value (VEV) provided there is an upper limit on the cubic coupling, $|\mu_G| \lesssim \sqrt{\lambda_G} M_{G_H}$. We are primarily interested in the case where the physical mass of G_H , M_{G_H} , is above around 100 GeV.

The production of G_H at hadron colliders occurs mainly in pairs, due to the couplings (2.1) to gluons, via the tree-level diagrams shown in figure 1. This production depends only on the G_H mass, because the couplings to gluons are fixed by QCD gauge invariance. Single G_H production is possible at one loop through a cubic interaction [the last term in eq. (2.2)], but it is suppressed enough (by the loop factor and by an accidental cancellation discussed in section 2.2) to be neglected.

2.2 Decays of the scalar octet

The only renormalizable coupling of G_H that violates the Z_2 invariance under the $G_H \rightarrow -G_H$ transformation is the cubic term in eq. (2.2). Thus, the only decays of G_H allowed by the renormalizable couplings shown above occur at one or more loops and involve at least one vertex proportional to μ_G . At one loop, divergent terms from the triangle and

Figure 1. $G_H G_H$ production in hadronic collisions (u -channel G_H exchange is not shown). Curly lines represent gluons, while dashed lines represent scalar octets.

Figure 2. Scalar octet decay to gluons, due to the cubic G_H interaction of eq. (2.2). A diagram similar with the left one but with interchanged end points for the gluon lines is not shown.

bubble diagrams shown in figure 2 cancel each other, and the finite result for the width of G_H decaying into two gluons is

$$\Gamma_0(G_H \rightarrow gg) = \frac{15 \alpha_s^2 \mu_G^2}{128 \pi^3 M_{G_H}} \left(\frac{\pi^2}{9} - 1 \right)^2, \quad (2.3)$$

where α_s is the QCD coupling evaluated at M_{G_H} . This width is accidentally suppressed by the small numerical coefficient given in the paranthesis (see [18] for a similar case).

Decays of G_H into quarks induced by nonrenormalizable couplings may have large branching fractions because the above decay width into gluons is suppressed. Dimension-5 operators of this type involve either a Higgs doublet,

$$\frac{C_{ij}^d}{m_\psi} H G_H^a \bar{Q}_L^i T^a d_R^j + \frac{C_{ij}^u}{m_\psi} \tilde{H} G_H^a \bar{Q}_L^i T^a u_R^j + \text{H.c.}, \quad (2.4)$$

or a covariant derivative,

$$\frac{1}{m_\psi} \left(K_{ij}^d \bar{d}_R^i \gamma^\mu T^a d_R^j + K_{ij}^u \bar{u}_R^i \gamma^\mu T^a u_R^j + K_{ij}^Q \bar{Q}_L^i \gamma^\mu T^a Q_L^j \right) D_\mu G_H^a + \text{H.c.}, \quad (2.5)$$

where $Q_L^i \equiv (u_L^i, d_L^i)$, the indices $i, j = 1, 2, 3$ label the generations, $C_{ij}^{d,u}$ and $K_{ij}^{d,u,Q}$ are complex coefficients, and m_ψ is the mass of a heavy particle which generates these operators (if several particles of different masses contribute to these operators, then their effects can be included in the coefficients). Note that in the semi-classical approximation the operators (2.5) may be transformed into the operators (2.4) through integration by parts and the use of quark field equations.

There are also dimension-5 operators contributing to the $G_H \rightarrow gg$ decay,

$$\frac{\alpha_s}{4\pi m_\psi} d_{abc} G_H^a G_{\mu\nu}^b \left(\xi_G G^{c\mu\nu} + \xi'_G \epsilon^{\mu\nu\alpha\beta} G_{\alpha\beta}^c \right), \quad (2.6)$$

Figure 3. Effective couplings of a scalar octet to b quarks and the Higgs VEV (left diagram), or to a pair of gauge bosons (right diagram), induced in the presence of a heavy vectorlike quark ψ and leading to the $G_H \rightarrow b\bar{b}, gg, g\gamma, gZ$ decays.

and to the $G_H \rightarrow gZ$ or $g\gamma$ decays,

$$\frac{\sqrt{\alpha_s \alpha}}{4\pi \cos \theta_W m_\psi} G_H^a G_{\mu\nu}^a \left(\xi_B B^{\mu\nu} + \xi'_B \epsilon^{\mu\nu\alpha\beta} B_{\alpha\beta} \right) . \quad (2.7)$$

Here, $B_{\mu\nu}$ is the field tensor of the $U(1)_Y$ gauge boson ($-\sin \theta_W Z^\mu + \cos \theta_W A^\mu$, where A^μ is the photon field) and θ_W is the weak mixing angle. These operators are generated by loops involving some new particle of mass m_ψ whose coupling to G_H determines the dimensionless coefficients ξ_G, ξ'_G, ξ_B and ξ'_B . The decay of a scalar octet into a gluon and a photon has been studied in ref. [19, 20].

Operators that allow G_H to decay into a Higgs boson and gluons arise at dimension-7 or higher. For example, the operator $(D^\mu G_H^a) G_{\mu\nu}^a H^\dagger D^\nu H$ induces the $G_H \rightarrow ggh$ decay (the $G_H \rightarrow gh$ process is forbidden by angular momentum conservation). The operators that couple G_H to leptons also appear only at dimension-7 or higher, for example, $(D^\mu G_H^a) \bar{L}^i \gamma^\nu L^j G_{\mu\nu}^a$.

Given that the operators (2.4) involve the Higgs doublet, their coefficients are typically proportional to the quark mass. It is natural to assume that the dominant decay mode induced by operators (2.4) is $G_H \rightarrow t\bar{t}$ for $M_{G_H} \gtrsim 350$ GeV, and $G_H \rightarrow b\bar{b}$ for $10 \text{ GeV} \lesssim M_{G_H} \lesssim 350$ GeV. However, the relative normalization of the C_{ij}^d and C_{ij}^u coefficients depends on the underlying mechanism responsible for generating these operators, so that the decay to $b\bar{b}$ could dominate even for $M_{G_H} \gtrsim 350$ GeV.

To be concrete, we consider the following renormalizable model as an origin for the dimension-5 operators. A vectorlike quark, ψ , having the same gauge charges as b_R may mix with the down-type quarks via the following terms in the Lagrangian:

$$m_\psi \bar{\psi}_L \psi_R + \mu_{i\psi} \bar{\psi}_L d_R^i + y_{ij} H \bar{Q}_L^i d_R^j + \lambda_i H \bar{Q}_L^i \psi_R + \text{H.c.} , \quad (2.8)$$

where y_{ij} and λ_i are Yukawa couplings, $\mu_{i\psi}$ is a mass mixing parameter, and m_ψ is the ψ fermion mass in the small $\mu_{i\psi}$ limit. The scalar octet also has Yukawa couplings to ψ :

$$G_H^a (\eta_0 \bar{\psi}_L T^a \psi_R + \eta_i \bar{\psi}_L T^a b_R^i) + \text{H.c.} \quad (2.9)$$

The above interactions lead to effective couplings of G_H to standard model quarks, as shown in the first diagram of figure 3. Upon integrating ψ out, the operators (2.4) are

induced with coefficients

$$C_{ij}^u = 0 \quad , \quad C_{ij}^d \approx \lambda_i \eta_j^* \quad , \quad (2.10)$$

where we assumed for simplicity that $m_\psi \gg \mu_{i\psi}$. The coefficients C_{ij}^d are defined in the gauge eigenstate basis. Assuming that the two unitary matrices that diagonalize the down-type quark mass matrix are close to the identity matrix, the G_H couplings to down-type quarks in the mass eigenstate basis are approximately equal to C_{ij}^d . For $|\lambda_3| \gg |\lambda_1|, |\lambda_2|$ and $|\eta_3| \gg |\eta_1|, |\eta_2|$ the width for $G_H \rightarrow b\bar{b}$ is much larger than for any other fermion final states, and is given by

$$\Gamma(G_H \rightarrow b\bar{b}) \simeq \frac{|\lambda_3 \eta_3|^2 v^2}{32\pi m_\psi^2} M_{G_H} \quad , \quad (2.11)$$

where $v = 246$ GeV is the electroweak scale. If $\lambda_{1,2}$ and $\eta_{1,2}$ are not much smaller than λ_3 and η_3 (which is allowed if m_ψ is large enough to avoid constraints on flavor-changing processes, as discussed in section 2.3), then decays into other pairs of down-type quarks may also be important, including flavor-changing ones such as $G_H \rightarrow b\bar{s}$ or $s\bar{b}$; we do not explore further this possibility.

The ψ fermion also induces some of the dimension-5 operators given in eqs. (2.6) and (2.7), as shown in figure 3. The ensuing decay width into gluons, which adds to Γ_0 given in eq. (2.3), is

$$\Gamma_\psi(G_H \rightarrow gg) \simeq \frac{5 \eta_0^2 \alpha_s^2 M_{G_H}^3}{6 (16\pi)^3 m_\psi^2} |A(M_{G_H}^2/(4m_\psi^2))|^2 \quad , \quad (2.12)$$

where the function A arises from a momentum integral given in [21]:

$$A(x) = \frac{2}{x^2} [x + (x-1)\text{Arcsin}^2 \sqrt{x}] \quad , \quad \text{for } x \leq 1 \quad . \quad (2.13)$$

For $x \ll 1$, i.e. $m_\psi \gg M_{G_H}/2$, the function has the value $A \approx 4/3$, and the coefficients of the operators (2.6) are

$$\xi_G = \frac{\eta_0}{6\sqrt{2}} \quad , \quad \xi'_G = 0 \quad . \quad (2.14)$$

The ratio of the decay widths into $b\bar{b}$ and gg , neglecting the interference of the two diagrams contributing to $G_H \rightarrow gg$ in figures 2 and 3, is given by

$$\frac{\Gamma(G_H \rightarrow b\bar{b})}{\Gamma_\psi(G_H \rightarrow gg) + \Gamma_0(G_H \rightarrow gg)} = \frac{0.52}{\eta_0^2 + 3.02 \mu_G^2 m_\psi^2 / M_{G_H}^4} \left(\frac{|\lambda_3 \eta_3|}{10^{-3}} \frac{100 \text{ GeV}}{M_{G_H}} \frac{0.1}{\alpha_s} \right)^2 \quad . \quad (2.15)$$

Note that m_ψ may be significantly above the TeV scale, so that the vectorlike quark might not be produced even at the LHC. Thus, the branching fraction of $G_H \rightarrow b\bar{b}$ can be very small if $\mu_G m_\psi \gg M_{G_H}$. Alternatively, if ψ is not much heavier than G_H , and $\eta_0 = O(1)$, then the branching fraction of $G_H \rightarrow b\bar{b}$ can be large only for $|\lambda_3 \eta_3| \gtrsim 10^{-3}$.

The scalar octet has a very narrow width (several orders of magnitude less than its mass), as its contributions arise from loops or higher dimensional operators. Its decays are usually prompt, unless μ_G is very small and m_ψ is very large. For example, $\mu_G = 0$, $\eta_0 = 0$ and $m_\psi/|\lambda_3 \eta_3| \gtrsim 2.5 \times 10^5$ TeV give a decay length $c\tau \gtrsim 100 \mu\text{m}$ so that G_H decays

off the beam line. We will not investigate further these signatures involving dijet and $b\bar{b}$ resonances originating from displaced vertices.

The decays $G_H \rightarrow g\gamma, gZ$ have small branching fractions due to coupling and color suppressions [the coefficients of the operators (2.7) are comparable with those in eq. (2.14): $\xi_B = \xi_G/3$ and $\xi'_B = 0$]. For example,

$$\frac{\Gamma(G_H \rightarrow g\gamma)}{\Gamma_\psi(G_H \rightarrow gg)} = \frac{\alpha}{15\alpha_s} \approx 0.44\% \quad . \quad (2.16)$$

Nevertheless, a photon-plus-jet resonance that arises together with a dijet resonance of equal mass, is an interesting signature that follows from the decays of a G_H pair.

To summarize, within a simple renormalizable origin of the dimension-5 operators (2.4) it is natural to expect that the main decay modes are $G_H \rightarrow b\bar{b}$ and $G_H \rightarrow gg$, with the ratio of these decay widths given in eq. (2.15). This ratio depends on several unknown parameters, so that in what follows we will treat the branching fraction for $G_H \rightarrow b\bar{b}$ as a free parameter.

2.3 CP violation in B_s mixing from scalar octet exchange

The operators (2.4) can induce tree-level flavor-changing neutral-current (FCNC) processes mediated by the scalar octet. In particular, integrating out the octet gives the following $\Delta B = 2$ terms in the effective Lagrangian:

$$\frac{v^2}{2M_{G_H}^2 m_\psi^2} \left(\eta'_3 \lambda'_2 \bar{b}_L T^a s_R + \eta'_2 \lambda'_3 \bar{b}_R T^a s_L \right)^2, \quad (2.17)$$

where the quark fields shown here are mass eigenstates, and η'_i and λ'_i are the Yukawa couplings of eqs. (2.8) and (2.9) transformed to the mass eigenstate basis.

Tree-level exchange of octet bosons has been proposed before [22] as a possible explanation for the anomalous dimuon charge asymmetry reported by the D0 Collaboration [23]. The novel feature in our case is that the scalar octet has dimension-5 couplings to the standard model quarks. Nevertheless, we now show that the effect of G_H is large enough if the G_H mass is below a few hundred GeV.

Using the relation $T_{ij}^a T_{kl}^a = -\delta_i^j \delta_k^l / 6 + \delta_i^l \delta_j^k / 2$, and substituting the matrix elements of the ensuing operators from ref. [24], we find the matrix element of the Hamiltonian associated with operator (2.17):

$$\begin{aligned} \langle \bar{B}_s | \mathcal{H}^{\text{NP}} | B_s \rangle = & \frac{v^2 M_{B_s}^4 f_{B_s}^2}{96 M_{G_H}^2 m_\psi^2 (m_b + m_s)^2} \left[\left((\eta'_2 \lambda'_3)^2 + (\eta'_3 \lambda'_2)^2 \right) \left(B_3 + \frac{5}{3} B_2 \right) \right. \\ & \left. + 4\eta'_3 \lambda'_2 \eta'_2 \lambda'_3 (B_5 - B_4) \right] . \end{aligned} \quad (2.18)$$

The decay constant, computed on the lattice with 2+1 flavors [25], is $f_{B_s} = 231 \pm 15$ MeV. The “bag” parameters have been estimated on the lattice in the quenched approximation [24]: $B_2 = 0.80$, $B_3 = 0.93$, $B_4 = 1.16$ and $B_5 = 1.75$.

If $\eta'_3 \lambda'_2$ or $\eta'_2 \lambda'_3$ have complex phases, then the G_H exchange induces CP violation in $B_s - \bar{B}_s$ mixing. Parametrizing

$$\langle \bar{B}_s | \mathcal{H}^{\text{NP}} | B_s \rangle \equiv \left(C_{B_s} e^{-i\phi_s} - 1 \right) 2M_{B_s} (M_{12}^{\text{SM}})^* \quad (2.19)$$

where $M_{12}^{\text{SM}} \simeq (9.0 \pm 1.4) \text{ ps}^{-1}$ [22, 26], and using eq. (2.18) we find that the vectorlike quark mass required to induce the CP-violating phase ϕ_s is

$$m_\psi = \frac{v f_{B_s} M_{B_s}^{3/2} \left| 2.3 [(\eta'_2 \lambda'_3)^2 + (\eta'_3 \lambda'_2)^2] + 2.4 \eta'_3 \lambda'_2 \eta'_2 \lambda'_3 \right|^{1/2}}{8 M_{G_H} (m_b + m_s) (3 M_{12}^{\text{SM}})^{1/2} (C_{B_s}^2 + 1 - 2 C_{B_s} \cos \phi_s)^{1/4}}. \quad (2.20)$$

The B_s mass difference of $\Delta M_s = 17.70 \pm 0.08 \text{ ps}^{-1}$, obtained from combining in quadrature the CDF [27] and LHCb [28] measurements, gives the constraint $C_{B_s} = 0.98 \pm 0.17$. The D0 dimuon asymmetry requires ϕ_s near $-\pi/2$. For illustration, let us assume $\eta'_3 \lambda'_2 = \eta'_2 \lambda'_3$, $C_{B_s} = 1$ and $\phi_s = -\pi/2$, which gives

$$m_\psi \approx 1.1 \text{ TeV} \frac{|\eta'_3 \lambda'_2|}{10^{-2}} \frac{100 \text{ GeV}}{M_{G_H}}. \quad (2.21)$$

Thus, for $|\eta'_3 \lambda'_2| = |\eta'_2 \lambda'_3| \sim 10^{-2}$, the D0 dimuon asymmetry suggests that the vectorlike quark has a mass within the reach of the LHC.

3 Tevatron phenomenology of scalar octets

Scalar octets are produced in pairs at hadron colliders (see figure 1) and decay with a large branching fraction into $b\bar{b}$. The partonic cross sections are given by [8, 16, 17].

$$\sigma(q\bar{q} \rightarrow G_H G_H) = \frac{2\pi\alpha_s^2}{9\hat{s}} \beta^3, \quad (3.1)$$

$$\sigma(gg \rightarrow G_H G_H) = \frac{3\pi\alpha_s^2}{32\hat{s}} \left[27\beta - 17\beta^3 + 3(\beta^4 + 2\beta^2 - 3) \ln \left(\frac{1+\beta}{1-\beta} \right) \right], \quad (3.2)$$

where $\beta = (1 - 4M_{G_H}^2/\hat{s})^{1/2}$ is the velocity of the octet in the center of mass frame. Thus, the signal consists of a pair of narrow $b\bar{b}$ resonances of same mass, M_{G_H} . Including the interactions (2.2) into the MadGraph/MadEvent v5 package [29, 30], we find that the processes shown in figure 1, convoluted with the CTEQ 6L1 [31] parton distribution functions (PDFs), give the cross section for $p\bar{p} \rightarrow G_H G_H$ at the Tevatron shown by the solid line in figure 4. The factorization and renormalization scales used by default in MadGraph/MadEvent are equal and depend on the event: $(M_{G_H}^2 + p_T^2(G_1))^{1/2} (M_{G_H}^2 + p_T^2(G_2))^{1/2}$, where $p_T(G_1)$ and $p_T(G_2)$ are the transverse momenta of the two G_H produced in each event. This cross section depends only on M_{G_H} , and is consistent with the result shown in figure 3 of ref. [7].

3.1 $3b$ versus $4b$ signals

The QCD background is dramatically reduced by selecting events containing four jets, all b tagged, and then imposing that two of them have an invariant mass close to that of the other two b jets [7]. We point out, however, that the sensitivity to the pair of octets could be improved if, instead of requiring four b jets, only the three jets of largest transverse momenta are b tagged and required to pass the cuts. In this case only two of the b jets will

Figure 4. Leading-order cross section for production of a G_H pair (solid line) at the Tevatron. Also shown are the cross sections after cuts for producing a pair of G_H followed by each decaying with 100% branching fraction into $b\bar{b}$ and requiring that 3 (dashed line) or 4 (dotted line) jets pass the cuts and are b -tagged.

M_{G_H} (GeV)	partonic level		radiation included		b -tagging included	
	3j	4j	3j	4j	3b	4b
100	91.1	62.1	76.9	38.9	9.4	1.0
150	95.8	78.1	93.2	65.5	15.2	2.4
200	97.4	83.1	96.9	77.8	18.1	3.0
250	98.2	85.0	98.4	84.3	19.8	3.9
300	98.6	86.8	99.0	87.8	19.9	3.8
350	98.9	86.7	99.4	90.6	20.0	4.2

Table 1. Efficiency in % for the three and four jets to pass the basic cuts ($p_T > 20$ GeV, $|\eta| < 2$): at parton level (columns 2,3), after radiation and detector effects included through PYTHIA and PGS (columns 4,5), and after b tagging by PGS (columns 6,7).

form a resonance and thus the background is substantially higher, but the signal is also larger, such that the $S/\sqrt{B}$ significance may increase.

To see this, first note that the b tagging efficiency is about 50%, leading to a factor of 2 suppression of the $4b$ signal compared to the $3b$ signal. The $4b$ signal is further suppressed because the 4th jet is rather soft, especially for low M_{G_H} , and does not always pass the basic cuts, such as $p_T > 20$ GeV, imposed by the CDF [14] and D0 [35] searches. To estimate the ensuing suppression of the $4b$ signal, we use MadGraph/MadEvents to generate parton-level events at tree level for $p\bar{p} \rightarrow G_H G_H \rightarrow b\bar{b}b\bar{b}$ at a center-of-mass energy of 1.96 TeV. Imposing the basic cuts used by CDF [14], namely $p_T > 20$ GeV and $|\eta| < 2$ for each b jet, we obtain the result shown in columns 2 and 3 of table 1.

Initial and final state radiation further soften the 4th jet. We estimate this effect by processing the parton-level events with PYTHIA [32] for showering and hadronization, and

Figure 5. Fraction of signal events having the invariant mass of the i th and j th jets, $m_{ij} \equiv m(b_i, b_j)$, within 20% of the octet mass M_{G_H} . The fractions for combinations with the 4th jet are below 1% and are not shown here.

PGS [34] (with the default CDF detector card) for detector simulation. The result is shown in columns 4 and 5 of table 1. The b -tagging efficiency also decreases for softer jets, so that the efficiency for four b jets to pass the basic cuts is significantly smaller than that for three b jets, especially for low M_{G_H} (see the last column of table 1). In figure 4 we show the signal cross sections for 3 and 4 b -tagged jets after basic kinematic cuts.

3.2 Properties of the $3b$ signal

Given that the $3b$ signal is larger than the $4b$ signal by a factor ranging between 9 and 5 when M_{G_H} varies between 100 and 350 GeV (see last two columns of table 1), it is useful to focus on the $3b$ final state. A clear feature of this signal is that two of the three b jets form a narrow resonance of mass close to M_{G_H} . Let us label these jets by b_i , $i = 1, 2, 3$. Within each event we take b_1, b_2, b_3 to be the jets of largest, 2nd-largest and 3rd-largest transverse momenta (p_T), respectively. In the limit where M_{G_H} is very large, close to $\sqrt{s}/2$, the octets are produced mostly at rest and the p_T of a jet is fixed by the angle between the two jets from the decay of an octet, so that b_1 and b_2 form a mass peak.

For masses of interest at the Tevatron, between 100 and 400 GeV, the p_T 's of the octets are comparable with the p_T 's of the jets in the rest frame of the corresponding octet. As a result, the mass peak is formed in some events by b_1 and b_2 while in other events by b_1 and b_3 , or even by the b_2 and b_3 jets. We refer to these three types of events as (b_1, b_2) , (b_1, b_3) , and (b_2, b_3) respectively. To estimate the fractions of events falling into these categories, we impose a $|m_{ij} - M_{G_H}| < 0.2M_{G_H}$ cut, where $m_{ij} \equiv m(b_i, b_j)$ is the invariant mass of the b_i and b_j jets. We show the efficiency of the combination of basic and invariant mass cuts in figure 5. Given that (b_1, b_2) and (b_1, b_3) have comparable efficiencies, it is useful to use both m_{12} and m_{13} as discriminants between the octet signal and the QCD background. Note that the ratio of the efficiencies for (b_1, b_2) and (b_1, b_3) increases with M_{G_H} , which is explained by the large M_{G_H} argument presented above.

So far there have been no collider searches for the pair-produced octet. However, the CDF and D0 Collaborations have searched for a $3b$ signal predicted in the MSSM at large $\tan\beta$: a b quark from the proton (in the 5-flavor PDF scheme) radiates off a heavy Higgs boson which then decays into a $b\bar{b}$ pair. This is the same final state as the $3b$ one due to octets discussed here, but we will show that the kinematic distributions are different.

The CDF search [14], with 2.2 fb^{-1} of data, shows an intriguing excess of events with m_{12} in the 125–155 GeV range. The probability for this excess to arise from a fluctuation of the standard model background is 0.9%, and increases to 5.7% when the whole range of invariant masses is taken into account.

The D0 search [35], with 5.2 fb^{-1} of data, rules out the presence of MSSM Higgs bosons with couplings large enough to account for the CDF excess. The tension between the CDF and D0 results may be due to a statistical fluctuation. An alternative explanation, however, is that the D0 search is less sensitive to the octet-induced signal; this is a consequence of the optimization of the signal within the D0 search for the MSSM Higgs bosons through the use of a likelihood discriminant. Some of the kinematic variables included in the likelihood discriminant (e.g., the angular separation $\Delta\phi$ between the two b jets that are most likely to originate from the decay of the new particle) may discriminate against events due to $G_H G_H$ production.

Motivated by the CDF search, we first study the kinematics of an octet having a mass $M_{G_H} = 140\text{ GeV}$, and impose the basic cuts $p_T > 20\text{ GeV}$ and $|\eta| < 2$ for each b jet. In figure 6, left panel, we show the number of events as a function of m_{12} and m_{13} . Although the peak of the m_{12} distribution is close to M_{G_H} , the spectrum is broad such that our mass window cut $|m(b_1, b_2) - M_{G_H}| < 0.2M_{G_H}$ keeps only around 50% of signal events. The m_{13} distribution also shows a clear peak, which is narrower than the m_{12} one, and at a lower invariant mass because radiation effects make the 3rd jet softer. The transverse energy distributions of the two leading jets (see the right panel of figure 6) also have clear peaks. However, those peaks appear to be too broad to provide a convincing explanation for the excess of the leading two jet transverse energy distributions at CDF [15]. We return to this issue in section 4.2.

To study the octet discovery limit at the Tevatron, we impose high p_T cuts on the three leading b jets, of 150, 120, and 100 GeV, respectively. For $M_{G_H} = 300\text{ GeV}$, the signal has an acceptance around 3%. After imposing the mass window cut there are approximate 6 signal events for 20 fb^{-1} with approximately zero background. Thus, using only the three leading b jets, we estimate that the Tevatron can discover scalar octets of mass up to $\sim 300\text{ GeV}$.

3.3 Effects of bound states

Since the octet is much heavier than the QCD confinement scale, the QCD interaction approximately generates a Coulomb-like potential between two octets. There are three attractive channels which lead to bound states: 1 , 8_A (antisymmetric) and 8_S (symmetric) from $8 \otimes 8$, because they have positive differences of the quadratic Casimir between the initial and final states. Defining $C_f = [C_2(8) + C_2(8) - C_2(f)]/2$, we have $C_1 = 3$ and

Figure 6. Kinematic distributions resulting from $G_H G_H$ production at the Tevatron due to gluon couplings, for an octet mass of 140 GeV (cross section is 6.6 pb) and a $G_H \rightarrow b\bar{b}$ branching fraction of 50%. Left panel: invariant mass distributions of the two leading b jets (dotted red line), and of the 1st and 3rd leading b jets (solid blue line). Right panel: transverse energy distributions of the 1st (dashed black line) and 2nd (solid purple line) leading b jets.

$C_{8_A} = C_{8_S} = 3/2$. The potential from 1-gluon exchange is

$$V(r) = -C_f \frac{\alpha_s(1/r)}{r}, \quad (3.3)$$

where α_s is evaluated at the energy scale of the Bohr radius r of the bound state. At one loop,

$$\alpha_s(1/r) = \frac{1}{\alpha_s^{-1}(M_Z) - b_0 \ln(rM_Z)}. \quad (3.4)$$

Here, $b_0 = (11N_c - 2N_f)/(6\pi)$ with $N_f = 5$ for $m_b < 1/r < m_t$.

Solving the Schrödinger equation numerically for the potential (3.3), using the above running coupling, we find $r_0 \approx 0.07 \text{ GeV}^{-1}$, the binding energy $E_b(8) \approx 2.0 \text{ GeV}$, and the squared wavefunction at the origin $|\psi_8(0)|^2 \approx (9.2 \text{ GeV})^3$. For the singlet channel we find $E_b(1) \approx 6.5 \text{ GeV}$ and $|\psi_1(0)|^2 \approx (16.9 \text{ GeV})^3$.

For the pure Coulomb potential, with $\alpha_s = \alpha_s(1/r_0)$ fixed, we find $E_b = C_f^2 \alpha_s^2 M_{G_H}/4 = 2.2 \text{ GeV}$ and $|\psi(0)|^2 = C_f^3 \alpha_s^3 M_{G_H}^3/(8\pi) \approx (12.4 \text{ GeV})^3$. Hence, the effect of the running is about 10%; we will use the running potential in what follows.

Our calculation of the bound-state wavefunction and binding energy has neglected the octet width. Let us now estimate the bound-state effects on octet production for $\Gamma(G_H) \ll E_b$. The S -wave bound states of two scalar fields have the quantum numbers, J^{PC} , of 0^{++} for the singlet, 0^{+-} for 8_A and 0^{++} for 8_S . Their couplings to quarks are suppressed by the quark mass because of the chirality flip, so the dominant production at hadron colliders is from gluons.

The falling apart width of a bound state is $\Gamma(\mathcal{B}_f \rightarrow 4b) \approx 2\Gamma_G$. The $\mathcal{B}_f \rightarrow gg$ widths are calculated using the non-relativistic relation in the low-velocity limit, $\Gamma(\mathcal{B}_f \rightarrow X) = v\sigma(G_H G_H \rightarrow X)|\psi(0)|^2$ [36–38]:

$$\Gamma(1 \rightarrow gg) = \frac{9\pi\alpha_s^2(M_{\mathcal{B}_1})}{2M_G^2} |\psi_1(0)|^2, \quad \Gamma(8_S \rightarrow gg) = \frac{9\pi\alpha_s^2(M_{\mathcal{B}_8})}{8M_G^2} |\psi_{8_S}(0)|^2, \quad (3.5)$$

where the bound-state mass is $M_{\mathcal{B}_f} = 2M_{G_H} - E_b(f)$. Note that 8_A does not decay into two gluons. For $M_{G_H} = 150$ GeV, we have $\Gamma(1 \rightarrow gg) \approx 0.04$ GeV and $\Gamma(8_S \rightarrow gg) \approx 0.002$ GeV.

In the narrow width approximation, the partonic production cross sections of the bound states are

$$\hat{\sigma}_{gg \rightarrow 1}(\hat{s}) = \frac{\pi^2 \Gamma(1 \rightarrow gg)}{8 M_{\mathcal{B}_1}} \delta(\hat{s} - M_{\mathcal{B}_1}^2), \quad \hat{\sigma}_{gg \rightarrow 8_S}(\hat{s}) = \frac{\pi^2 \Gamma(8_S \rightarrow gg)}{M_{\mathcal{B}_8}} \delta(\hat{s} - M_{\mathcal{B}_8}^2) \quad (3.6)$$

Using the gluon PDF, we obtain the production cross section at the Tevatron:

$$\sigma(p\bar{p} \rightarrow \mathcal{B}_f) = \frac{\pi^2 \Gamma(1 \rightarrow gg)}{8 M_{\mathcal{B}_f} s} \int_{M_{\mathcal{B}_f}^2/s}^1 \frac{dx}{x} f_g(x) f_g\left(\frac{M_{\mathcal{B}_f}^2}{xs}\right). \quad (3.7)$$

For $M_{G_H} = 150$ GeV, using the Mathematica MSTW 2008 PDFs [39], the cross sections are $\sigma(p\bar{p} \rightarrow 1) \approx 190$ fb and $\sigma(pp \rightarrow 8_S) \approx 57$ fb, which are a few percent of the octet pair production and can be neglected.

3.4 Supersymmetric Higgs bosons

Let us now compare the kinematic distributions of the multi- b -jet final states arising from scalar octet production with those due to the Higgs bosons of the MSSM. In Two-Higgs-doublet models, such as the Higgs sector of the MSSM, there are two color-singlet and electrically-neutral spin-0 particles (the heavy Higgs bosons H^0 and A^0) that may have large couplings to the b quark. In the 5-flavor PDF scheme, where the proton includes a b -quark, the emission of a heavy Higgs boson from a b quark line leads to a $3b$ final state (see diagram next to table 2). Equivalently, in the 4-flavor PDF scheme, where there is no b inside the proton, production of a $b\bar{b}$ pair followed by emission of a heavy Higgs boson gives four b jets that may appear as a $3b$ final state when one of the jets does not pass the cuts. At large $\tan\beta$ and for $M_A \gg M_Z$, the production cross sections of H^0 and A^0 are approximately equal, and are related to the standard model Higgs production by $\sigma(b\bar{b}H^0) = \sigma(b\bar{b}A^0) = \tan^2\beta \sigma(b\bar{b}h_{\text{SM}}^0)$ [40].

In order to study the multi- b Higgs signal, we impose the same cuts as in the case of scalar octets: $p_T > 20$ GeV and $|\eta| < 2$ for each b jet. Compared to the octet case, fewer events pass these cuts because one of the jets does not arise from the decay of a heavy particle. For example, the fourth b jet has ~ 3 times smaller chance to pass the basic cuts than in the octet case (see rows 2 and 3 of table 2). The invariant mass of the leading and the third jets is also less likely to be within the mass window $|m_{13} - M_{G_H}| < 0.2M_{G_H}$ than in the octet case (see last two rows of table 2). Thus, the m_{13} distribution is a good discriminant between the octet and the MSSM Higgs bosons, especially when the resonance is heavy. We show the invariant mass distributions for the b_1, b_2 and b_1, b_3 jet pair in figure 7 for heavy Higgs bosons of mass 140 GeV and a leading-order production cross section (summed over A^0 and H^0) corresponding to $\tan\beta = 40$. Comparing figures 7 and 6, we can see that the MSSM Higgs bosons lead to an m_{13} distribution with a peak that is both lower and located at a smaller value than the peak due to the scalar octet.

If a large excess of events over the standard model background will be established in the $3b$ final state, then an MSSM interpretation will typically require an excess of events in

M (GeV)	150	200	250	300	350
G_H	15.8	16.6	19.7	19.1	21.0
H^0, A^0	4.3	6.1	7.4	7.7	7.7
G_H	1.0	1.0	0.86	0.79	0.73
H^0, A^0	0.50	0.25	0.16	0.10	0.10

Table 2. Ratio (in %) of the numbers of events with $4b$ or $3b$, for the MSSM (representative process shown on the right-hand side) at $\tan\beta = 40$ (3rd row) and QCD $G_H G_H$ production (2nd row). Also, the ratio of numbers of events that pass the m_{12} or m_{13} invariant-mass window cut, $|m_{ij} - M_{G_H}| < 0.2M_{G_H}$, for the MSSM (5th row) and for the octet (4th row).

Figure 7. Same as figure 6, but for MSSM Higgs production at $\tan\beta = 40$ (the leading-order production cross section is around 5.4 pb).

the $b\tau^+\tau^-$ final state due to the decays of H^0 and A^0 into tau leptons (albeit the correlation is sensitive to loop corrections [41]). By contrast, G_H can not decay into leptons-only final states because it is a color octet, so that no $b\tau^+\tau^-$ excess is predicted in the octet models.

4 Renormalizable coloron model

Let us now analyze a minimal renormalizable model that includes a spin-1 color-octet particle [10–13], called “coloron”. We will show that this model includes a scalar octet, identical with the G_H studied in sections 2 and 3 except that here it can be resonantly produced in pairs, through an s -channel coloron, G'_μ . The gauge symmetry of this model is an extension of QCD, $SU(3)_1 \times SU(3)_2$. This symmetry is spontaneously broken down to the QCD gauge group $SU(3)_c$ by the VEV of a complex scalar field, Σ , which transforms as $(3, \bar{3})$ under $SU(3)_1 \times SU(3)_2$.

4.1 Interactions and masses

The most general renormalizable potential of Σ is

$$V(\Sigma) = -m_\Sigma^2 \text{Tr}(\Sigma\Sigma^\dagger) - \mu (\det \Sigma + \text{H.c.}) + \frac{\lambda}{2} \left[\text{Tr}(\Sigma\Sigma^\dagger) \right]^2 + \frac{\kappa}{2} \text{Tr}(\Sigma\Sigma^\dagger\Sigma\Sigma^\dagger) \quad , \quad (4.1)$$

where without loss of generality we take $\mu > 0$. Note that the second term above is $\text{SU}(3)_1 \times \text{SU}(3)_2$ invariant because it can be written as

$$\det \Sigma = \frac{1}{6} \epsilon^{ijk} \epsilon^{i'j'k'} \Sigma_{ii'} \Sigma_{jj'} \Sigma_{kk'} \quad . \quad (4.2)$$

We assume $m_\Sigma^2 > 0$ so that Σ acquires a VEV:

$$\langle \Sigma \rangle = \frac{f_\Sigma}{\sqrt{6}} \mathbb{I}_3 = \frac{\sqrt{4(\kappa + 3\lambda) m_\Sigma^2 + \mu^2 + \mu}}{2(\kappa + 3\lambda)} \mathbb{I}_3 \quad , \quad (4.3)$$

where $\mathbb{I}_3$ is the unit 3×3 matrix. The potential is bounded from below provided $3\lambda + \kappa > 0$.

Expanding Σ around this vacuum, we find that its 18 degrees of freedom (Σ is a 3×3 complex matrix) are grouped into four real scalar fields: two octets (G_H^a and G_G^a , $a = 1, \dots, 8$) and two singlets (ϕ_R and ϕ_I) under $\text{SU}(3)_c$,

$$\Sigma = \frac{1}{\sqrt{6}} (f_\Sigma + \phi_R + i\phi_I) \mathbb{I}_3 + (G_H^a + iG_G^a) T^a \quad . \quad (4.4)$$

G_G^a are the Nambu-Goldstone bosons associated with the broken generators of $\text{SU}(3)_1 \times \text{SU}(3)_2$, and become the longitudinal degrees of freedom of the coloron. Previous studies of this model have assumed that the other degrees of freedom of Σ are heavy enough to be neglected². It turns out, however, that within a renormalizable model valid for a range of scales above f_Σ , the G_H^a , ϕ_R and ϕ_I scalars are lighter than the coloron. Thus, it is interesting to study their properties. We refer to the renormalizable model that includes all the scalar particles contained in Σ and the $\text{SU}(3)_1 \times \text{SU}(3)_2$ gauge symmetry as ReCoM.

For $\mu \rightarrow 0$, there is a $\text{U}(1)_1 \times \text{U}(1)_2$ global symmetry broken by $\langle \Sigma \rangle$ down to the diagonal $\text{U}(1)$ subgroup, and the associated Nambu-Goldstone boson is ϕ_I , which becomes massless. For any μ , the squared mass of ϕ_I is given by

$$M_{\phi_I}^2 = \sqrt{\frac{3}{2}} \mu f_\Sigma \quad . \quad (4.5)$$

For $\kappa \rightarrow 0$ and $\mu \rightarrow 0$ the potential (4.1) has a global $\text{SO}(18)$ symmetry, which is spontaneously broken by the Σ VEV down to $\text{SO}(17)$, so that G_H and ϕ_I are massless Nambu-Goldstone bosons. Hence, the quartic term in $V(\Sigma)$ proportional to λ does not contribute to the G_H mass. For any κ and μ , the squared mass of G_H is

$$M_{G_H}^2 = \frac{1}{3} (2M_{\phi_I}^2 + \kappa f_\Sigma^2) \quad . \quad (4.6)$$

²An exception is the $\text{SU}(3)_1 \times \text{SU}(3)_2$ linear σ -model description of heavy-light mesons in QCD [42, 43].

The mass of ϕ_R also follows from eq. (4.1):

$$M_{\phi_R}^2 = \frac{1}{3} [(\kappa + 3\lambda) f_\Sigma^2 - M_{\phi_I}^2] \quad . \quad (4.7)$$

eq. (4.3) implies $f_\Sigma \geq \sqrt{6}\mu/(\kappa + 3\lambda)$, so that $M_{\phi_R}^2 \geq 0$.

The kinetic term of Σ , normalized such that all its component fields shown in eq. (4.4) have canonical kinetic terms, is given by

$$\text{Tr} \left(D_\mu \Sigma^\dagger D^\mu \Sigma \right) \quad , \quad (4.8)$$

where the $\text{SU}(3)_1 \times \text{SU}(3)_2$ covariant derivative is

$$D^\mu \Sigma = \partial^\mu \Sigma - i h_1 G_1^{\mu a} T^a \Sigma + i h_2 G_2^{\mu a} \Sigma T^a \quad . \quad (4.9)$$

The kinetic term leads to a mass-square matrix for the two gauge fields, $G_1^{\mu a}$ and $G_2^{\mu a}$, proportional to f_Σ^2 . Upon diagonalization, one linear combination becomes the massless QCD gluon,

$$G^\mu = \cos \theta G_1^\mu + \sin \theta G_2^\mu \quad , \quad (4.10)$$

while the orthogonal linear combination is a massive spin-1 octet, the coloron:

$$G'^\mu = \sin \theta G_1^\mu - \cos \theta G_2^\mu \quad . \quad (4.11)$$

The mixing angle depends only on the $\text{SU}(3)_1 \times \text{SU}(3)_2$ gauge couplings, h_1 and h_2 : $\tan \theta = h_1/h_2$. The QCD gauge coupling is given by $g_s = h_1 \cos \theta = h_2 \sin \theta$, and the mass of the coloron is

$$M_{G'} = \frac{\sqrt{2} g_s}{\sqrt{3} \sin 2\theta} f_\Sigma \quad . \quad (4.12)$$

This mass is larger than the VEV f_Σ if $\tan \theta \lesssim 0.65$. For $\tan \theta \ll 1$ and $\lambda, \kappa \lesssim O(1)$, the coloron is heavier than the scalars. For example, the set of parameters

$$\tan \theta = 0.12 \quad , \quad f_\Sigma = 75 \text{ GeV} \quad , \quad \mu = 275 \text{ GeV} \quad , \quad \kappa = \lambda = 1.4 \quad , \quad (4.13)$$

and $\alpha_s(M_{G'}) \simeq 0.1$ gives the following mass spectrum:

$$M_{G'} \approx 290 \text{ GeV} \quad , \quad M_{G_H} \approx 140 \text{ GeV} \quad , \quad M_{\phi_I} \approx 159 \text{ GeV} \quad , \quad M_{\phi_R} \approx 46 \text{ GeV} \quad . \quad (4.14)$$

Substituting the decomposition of Σ , eq. (4.4), into the kinetic term (4.8), we find the following interactions of a coloron and two scalars:

$$\frac{\sqrt{2} g_s}{\sqrt{3} \sin 2\theta} G'^\mu_a (\phi_I \partial^\mu G_H^a - G_H^a \partial^\mu \phi_I) + \frac{g_s}{\tan 2\theta} f^{abc} G'^\mu_a G_H^b \partial^\mu G_H^c \quad . \quad (4.15)$$

These interactions induce G' decays into scalars with tree-level widths (in the case where the two-body decays are kinematically open) given by:

$$\begin{aligned} \Gamma(G'_\mu \rightarrow G_H G_H) &= \frac{\alpha_s}{8 \tan^2 2\theta} M_{G'} \left(1 - \frac{4 M_{G_H}^2}{M_{G'}^2} \right)^{3/2} \quad , \\ \Gamma(G'_\mu \rightarrow G_H \phi_I) &= \frac{\alpha_s}{18 \sin^2 2\theta} M_{G'} \left(1 - 2 \frac{M_{\phi_I}^2 + M_{G_H}^2}{M_{G'}^2} + \frac{(M_{\phi_I}^2 - M_{G_H}^2)^2}{M_{G'}^4} \right)^{3/2} \quad . \end{aligned} \quad (4.16)$$

If all standard model quarks belong to the fundamental representation of $SU(3)_1$, then G' couples to them as

$$g_s \tan \theta \bar{q} \gamma^\mu T^a G'_\mu{}^a q \quad . \quad (4.17)$$

The width of the coloron decay into quark pairs is

$$\Gamma(G'_\mu \rightarrow q \bar{q}) = \frac{\alpha_s}{6} \tan^2 \theta M_{G'} \left(1 - \frac{4m_q^2}{M_{G'}^2} \right)^{1/2} , \quad (4.18)$$

where we have not summed over quark flavors. The kinematic suppression in the last factor is relevant for $G'_\mu \rightarrow t \bar{t}$, if $M_{G'} > 2m_t$. The coloron effects on top-quark physics are rather sensitive to new quarks that mix with the top [44] and to the top charges under $SU(3)_1 \times SU(3)_2$ [10, 11].

QCD corrections to the above decay widths may be large, perhaps of the order of 50%, but computing them is beyond the scope of this paper. Assuming that the QCD corrections to $G'_\mu \rightarrow G_H G_H$ and $G'_\mu \rightarrow q \bar{q}$ decays are of the same size, the parameters given in eq. (4.13) lead to branching fractions of 76% and 24% for $G_H G_H$ and jj , respectively.

4.2 ReCoM and the CDF excess in the $3b$ final state

The total width of G'_μ is rather small, typically less than one percent of its mass for $\tan \theta \lesssim 0.3$ and for a G'_μ mass not more than a few percent above the $G_H G_H$ and $G_H \phi_I$ thresholds. We can then use the narrow width approximation to estimate the cross section for producing a G'_μ in the s -channel:

$$\sigma(q \bar{q} \rightarrow G') \approx \frac{8\pi^2 \alpha_s \tan^2 \theta}{9M_{G'}} \delta(\sqrt{s} - M_{G'}) \quad . \quad (4.19)$$

Convoluting this partonic cross section with the MSTW [39] PDFs, and then multiplying by the branching fractions derived from eqs. (4.16) and (4.18) we find the total cross sections shown in figure 8. The QCD corrections to these processes are also likely to be sizable, and need to be computed in the future.

The process $p\bar{p} \rightarrow G'_\mu \rightarrow jj$ is constrained by the CDF search for dijet resonances [45]: for $M_{G'} \approx 290$ GeV, the limit on the cross section times acceptance is about 100 pb. Given that the G'_μ production is proportional to $\tan^2 \theta$, figure 8 indicates that the CDF limit implies $\tan \theta \lesssim 0.2$ for a G_H mass of 140 GeV.

The decays of G_H into $b\bar{b}$ or gg proceed through higher-dimensional operators, similar to the case discussed in section 2. For example, a vectorlike down-type quark ψ , which transforms as a triplet under $SU(3)_2$, couples to Σ and a b quark:

$$\eta_3 \Sigma \bar{b}_R \psi_L + \text{H.c.} \quad (4.20)$$

Integrating out ψ we obtain at tree level the first operator of eq. (2.5), which mediates the $G_H \rightarrow b\bar{b}$ decay, suppressed by two powers of the η_3 coupling. A 1-loop diagram as in figure 3 leads to $G_H \rightarrow gg$. This decay also occurs at one loop independently of ψ , as shown in figure 2, due to the $\det \Sigma$ term from eq. (4.1).

Figure 8. Production cross section of G'_μ at the Tevatron times branching fractions, computed in the narrow width approximation and at leading order. The cross section for $p\bar{p} \rightarrow G'_\mu \rightarrow G_H G_H$ is given by the solid (red) line, and that for $p\bar{p} \rightarrow G'_\mu \rightarrow jj$ by the dotted (blue) line. The masses are taken to be 140 GeV for G_H and 290 GeV for G'_μ , and the decay $G'_\mu \rightarrow G_H \phi_I$ is assumed to be kinematically forbidden.

Figure 9. Coloron (G'_μ) resonance decaying to a pair of scalar octets (G_H), each giving rise to a $b\bar{b}$ resonance.

Compared to the QCD pair production of two G_H scalars, the kinematic distributions of b jets arising from $G' \rightarrow G_H G_H \rightarrow 4b$ (see diagram in figure 9) are changed dramatically. Especially when the mass of G' is close to twice of the G_H mass, those two G_H 's are mainly produced at rest and the jet E_T distributions are more peaked, as can be seen by comparing the right-hand panels of figures 6 and 10. As shown in figure 8, the resonant production of a G_H pair is an order of magnitude larger than the QCD $G_H G_H$ cross section (of about 4.6 pb for $M_{G_H} = 140$ GeV, as can be seen in figure 4). Hence, a smaller branching fraction (around 20%) of $G_H \rightarrow b\bar{b}$ gives a number of $4b$ events comparable to that due to the supersymmetric Higgs bosons for a mass of 140 GeV and $\tan\beta = 40$.

The CDF $3b$ search [14] shows an excess in the m_{12} distribution, as mentioned in section 3.2, and in addition the shapes of the E_T distributions of the b jets are shifted towards larger values compared to the standard model background [15]. The dotted (blue) line in figure 11 shows the background from the best CDF fit to the data, and the shown data points include only statistical errors. The 40–50 GeV bin of the leading-jet E_T distribution

Figure 10. Same as figure 6, but for resonant $G_H G_H$ production via G' and a $G_H \rightarrow b\bar{b}$ branching fraction of 18%. The production cross section of $p\bar{p} \rightarrow G' \rightarrow G_H G_H$ is 50 pb for $M_{G_H} = 140$ GeV, $M_{G'} = 290$ GeV and $\tan \theta = 0.12$.

Figure 11. Comparison of the signal-plus-background distributions with CDF data [15]. The solid (red) line is a combination of the background (multiplied by 0.93) from CDF and both the resonant (via a G'_μ) and QCD production of two G_H scalars. The branching fraction of $G_H \rightarrow b\bar{b}$ is taken to be 18% as a result of fitting to the m_{12} and E_T distributions of the two leading jets. The dotted (blue) line, taken from [15], is the CDF background-only fit to the data. The black points are CDF data with only statistic error bars. The G'_μ mass is chosen to be 290 GeV and the G_H mass is 140 GeV.

is about 3.5σ below the background, suggesting that the background normalization is too high. The 70 – 80 GeV bin is about 2.5σ above the potentially overestimated background, suggesting the presence of some new heavy particles which produce hard b jets. The E_T distribution of the second jet has similar features.

In our ReCoM there are four parameters relevant for fitting the CDF invariant mass and jet E_T distributions: the G'_μ and G_H masses, the mixing angle θ and the branching fraction of $G_H \rightarrow b\bar{b}$. Since the peaks of the E_T distributions of the two leading b jets match the excess bins (70 – 80 GeV for the first jet and 50 – 60 GeV for the second jet, see figure 11), we fix the masses of G'_μ and G_H to be 290 GeV and 140 GeV, respectively (nearly values with similar splitting, such as 305 GeV and 150 GeV, work equally well). Figure 8

Figure 12. Representative diagrams for W boson production in association with a coloron (G'_μ) decaying to a pair of scalar octets (G_H), each giving rise to a dijet resonance.

shows that the production cross section is insensitive to $\tan\theta$ for $0.1 < \tan\theta < 0.2$; for illustration, we fix $\tan\theta = 0.12$. Taking into account the unknown overall normalization of the background, we fit the m_{12} (first 10 bins) and E_T (first 9 bins for each of the two leading jets) distributions reported by CDF [15]. The total χ^2 reaches the minimum for the background normalization factor of 0.93 and $B(G_H \rightarrow b\bar{b}) = 18\%$. The χ^2 per degree of freedom is 0.75 (p -value of 82%) from our signal-plus-background fit, while the background-only fit has $\chi^2/dof = 1.74$ (p -value of 1%). In our simple fit we have not included systematic errors, and we have not allowed different normalization factors for different components of the background. Nevertheless, this successful fit indicates that it would be important for the CDF and D0 Collaborations to compare their multi- b data with the ReCoM predictions.

The fitted $G_H \rightarrow b\bar{b}$ branching fraction implies that the $G_H \rightarrow gg$ branching fraction is large, around 82%. Thus, ReCoM predicts that the number of $bbjj$ events with equal $b\bar{b}$ and jj invariant masses is about 9 times larger than the number of $4b$ events satisfying the similar equal-mass condition. Another prediction is that the invariant mass of the four jets has a peak close to the coloron mass of 290 GeV.

4.3 Dijet resonances plus a W boson and other signals

Another test of our ReCoM is associated production of G'_μ with weak gauge bosons. Feynman diagrams for producing G'_μ plus W^+ are shown in figure 12. We find that the partonic cross section for this process is

$$\sigma(u\bar{d} \rightarrow G'_\mu W^+) = \frac{8\pi\alpha_s \tan^2\theta}{9\hat{s}^2 \sin^2\theta_W} \left[\frac{\hat{s}^2 + (M_{G'}^2 + M_W^2)^2}{\hat{s} - M_{G'}^2 - M_W^2} \ln\left(\frac{\hat{s} - M_{G'}^2 - M_W^2 + M_\beta^2}{2M_{G'}M_W}\right) - M_\beta^2 \right], \quad (4.21)$$

where

$$M_\beta^2 = \left[(\hat{s} - M_{G'}^2 - M_W^2)^2 - 4M_{G'}^2 M_W^2 \right]^{1/2}. \quad (4.22)$$

The leading-order cross section for the $p\bar{p} \rightarrow G'_\mu W$ process at the Tevatron, computed using MadGraph, is shown in figure 13 for $\tan\theta = 0.12$, and is around 150 fb for $M_{G'} = 290$ GeV.

Figure 13. Leading-order production cross section (solid black line) of G'_μ in association with a W boson at the Tevatron for $\tan \theta = 0.12$. The cross section times the coloron branching fraction is shown for the G'_μ decay into a pair of jets (dotted line), and into $G_H G_H$ (solid red line).

Figure 14. Invariant mass distributions for the two leading jets (m_{12} , solid line) and for the first and third jet (m_{13} , dashed line), in events arising from $G'_\mu + W$ production at the Tevatron. All coloron decay channels and only the $W \rightarrow e\nu, \mu\nu$ decays are included. The first peak is mostly due to the $G'_\mu \rightarrow G_H G_H \rightarrow 4j$ decay (its location is sensitive only to the G_H mass), while the second peak in m_{12} is due to the direct $G'_\mu \rightarrow 2j$ decay. The branching fractions for those two channels are 76% and 24%, respectively, for $M_{G_H} = 140$ GeV, $M_{G'} = 290$ GeV and $\tan \theta = 0.12$. A slightly larger G'_μ mass increases the first peak, and a larger $\tan \theta$ increases the second peak.

Since the coloron can cascade decay into four jets via two G_H scalars, or directly decay into two jets, it is useful to analyze the invariant mass distribution of the two leading jets in the final states that include a W decaying to $\ell\nu$, with $\ell = e, \mu$. Following the CDF search in this channel [46], we impose the $p_T(\text{jets}) > 15$ GeV, $p_T(\ell) > 20$ GeV, $\cancel{E}_T > 25$ GeV cuts

on the signal events, using the MadGraph/MadEvents to Pythia to PGS chain as in section 3. For the signal events from $G'_\mu \rightarrow G_H G_H \rightarrow 4j$, the acceptance for passing those cuts is around 63% for inclusive 2 jets or 3 jets in the final state. For the signal events from $G'_\mu \rightarrow 2j$, the acceptances are around 59% for inclusive 2-jet events and 28% for inclusive 3-jet events.

In figure 14, we show the invariant mass distributions of a pair of jets in the multi-jet-plus- $\ell\nu$ events resulting from a simulation of the ReCoM signal. As expected, the invariant mass of the two leading jets (m_{12}) has two peaks corresponding roughly to the masses of G_H and G'_μ , chosen to be 140 GeV and 290 GeV, respectively. The height of the first peak is relatively insensitive to $\tan\theta$, since $\sigma(p\bar{p} \rightarrow G'_\mu W) \times B(G'_\mu \rightarrow G_H G_H)$ is fairly constant for a range of $\tan\theta$ values, which is similar to the case of single G'_μ production shown by the solid (red) line in figure 8. The height of the second peak is strongly sensitive to $\tan\theta$; especially for small values of $\tan\theta$, the cross section times the $B(G'_\mu \rightarrow 2j)$ branching fraction is proportional to $\tan^4\theta$.

The invariant mass of the leading and third jets (m_{13}) has a single peak (see the dashed line in figure 14), near the G_H mass, because the direct $G'_\mu \rightarrow 2j$ decay does not contribute at leading order to this distribution. The peak in m_{13} is a distinctive feature of ReCoM, allowing to differentiate it from the low-scale technicolor model [47–49] that also predicts final states involving a dijet resonance and a W boson. Even more dramatic would be the observation of two different dijet resonances of equal mass in association with a W , as shown in figure 12, but in that case it is likely that a computation of next-to-leading order effects is necessary, in order to include the case where an extra jet is radiated with a p_T larger than that of the third or fourth jet originating from G_H decays.

G'_μ production in association with a Z boson or photon may also be interesting. For example, Z -plus-jets events at the LHC, with the leading jets forming a resonance and the Z decaying into charged leptons, could allow a sufficient separation of the ReCoM signal from the large standard model background.

Throughout this section we have assumed for simplicity that ϕ_I is too heavy to be produced in coloron decays. However, for a range of ReCoM parameters [see eqs. (4.5) and (4.6)] ϕ_I is lighter than G_H so that $G'_\mu \rightarrow G_H \phi_I$ is the dominant decay mode of the coloron. In that case, a larger coloron production cross section (i.e., a larger $\tan\theta$) is allowed by the dijet searches because the branching fraction for $G'_\mu \rightarrow q\bar{q}$ is suppressed. If ϕ_I decays predominantly into light jets, than the $G'_\mu \rightarrow G_H \phi_I$ decay contributes to the $4j$ and $W + 4j$ signals (increasing the height of the first peak in figure 14) without necessarily affecting the multi- b signal.

5 Conclusions

Four-jet final states at the Tevatron and the LHC are predicted in various theories for physics beyond the standard model. Even though the QCD background is very large, the presence of invariant mass peaks allows the separation of the signal from the background [7, 8]. When the new particles decaying into jets have spin 0, it is natural to expect that a significant fraction of the jets comes from b quarks because the spin-0 couplings to standard

model fermions are typically proportional to mass. Thus, b -tagging can further decrease the background.

In this paper we have investigated the properties of weak-singlet, color-octet scalars (“scalar octets”). Electroweak gauge invariance prevents these scalars from coupling at renormalizable level to standard model fermions, but dimension-5 couplings to quarks are allowed and lead to decays mostly into $b\bar{b}$ and gluon pairs. The exchange of a scalar octet of mass below a few hundred GeV could lead to large CP violation in B_s mixing, as indicated by the D0 dimuon asymmetry (see section 2.3), if a vectorlike quark that induces the dimension-5 couplings has mass below the TeV scale.

The scalar octets are pair produced through their coupling to gluons, which is fixed by QCD gauge invariance, or via an s -channel resonance due to a spin-1 color-octet particle. We have shown that the simplest gauge invariant origin of such a resonance, namely the renormalizable version (“ReCoM”) of the coloron model based on the $SU(3)_1 \times SU(3)_2$ gauge extension of QCD [10–13], automatically includes a large coupling of a coloron to a pair of scalar octets. The parameters in this model that affect the $4b$ signal are the ratio $\tan\theta$ of gauge couplings, which controls both the production and branching fractions of the coloron, the masses of the coloron and scalar octet, and the branching fraction of the scalar octet into $b\bar{b}$. We have focused on the case where only three b jets pass some basic cuts, so that a large part of the signal is preserved, at the expense of not being able to reduce the background as efficiently as when both $b\bar{b}$ are present.

Both D0 and CDF have searched for $3b$ final states present in the MSSM at large $\tan\beta$. The D0 search involves a likelihood discriminant which is optimized for the MSSM topology, and therefore cannot be applied to our $3b$ signal. The CDF preliminary result in the $3b$ final state, using the invariant mass distribution m_{12} of the two leading jets, has some excess of events consistent with a new particle of mass in the 140 – 150 GeV range and decaying to $b\bar{b}$. The background normalizations are taken as free parameters, and are fitted so that the deviation in m_{12} is minimized. However, the shapes of transverse energy distributions of the leading and second jets do not fit well the standard model background [15]. We have shown here that the ReCoM changes the shapes of these distributions such that they agree well with the data (see figure 11). The best fit of the ReCoM to the CDF preliminary results indicates that the branching fraction of the scalar octet into two gluons is approximately 3 times larger than the branching ratio into $b\bar{b}$.

Perhaps this good fit is only an accident, and the background modeling performed by CDF can be modified such that the standard model fit to the data improves. Nevertheless, the success of the ReCoM in describing the CDF data is intriguing enough to warrant its experimental study. Specifically, the CDF Collaboration could fit the background plus the ReCoM signal to the $3b$ data for the m_{12} , E_{T1} and E_{T2} distributions. The D0 Collaboration could use the scalar octet kinematics to define the likelihood discriminant and to check whether the ReCoM is consistent with their observables. Both collaborations could look for a $4b$ signal exhibiting a pair of resonances of equal mass. They could also search for the $bbjj$ signal from two resonances as a further test of the ReCoM.

The cross section for coloron production is large enough so that even the process where a W boson is radiated from the initial state (see figure 12), which has smaller backgrounds,

leads to a sufficient number of events to be tested at the Tevatron. The signatures include a W decaying leptonically and four jets, with the invariant mass distribution for two jets peaking at the same location as for the other two jets, near the scalar octet mass of $140 - 150$ GeV. When the two leading jets come from the same scalar octet, the signature is a W boson plus a dijet resonance, which may explain an excess in the CDF data from an inclusive search in this channel [46]. ReCoM predicts the presence of a second mass peak of the two leading jets, due to the direct decay of the coloron into quarks (see figure 14).

The ATLAS and CMS experiments can search for scalar octets with masses between a few hundred GeV and a couple of TeV, impressively extending the Tevatron reach. In the case of a scalar octet with mass of about $140 - 150$ GeV, relevant for the CDF excess, it is hard to overcome the QCD background at the LHC, so that the Tevatron experiments should attempt to discover or rule out its existence.

Acknowledgments

We have benefited from insightful conversations on experimental aspects with Dan Amidei, Henry Frisch, Jonathan Hays and Tom Wright, and on theoretical aspects with Estia Eichten, Walter Giele, Roni Harnik, Patrick Fox, Gordan Krnjaic, Paul MacKenzie and Michael Peskin. SLAC is operated by Stanford University for the US Department of Energy under contract DE-AC02-76SF00515. Fermilab is operated by Universities Research Association Inc. under contract No. DE-AC02-76CH02000 with the DOE.

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] For a review, see C.T. Hill and E.H. Simmons, *Strong dynamics and electroweak symmetry breaking*, *Phys. Rept.* **381** (2003) 235 [Erratum *ibid.* **390** (2004) 553] [[hep-ph/0203079](#)] [[SPIRES](#)].
- [2] G. Burdman, B.A. Dobrescu and E. Ponton, *Resonances from two universal extra dimensions*, *Phys. Rev. D* **74** (2006) 075008 [[hep-ph/0601186](#)] [[SPIRES](#)].
- [3] C. Kilic, T. Okui and R. Sundrum, *Colored Resonances at the Tevatron: Phenomenology and Discovery Potential in Multijets*, *JHEP* **07** (2008) 038 [[arXiv:0802.2568](#)] [[SPIRES](#)].
- [4] C. Kilic, T. Okui and R. Sundrum, *Vectorlike Confinement at the LHC*, *JHEP* **02** (2010) 018 [[arXiv:0906.0577](#)] [[SPIRES](#)].
- [5] Y. Bai and R.J. Hill, *Weakly Interacting Stable Pions*, *Phys. Rev. D* **82** (2010) 111701 [[arXiv:1005.0008](#)] [[SPIRES](#)].
- [6] T. Plehn and T.M.P. Tait, *Seeking Sgluons*, *J. Phys. G* **36** (2009) 075001 [[arXiv:0810.3919](#)] [[SPIRES](#)].
- [7] B.A. Dobrescu, K. Kong and R. Mahbubani, *Massive color-octet bosons and pairs of resonances at hadron colliders*, *Phys. Lett. B* **670** (2008) 119 [[arXiv:0709.2378](#)] [[SPIRES](#)].

- [8] R.S. Chivukula, M. Golden and E.H. Simmons, *Multi-jet physics at hadron colliders*, *Nucl. Phys. B* **363** (1991) 83 [SPIRES].
- [9] C. Kilic, S. Schumann and M. Son, *Searching for Multijet Resonances at the LHC*, *JHEP* **04** (2009) 128 [arXiv:0810.5542] [SPIRES].
- [10] C.T. Hill, *Topcolor: Top quark condensation in a gauge extension of the standard model*, *Phys. Lett. B* **266** (1991) 419 [SPIRES].
- [11] C.T. Hill and S.J. Parke, *Top production: Sensitivity to new physics*, *Phys. Rev. D* **49** (1994) 4454 [hep-ph/9312324] [SPIRES].
- [12] R.S. Chivukula, A.G. Cohen and E.H. Simmons, *New strong interactions at the Tevatron?*, *Phys. Lett. B* **380** (1996) 92 [hep-ph/9603311] [SPIRES].
- [13] E.H. Simmons, *Coloron phenomenology*, *Phys. Rev. D* **55** (1997) 1678 [hep-ph/9608269] [SPIRES].
- [14] CDF collaboration, *Search for Higgs bosons produced in association with b quarks*, Note 10105, June 2010,
http://www-cdf.fnal.gov/physics/new/hdg//Results_files/results/3b_susyhggs-jun10.
- [15] http://www-cdf.fnal.gov/physics/new/hdg//Results_files/results/3b_susyhggs-jun10/more_plots.html
- [16] A.V. Manohar and M.B. Wise, *Flavor changing neutral currents, an extended scalar sector and the Higgs production rate at the LHC*, *Phys. Rev. D* **74** (2006) 035009 [hep-ph/0606172] [SPIRES].
- [17] M. Gorbunov, T.J. Khoo, D.J. Phalen, A. Pierce and D. Tucker-Smith, *Color-octet scalars at the LHC*, *Phys. Rev. D* **77** (2008) 095003 [arXiv:0710.3133] [SPIRES].
- [18] M.I. Gresham and M.B. Wise, *Color Octet Scalar Production at the LHC*, *Phys. Rev. D* **76** (2007) 075003 [arXiv:0706.0909] [SPIRES].
- [19] Y. Bai and A. Martin, *Topological Pions*, *Phys. Lett. B* **693** (2010) 292 [arXiv:1003.3006] [SPIRES].
- [20] A.R. Zerwekh, C.O. Dib and R. Rosenfeld, *A New signature for color octet pseudoscalars at the CERN LHC*, *Phys. Rev. D* **77** (2008) 097703 [arXiv:0802.4303] [SPIRES].
- [21] T.G. Rizzo, *Gluon final states in Higgs boson decay*, *Phys. Rev. D* **22** (1980) 178 [Addendum *ibid.* **22** (1980) 1824]. [SPIRES].
- [22] B.A. Dobrescu, P.J. Fox and A. Martin, *CP violation in B_s mixing from heavy Higgs exchange*, *Phys. Rev. Lett.* **105** (2010) 041801 [arXiv:1005.4238] [SPIRES].
- [23] D0 collaboration, V.M. Abazov et al., *Evidence for an anomalous like-sign dimuon charge asymmetry*, *Phys. Rev. D* **82** (2010) 032001 [arXiv:1005.2757] [SPIRES].
- [24] D. Becirevic, V. Giménez, G. Martinelli, M. Papinutto and J. Reyes, *B -parameters of the complete set of matrix elements of $\Delta B = 2$ operators from the lattice*, *JHEP* **04** (2002) 025 [hep-lat/0110091] [SPIRES].
- [25] HPQCD collaboration, E. Gamiz, C.T.H. Davies, G.P. Lepage, J. Shigemitsu and M. Wingate, *Neutral B Meson Mixing in Unquenched Lattice QCD*, *Phys. Rev. D* **80** (2009) 014503 [arXiv:0902.1815] [SPIRES].
- [26] A. Lenz and U. Nierste, *Theoretical update of $B_s - \bar{B}_s$ mixing*, *JHEP* **06** (2007) 072 [hep-ph/0612167] [SPIRES].

- [27] CDF collaboration, A. Abulencia et al., *Observation of $B/s0$ anti- $B/s0$ oscillations*, *Phys. Rev. Lett.* **97** (2006) 242003 [[hep-ex/0609040](#)] [[SPIRES](#)].
- [28] LHCb collaboration, *Measurement of Δm_s in the decay $B_s^0 \rightarrow D_s^-(K^+K^-\pi^-)(3)\pi$* , note LHCb-CONF-2011-005, April 2011.
- [29] J. Alwall et al., *MadGraph/MadEvent v4: The New Web Generation*, *JHEP* **09** (2007) 028 [[arXiv:0706.2334](#)] [[SPIRES](#)].
- [30] MadGraph v5.1.2.3, <http://madgraph.phys.ucl.ac.be/>.
- [31] J. Pumplin et al., *New generation of parton distributions with uncertainties from global QCD analysis*, *JHEP* **07** (2002) 012 [[hep-ph/0201195](#)] [[SPIRES](#)].
- [32] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 Physics and Manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)] [[SPIRES](#)].
- [33] CDF collaboration, F. Abe et al., *Search for new particles decaying to b anti- b in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, *Phys. Rev. Lett.* **82** (1999) 2038 [[hep-ex/9809022](#)] [[SPIRES](#)].
- [34] J.S. Conway, *Pretty Good Simulation of high-energy collisions*, 090401 release, <http://physics.ucdavis.edu/~conway/research/software/pgs/pgs4-general.htm>.
- [35] D0 collaboration, V.M. Abazov et al., *Search for neutral Higgs bosons in the multi- b -jet topology in 5.2fb-1 of $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV*, *Phys. Lett. B* **698** (2011) 97 [[arXiv:1011.1931](#)] [[SPIRES](#)].
- [36] K. Cheung and W.-Y. Keung, *Split supersymmetry, stable gluino and gluinonium*, *Phys. Rev. D* **71** (2005) 015015 [[hep-ph/0408335](#)] [[SPIRES](#)].
- [37] Y. Kats and M.D. Schwartz, *Annihilation decays of bound states at the LHC*, *JHEP* **04** (2010) 016 [[arXiv:0912.0526](#)] [[SPIRES](#)].
- [38] C. Kim and T. Mehen, *Color Octet Scalar Bound States at the LHC*, *Phys. Rev. D* **79** (2009) 035011 [[arXiv:0812.0307](#)] [[SPIRES](#)].
- [39] A.D. Martin, W.J. Stirling, R.S. Thorne and G. Watt, *Parton distributions for the LHC*, *Eur. Phys. J. C* **63** (2009) 189 [[arXiv:0901.0002](#)] [[SPIRES](#)].
- [40] A. Djouadi, *The Anatomy of electro-weak symmetry breaking. II. The Higgs bosons in the minimal supersymmetric model*, *Phys. Rept.* **459** (2008) 1 [[hep-ph/0503173](#)] [[SPIRES](#)].
- [41] M.S. Carena, S. Heinemeyer, C.E.M. Wagner and G. Weiglein, *MSSM Higgs boson searches at the Tevatron and the LHC: Impact of different benchmark scenarios*, *Eur. Phys. J. C* **45** (2006) 797 [[hep-ph/0511023](#)] [[SPIRES](#)].
- [42] W.A. Bardeen and C.T. Hill, *Chiral dynamics and heavy quark symmetry in a solvable toy field theoretic model*, *Phys. Rev. D* **49** (1994) 409 [[hep-ph/9304265](#)] [[SPIRES](#)].
- [43] W.A. Bardeen, E.J. Eichten and C.T. Hill, *Chiral Multiplets of Heavy-Light Mesons*, *Phys. Rev. D* **68** (2003) 054024 [[hep-ph/0305049](#)] [[SPIRES](#)].
- [44] B.A. Dobrescu, K. Kong and R. Mahbubani, *Prospects for top-prime quark discovery at the Tevatron*, *JHEP* **06** (2009) 001 [[arXiv:0902.0792](#)] [[SPIRES](#)].
- [45] CDF collaboration, T. Aaltonen et al., *Search for new particles decaying into dijets in proton- antiproton collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. D* **79** (2009) 112002 [[arXiv:0812.4036](#)] [[SPIRES](#)].
- [46] <http://www-cdf.fnal.gov/physics/ewk/2010/WW.WZ/index.html>.

- [47] E. Eichten and K.D. Lane, *Low - scale technicolor at the Tevatron*,
Phys. Lett. B **388** (1996) 803 [[hep-ph/9607213](#)] [[SPIRES](#)].
- [48] E. Eichten and K. Lane, *Low-scale technicolor at the Tevatron and LHC*,
Phys. Lett. B **669** (2008) 235 [[arXiv:0706.2339](#)] [[SPIRES](#)].
- [49] K. Lane and A. Martin, *An Effective Lagrangian for Low-Scale Technicolor*,
Phys. Rev. D **80** (2009) 115001 [[arXiv:0907.3737](#)] [[SPIRES](#)].