On flux quantization in F-theory II: unitary and symplectic gauge groups

Andrés Collinucci^{a,b} and Raffaele Savelli^c

^a*Theory Group, Physics Department, CERN,
CH-1211 Geneva 23, Switzerland*

^b*Physique Théorique et Mathématique Université Libre de Bruxelles,
C.P. 231, 1050 Bruxelles, Belgium*

^c*Max-Planck-Institut für Physik,
Föhringer Ring 6, 80805 Munich, Germany*

E-mail: collinucci.phys@gmail.com, raffaele.savelli@gmail.com

ABSTRACT: We study the quantization of the M-theory G-flux on elliptically fibered Calabi-Yau fourfolds with singularities giving rise to unitary and symplectic gauge groups. We seek and find its relation to the Freed-Witten quantization of worldvolume fluxes on 7-branes in type IIB orientifold compactifications on Calabi-Yau threefolds. By explicitly constructing the appropriate four-cycles on which to calculate the periods of the second Chern class of the fourfolds, we find that there is a half-integral shift in the quantization of G-flux whenever the corresponding dual 7-brane is wrapped on a non-spin submanifold. This correspondence of quantizations holds for all unitary and symplectic gauge groups, except for SU(3), which behaves mysteriously.

We also perform our analysis in the case where, in addition to the aforementioned gauge groups, there is also a ‘flavor’ U(1)-gauge group.

KEYWORDS: F-Theory, D-branes, M-Theory

ARXIV EPRINT: [1203.4542](https://arxiv.org/abs/1203.4542)

Contents

1	Introduction	1
2	Summary of results	3
3	Detection of the Freed-Witten anomaly in F-theory: Sp-series	4
3.1	Sen limit	4
3.2	F-theory lift	6
3.2.1	General idea	6
3.2.2	Formalization	10
4	Detection of the Freed-Witten anomaly in F-theory: SU-series	14
4.1	Sen limit	14
4.2	F-theory lift	15
4.2.1	The $SU(2N)$ family	15
4.2.2	The $SU(2N+1)$ family	17
4.3	The example of $SU(5)$	20
5	U(1)-restrictions	22
6	Outlook	23
A	Geometry of $SU(N)$ F-theory configurations	24
B	The $SU(3)$ case	25

1 Introduction

The investigation of global aspects of F-theory compactifications has received much attention in the past few years [1–9]. Although primarily driven by phenomenological aspirations, these studies have furthered our understanding of fundamental aspects in general F-theory configurations. One such notable aspect concerns G-flux [10–15]. A detailed knowledge of these objects and of their properties is crucial for both model building issues, like generation of chiral matter in the gauge theory sector and moduli stabilization in the gravitational sector, and more fundamental questions regarding the quantum consistency of the UV theory. An essential tool which has driven the recent development of the field is the duality between F and M-theory [16], which allows for a rigorous and elegant definition of F-theory through its long-wavelength limit (eleven-dimensional supergravity) (for reviews see [17, 18]).

However, many questions remain unanswered. In this paper we will address the quantization rule for the M-theory G_4 flux [19] in its relation to the Freed-Witten quantization condition for gauge fluxes on 7-brane configurations [20, 21]. In [22], the authors studied the link between the half-integral shifts in the quantization condition of G_4 and in the quantization condition of the dual 7-brane gauge flux. While proving that F-theory configurations with smooth Weierstrass representation never display such a shift, they were able to indirectly analyze the singular cases. Indeed, they matched the quantization conditions of G_4 and of its dual 7-brane flux by using the type IIB weak coupling limit of F-theory, and comparing the M2 and D3 induced charges. However, the analysis focused on symplectic gauge groups, which are more tractable at weak coupling.

In this paper, we generalize our previous studies, and render them physically more intuitive. We will in fact find a purely M-theoretic way of detecting the membrane anomaly. Moreover we will show that our method provides a direct lift to M-theory of the usual strategy adopted to detect Freed-Witten anomaly in perturbative D-brane physics. More precisely, a D7-brane wrapping a non-spin manifold carries a half-integrally quantized flux. Given the 2-cycle on which the period of the flux is half-integral, we identify the corresponding 4-cycle in M-theory where the period of the second Chern class of the elliptic fourfold is an odd integer, thus modifying the quantization rule of G_4 .

The authors of [23] have shown that no shift in the quantization of G_4 can be detected by integrating c_2 on holomorphic 4-cycles given by complete intersections of divisors, or on fibrations of exceptional $\mathbb{P}^1$'s over matter curves. Our strategy will be to look at holomorphic 4-cycles that are not complete intersections of divisors with the Calabi-Yau hypersurface, much in the same spirit of [14]. Their explicit description in terms of vanishing loci of algebraic equations is not present in a general point of the moduli space of the fourfold.

By mildly constraining the complex structure of the Weierstrass model, we are able to find those 4-cycles, which are generically not holomorphic, and therefore geometrically not tractable. In fact, we will propose two alternative methods to perform such a constraint, based on two different Ansätze on the coefficients of the ‘Tate model’. For both strategies, we will provide a clear physical interpretation in type IIA string theory. One of them allows us to make a more direct treatment of the symplectic series, hence extending and clarifying the results of [22], while the other one gives us the possibility to extend the same analysis to the whole tower of unitary gauge groups. This is indeed a very important generalization in light of the promising Grand-Unified-Theory models which are mainly based on the existence of an $SU(5)$ gauge stack of 7-branes.

A notable exception in our investigation is represented by the $SU(3)$ configuration, for which neither prescription works. We will present the details of this model in an appendix and explain the reasons of this anomalous behavior. Finally we will extend our analysis to models with flavor $U(1)$ -gauge groups, whose relevance in the generation of chiral matter on D-brane intersections is well-known [24].

The paper is organized as follows: After summarizing our results in section 2, we address the Sp -series, finding appropriate four-cycles on which to detect the half-integral shift in the quantization of G_4 . We relate these directly to the two-cycles on which the Freed-Witten shift of the corresponding D7 gauge flux is detected.

In section 4, we proceed with the case of unitary groups, which are generically less tractable from the type IIB point of view. Nevertheless, we find that the quantization of G_4 still mimics that of gauge flux of the corresponding D7-stack. To make our formulae more palatable, we work out explicitly the interesting case of $SU(5)$.

In section 5 we take our investigation one step further by adding singularities that give rise to ‘flavor’ $U(1)$ gauge groups. We conclude with an outlook in section 6.

In appendix A we give a general description of the resolution of CY fourfolds with arbitrary $SU(N)$ singularities over arbitrary base threefolds.

In appendix B we say a few words about the special case of $SU(3)$ gauge group.

2 Summary of results

Let us collect here the main results of this paper for the convenience of the reader.

We have found non-complete intersection, integral 4-cycles of the F-theory fourfold on which we are able to measure odd values of the second Chern class and hence half-integral shifts in the quantization condition of G_4 . We have performed this analysis for both the symplectic and the unitary series of gauge groups. We find half-integral shifts precisely when the corresponding 7-brane stack is wrapping a non-spin manifold, thus directly connecting this topological effect with the Freed-Witten quantization condition in type II string theory.

The general setup consists of a 7-brane stack with $Sp(N)$ gauge group wrapped on a divisor $\mathcal{D}$, and a ‘remainder’ 7-brane that saturates the $C_{(0)}^{RR}$ -tadpole (shaped like a *Whitney umbrella*, as shown in [25]). Given a holomorphic curve $\mathcal{C} \subset \mathcal{D}$, we explicitly construct a holomorphic surface $C_{(4)}$ in the resolved F-theory CY fourfold $\tilde{Z}_4$, such that

$$\int_{C_{(4)}} c_2(\tilde{Z}_4) = \int_{\mathcal{C}} [7 c_1(B_3) - (2N - 1) \mathcal{D}] . \quad (2.1)$$

This 4-cycle has the geometry of a $\mathbb{P}^1$ fibered over the curve $\mathcal{C} \subset \mathcal{D}$ and it lifts a loop of type IIA open strings stretching between one D-brane of the stack and the Whitney umbrella D-brane. The latter is always spin in the $Sp(N)$ -case. Hence, if $\mathcal{D}$ is not spin, such that its first Chern class has an odd period on $\mathcal{C}$, then $c_2(\tilde{Z}_4)$ will have an odd period on our 4-cycle $C_{(4)}$.

For the $SU(2N)$ -series with $N \geq 2$ we construct the 4-cycle in the same way. The general formula for the integrated second Chern class is now

$$\int_{C_{(4)}} c_2(\tilde{Z}_4) = \int_{\mathcal{C}} (6 c_1(B_3) - (2N - 1) \cdot \mathcal{D}) . \quad (2.2)$$

For the $SU(2N + 1)$ -series with $N \geq 2$ we construct a different type of 4-cycle $C'_{(4)}$, which however also consists in fibering a $\mathbb{P}^1$ over $\mathcal{C}$. Physically, this other 4-cycle is lifting a loop of closed, non-orientable type IIA strings intersecting one D-brane of the stack on $\mathcal{C}$. The general result for the integrated second Chern class is

$$\int_{C'_{(4)}} c_2(\tilde{Z}_4) = \int_{\mathcal{C}} \mathcal{D} . \quad (2.3)$$

It is worth remarking that this second strategy works fine also for $SU(2N)$ with $N \geq 3$.

We finally have investigated flux quantization in F-theory models with the so-called $U(1)$ -restriction [24]. For gauge groups $SU(N \geq 5)$ nothing changes with respect to the previous discussion provided we treat all of these cases with the second type of 4-cycle, $C'_{(4)}$.

For the Sp -series we conjecture that odd-rank Sp groups lead to even second Chern classes and viceversa. This is because in the odd-rank cases the two branches of the ‘flavour brane’ are separately non-spin and the induced flux on them is also half-integral. This effect cancels the analog effect arising on the non-abelian stack. In contrast, in the even-rank cases, the branches of the ‘flavour brane’ are spin and the induced gauge flux is integrally quantized. For this reason we still find, by explicit computation, an odd second Chern class also for the $U(1)$ -restricted $SU(4)$ -model.

3 Detection of the Freed-Witten anomaly in F-theory: Sp -series

In [22] it has been argued that, for M-theory on elliptically fibered Calabi-Yau fourfolds with a codimension one singularity of Kodaira type I_{2N}^{ns} , the G_4 flux is quantized in terms of half-integers if and only if the singular locus does not admit spin structures. This is the F-theory counterpart of the Freed-Witten quantization condition for the gauge flux on the D7-brane $Sp(N)$ -stack¹ which wraps a non-spin submanifold of the Calabi-Yau threefold in the weakly coupled type IIB string picture. However, it has not been clarified yet on which 4-cycles of the fourfold one is actually able to detect the shifted quantization condition. In fact, it turns out that such cycles are not manifest as algebraic submanifolds in a generic point of the moduli space of the fourfold. In the same spirit as in [14], in this section we will identify them by mildly constraining the complex structure moduli of the fourfold. As we will see, the way to perform the restriction is suggested by the description of the system in the weak coupling limit, which for this reason we will present first.

Throughout this paper, we will use the phrase *Freed-Witten anomaly*, to refer to the half-integral shift in the quantization of worldvolume flux, as opposed to the anomaly related to H_3 -fluxes that restrict non-trivially to the brane.

3.1 Sen limit

As reviewed in [22], Sen’s weak coupling limit of an F-theory configuration is described in terms of a Calabi-Yau threefold equipped with an orientifold involution. If B_3 is the base manifold of the F-theory elliptic fourfold, the type IIB Calabi-Yau threefold is the double cover of B_3 defined by the following equation

$$X_3 : (\xi - a_1)(\xi + a_1) = 4a_2. \quad (3.1)$$

in an ambient four-dimensional manifold. The ambient variety is obtained by adding to B_3 the homogeneous coordinate ξ of degree $c_1(B_3)$. The polynomials $a_{1,2}$, which depend on the coordinates of B_3 , are sections of $K_{B_3}^{-1}$, $K_{B_3}^{-2}$ respectively (they are the first Tate coefficients of the elliptic fourfold defining equation, see eq. (3.9) below). The geometrical

¹We will always use for symplectic groups the notation where N refers to the rank.

action of the orientifold involution is defined by the operation $\xi \rightarrow -\xi$ and the equation $\xi = 0$ determines the locus wrapped by an $O7^-$ -plane.

If the F-theory fourfold has an $\mathrm{Sp}(N)$ singularity over the locus $\{D = 0\} \subset B_3$ with D an homogeneous polynomial, in the weak coupling limit there is a stack of $2N$ orientifold-invariant D7-branes wrapping the manifold given by the complete intersection

$$S_2 : \begin{cases} D = 0 \\ (\xi - a_1)(\xi + a_1) = 4a_2 \end{cases} . \quad (3.2)$$

Being transverse to the $O7$ -plane, the stack accommodates an $\mathrm{Sp}(N)$ gauge theory. Consider the case in which the submanifold (3.2) is non-spin. Then, there must exist 2-cycles of the D7-stack on which $c_1(S_2)$ has an odd period. However, these may have no algebraic representatives in a generic region of the moduli space of the D7-divisor. That is indeed the case here, and the mildest assumption we can make in order that those Riemann surfaces show up is as follows. Let us constrain the polynomials D and a_2 to be of the form

$$\begin{aligned} D &\equiv P \hat{D} + Q \tilde{D} \\ a_2 &\equiv P \hat{a}_2 + Q \tilde{a}_2 , \end{aligned} \quad (3.3)$$

while keeping a_1 completely generic. Here P, Q are generic polynomials of *odd* degree² on X_3 . If $\hat{D}, \tilde{D}$ were constants, we could redefine things such that for example D simply coincides with P , but in general this is not the case. It is now easy to see that thanks to (3.3) some integral (1,1)-cycles of the D7-stack are manifest in our Calabi-Yau threefold and they can be written in the ambient four-dimensional manifold as

$$C_{(2)}^\pm : \begin{cases} P = 0 \\ Q = 0 \\ \xi = \pm a_1 \end{cases} . \quad (3.4)$$

As it is also the case in [14], the subvarieties (3.4) are complete intersections neither with the Calabi-Yau threefold equation (3.1), nor with the divisor $\{D = 0\}$ representing the D7-stack. Rather, they are codimension three subspaces of the ambient, which are automatically contained in S_2 . Moreover, one is the image of the other under the orientifold map. By construction, $c_1(S_2)$ may now integrate to an odd number over $C_{(2)}^\pm$. In contrast, every 2-cycle which is complete intersection with X_3 gives certainly an even result for the integral of $c_1(S_2)$, since X_3 is an hypersurface of even class ($2c_1(B_3)$) in the ambient manifold. Notice, finally, that the constraints (3.3) do not introduce any singularity in X_3 , as can be verified by computing the gradient of eq. (3.1) subjected to the ansatz (3.3).

As an example, let us consider the toy model discussed in ref. [22], namely $B_3 = \mathbb{P}^3$ with homogeneous coordinates $x_1, \dots, x_4$ and hyperplane class H . The associated double-cover type IIB Calabi-Yau threefold is the octic hypersurface $W\mathbb{P}_{1,1,1,1,4}^4$ [8], where ξ has degree four, and the hypersurface equation is given by

$$x_1^8 - x_2^8 + x_3^8 - x_4^8 - \xi^2 = 0 . \quad (3.5)$$

²More precisely, we require the Poincaré-dual classes of the submanifolds $\{P = 0\} \subset X_3$ and $\{Q = 0\} \subset X_3$ be odd classes.

Let the D7-brane be located at $D = \sum_{i=1}^4 a_i x_i$. If we wanted to determine, whether or not this brane is spin, we would have to integrate its first Chern class $c_1(D) = H$ on every possible 2-cycle, and check for even/odd results. Any 2-cycle defined as a complete intersection of D with another divisor will give an even answer, due to the intersection number $H^3 = 2$ of this CY threefold. Yet, this brane is *not* spin.

In order to detect this, one must search for holomorphic curves that cannot even be written as complete intersections of two divisors with X_3 , but as complete intersections of three divisors in the ambient fourfold. Let us take for simplicity the curve C

$$C: \quad \xi = x_4^4 \quad \cap \quad x_1 = x_2 \quad \cap \quad x_3 = 0. \quad (3.6)$$

This curve is a $\mathbb{P}^1$ that is automatically contained in X_3 . If we now restrict the D7-brane moduli to take the form:

$$D = a_1(x_1 - x_2) + a_3 x_3, \quad (3.7)$$

then it will pass through C , as opposed to just intersecting it. Now we may integrate $c_1(D)$ over this curve, and we will find that

$$\int_C c_1(D) = 1. \quad (3.8)$$

Although we have ‘forced’ this curve upon the D7-brane by constraining the moduli, it is always present in the divisor’s homology, just not holomorphically.

3.2 F-theory lift

3.2.1 General idea

The discussion above suggests the form of the 4-cycles on which we could detect the oddness of the second Chern class of the Calabi-Yau fourfold in an $\mathrm{Sp}(N)$ -singular F-theory configuration. We define the elliptically fibered Calabi-Yau fourfold with an I_{2N}^{ns} -singularity on $\{D = 0\} \subset B_3$ as the hypersurface

$$Z_4^{\mathrm{Sp}(N)} : Y^2 + a_1 XYZ + a_{3,N} D^N Y Z^3 = X^3 + a_2 X^2 Z^2 + a_{4,N} D^N X Z^4 + a_{6,2N} D^{2N} Z^6, \quad (3.9)$$

in the ambient fivefold given by a $W\mathbb{P}_{XYZ}^2$ -bundle over B_3 . Here $a_{i,j}$ is a section of the line bundle $K_{B_3}^{-i} \otimes \mathcal{L}_D^{-j}$, where $\mathcal{L}_D$ indicates the line bundle defined by the vanishing locus of its section D .

In order to compute any topological quantity, we have to completely resolve the Calabi-Yau fourfold. The resolution procedure, as prescribed by [26], leads via a series of blow-ups to a resolved fiber over the singular locus which exactly reproduces the extended (or affine) Dynkin diagram of the gauge group corresponding to the singularity one starts with. For non-split singularities, like the ones in the Sp -series, one has to take into account non-trivial monodromies acting on the various components of the resolved fiber as we go around non-contractible paths of the singular locus. The easiest such situation, with only two exceptional divisors, is depicted in figure 1, where E_1 and E_2 are the two components of the first exceptional divisor and they get exchanged by monodromy, while the extended

Figure 1. This shows the extended Dynkin diagram of $Sp(2)$ (right) obtained by folding the extended Dynkin diagram of $SU(4)$ (left). The monodromy responsible for the folding acts as a reflection with respect to the vertical dashed line drawn on the left figure.

node E_{-1} and the last exceptional divisor E_3 are left invariant. The green lines in figure 1 represent D6-branes in the type IIA picture. The resolved regime corresponds indeed to the Coulomb branch of the gauge theory living on the worldvolume of D6-branes. The latter are separated along the so called T-duality circle, i.e. one of the two fiber directions of the elliptic fibration, indicated in the figure by solid black lines. From this perspective, one realizes that the nodes of the diagram are nothing but *closed* M2 branes lifting the type IIA *open* strings stretched between two neighboring D6-branes. The additional direction is the so called M-theory circle, i.e. the other fiber direction of the elliptic fibration, which collapses on the D6-brane locations. Hence, in the non-split case, D6-branes as well undergo monodromies, in a way analogous to an orientifold involution. In fact, they are pairwise exchanged so to form couples of brane-image-brane (1 with 2 and 3 with 4 in the figure). Therefore, the left part of the figure displays the extended diagram of $SU(4)$ with the usual bunch of four (separated) D6-branes. If we quotient this configuration by the monodromy action, we obtain precisely the extended diagram of $Sp(2)$, with two (separated) stacks of brane-image-brane, depicted in the right figure as shadowed green lines.

The type IIA picture outlined above helps us finding in F-theory the 4-cycles which, lifting the 2-cycles (3.4), can be used to detect the half-quantization of the G_4 flux and thus the M-theory lift of the Freed-Witten anomaly. The general reasoning goes as follows.

The way we typically detect Freed-Witten anomalies [21] of D-branes wrapping holomorphic submanifolds is by considering a loop of open strings attached to a given D-brane and by integrating the first Chern class of the brane worldvolume over the 2-cycle generated by the loop of string boundaries. If the result is odd, we must switch on a half-quantized gauge flux on the D-brane to cancel the anomaly. However, if such a 2-cycle is taken to be holomorphic it must degenerate to a point which is the calibrated representative of the trivial homology class of the target space. Therefore, in order to measure Freed-Witten anomalies in this case, we are forced to consider two different D-branes, like the parallel D6s in figure 1, and look at loops of open strings stretched between them. These loops

will then form 3-chains with cylindrical shape having as boundaries holomorphic 2-cycles on each of the two D-branes. Analogously to the nodes of the Dynkin diagram discussed above, these 3-chains lift in M-theory to 4-cycles, regarded as loops of the closed M2-branes lifting the type IIA open strings considered.

Therefore, we are led to analyze the 4-cycles made by nodes of the Dynkin diagram fibered over the curve of the brane which we used to measure the Freed-Witten anomaly in type II string theory (see section 3.1). However, not every node works well to this end. In fact, any of the nodes depicted in the left part of figure 1 gives rise to a 4-cycle on which the second Chern class of the fourfold integrates to an even number. One can explicitly verify this statement using the details of the geometry discussed in the next subsection. But it is instructive to realize this from the type IIA point of view. Indeed, for instance, the integration of $c_2/2$ on the E_1 -fibration over the curve $\{P = Q = 0\} \subset B_3$ is equivalent modulo integers to the type IIA expression

$$\int_{\{P=Q=0\} \subset D6_1} F_1 - \int_{\{P=Q=0\} \subset D6_3} F_3, \quad (3.10)$$

where $F_{1,3}$ are the gauge fluxes on the $D6_{1,3}$ -branes respectively and the minus sign is due to the opposite orientations of the boundaries. If we take the D6-worldvolume to be non-spin, *both* F_1 and F_3 are half-integrally quantized to cancel the Freed-Witten anomaly. This means that the expression (3.10) is integral and thus the second Chern class of the fourfold integrates to an even number on the E_1 -fibration. The same conclusion holds for the E_2 -fibration (by considering the $D6_{2,4}$ -branes) and in general if the fiber is any Cartan node or the extended node. To be more precise, in the cases of the last Cartan node and of the extended node, the corresponding type IIA strings stretch between one D6-brane and its image $\widetilde{D6}$ under the monodromy. Hence the integration of $c_2/2$ for instance on the E_3 -fibration in the $Sp(2)$ example would be equivalent to the integer number

$$\int_{\{P=Q=0\} \subset D6_3} F_3 + \int_{\{P=Q=0\} \subset \widetilde{D6}_3} \tilde{F}_3, \quad (3.11)$$

where the plus sign arises because the gauge flux is odd under the orientifold involution.

The above logic suggests that a node which would allow us to measure an odd value for c_2 could be the one which lifts an open string stretching between a D-brane of the stack and an other D-brane which carries no flux (or an integrally quantized flux). The only such D-brane around is the Whitney-type brane. This leads us to constrain the complex structure of the fourfold in such a way that our curve $\{P = Q = 0\} \subset B_3$ is contained in both the D6-stack and the Whitney-umbrella brane. In other words, in order to identify the “detecting” 4-cycle, we have to require that the Freed-Witten 2-cycle $\{P = Q = 0\} \subset B_3$ becomes a branch of the matter curve represented by the intersection between the non-abelian stack and the I_1 -locus. Before formalizing this concept, let us explain the result in terms of diagrams. Starting from an $Sp(N)$ -singularity, our constraint on the complex structure of the blown-up fourfold will be such that there will be an enhancement to $SU(2N+1)$ along the curve $\{P = Q = 0\}$, as pictorially shown in figure 2. As it is manifest from the figure, the only relevant change for our purpose is the splitting of the last Cartan node (E_3 for

Figure 2. This shows the transition from the extended Dynkin diagram of $\text{Sp}(2)$ (left) to the extended Dynkin diagram of $\text{SU}(5)$ (right) happening along the curve $\{P = Q = 0\}$ due to the singularity enhancement. The fifth D-brane of the $\text{SU}(5)$ stack is given by the Whitney-type brane. The orange nodes are the fibers of the 4-cycles on which it is possible to detect the Freed-Witten anomaly.

$\text{Sp}(2)$) into two branches ($E_3^{(1)}$ and $E_3^{(2)}$), each of which lifts the open string stretching between one D-brane of the stack and the Whitney-type D-brane W . Due to the constraint made, the shape of the Whitney-umbrella will not be completely generic and an amount of integral flux would be necessarily generated on it to keep this configuration stable. However, since we are not focusing on the dynamics of the system here, but rather we are probing it to measure possible anomalies, we can assume that no flux is present on W . Therefore, focusing on the example of figure 2, the integration of $c_2/2$ on the $E_3^{(1)}$ -fibration over the curve $\{P = Q = 0\} \subset B_3$ is equivalent modulo integers to the type IIA expression

$$\int_{\{P=Q=0\} \subset \text{D6}_3} F_3, \quad (3.12)$$

which is half-integral if and only if the stack is non-spin. For the $E_3^{(2)}$ -fibration, we find the same expression in terms of the image flux $\tilde{F}_3$.

Let us finally remark that the symmetry enhancement along the whole matter curve does not have to be $\text{SU}(2N+1)$, which only appears along the branch $\{P = Q = 0\}$ due to the ansatz we will make. Indeed, for an I_{2N}^{ns} -Kodaira singularity, Tate's algorithm prescribes along the intersection with the I_1 -locus an enhancement to an I_{2N+1}^{ns} -Kodaira singularity, which corresponds to an unconventional gauge group [26]. As it will be clearer later, the reason for this slightly stronger requirement is to have a closed algebraic expression for the two branches of the last Cartan node. In contrast, by just asking

$$W = \hat{W}P + \tilde{W}Q, \quad (3.13)$$

where $\{W = 0\}$ defines the Whitney-type D-brane W , we would still split the last Cartan node, but non-trivial monodromies would mix the two branches and we could not single out one of them globally [14].

3.2.2 Formalization

Let us now put onto rigorous ground what we have previously discussed. In order to blow-up the $\mathrm{Sp}(N)$ -singularity of (3.9), we will follow the toric procedure adopted in [22]. First of all we introduce a new homogeneous coordinate σ to the ambient space and impose the additional equation $\sigma = D$. Then we add N toric vectors v_i (together with N projective relations), which, in order to stick with the notations of [26], we will label with odd numbers only,³ i.e. $1, \dots, 2N - 1$.

The Calabi-Yau fourfold will be defined as a complete intersection of two equations in an ambient sixfold X_6 , given by a projective bundle over B_3 . The ambient fiber over B_3 will be defined by the following assignment of projective weights

$$\begin{array}{c}
 W\mathbb{P}_{\sigma XYZ v_1 v_3 \dots v_{2N-3} v_{2N-1}}^3 : \begin{array}{ccccccccc}
 \sigma & X & Y & Z & v_1 & v_3 & \cdots & v_{2N-3} & v_{2N-1} \\
 0 & 2 & 3 & 1 & 0 & 0 & \cdots & 0 & 0 \\
 1 & 1 & 1 & 0 & -1 & 0 & \cdots & 0 & 0 \\
 1 & 2 & 2 & 0 & 0 & -1 & \cdots & 0 & 0 \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\
 1 & N-1 & N-1 & 0 & 0 & 0 & \cdots & -1 & 0 \\
 1 & N & N & 0 & 0 & 0 & \cdots & 0 & -1
 \end{array}
 \end{array} \quad (3.14)$$

The rows represent independent $\mathbb{C}^*$ -actions on the coordinates. In order to define the ambient space, we mod these out, much like in projective spaces. The first row indicates the pre-existing projective identification of $W\mathbb{P}_{XYZ}^2$.

Projective spaces are defined as $\mathbb{C}^n - \{0, \dots, 0\}/\mathbb{C}^*$, i.e. before modding out by the action, we must first delete the origin. In toric language, one would say that the *Stanley-Reisner ideal* consists of the element $z_1 \dots z_n$, meaning that these coordinates are forbidden from vanishing simultaneously.

The fiber defined above for the sixfold X_6 has the following SR ideal:

$$SR_{\mathrm{Sp}(N)} : \left\{ XYZ ; v_{2i-1}Z|_{i=1, \dots, N} ; v_{2i-1}XY|_{i=0, \dots, N-1} ; v_{2i-1}v_{2j-1} \left| \begin{array}{l} i, j=0, \dots, N \\ j-i > 1 \end{array} \right. \right\}, \quad (3.15)$$

where by convention we put $v_{-1} := \sigma$. The above is easily derived if we express the table (3.14) in the basis in which it is manifest that v_{2i-1} induces a blow-up along the locus $X = Y = v_{2i-3} = 0$, for $i = 1, \dots, N$. It is not difficult to realize that the last elements in (3.15), quadratic in the vs , arise by taking pairwise differences of the rows in table (3.14) expressed in the new basis. These elements are responsible for the appearance of the “unidimensional” structure of the extended Dynkin diagrams of the **C**-series. Calling $V \equiv \prod_{i=0}^N v_{2i-1}^{N-i}$, which has multi-degree $(0, 1, 2, \dots, N-1, N)$, the blown-up Calabi-Yau

³The reason is to distinguish them from the even-labeled ones which are further introduced to resolve split singularities, like the ones leading to unitary gauge groups (see section 4.2).

fourfold is given by the following intersection in the sixfold X_6 :

$$\tilde{Z}_4^{\text{Sp}(N)} : \begin{cases} Y(Y + a_1 X Z + a_{3,N} V Z^3) = X^3 \prod_{i=0}^N v_{2i-1}^i + \\ \quad + a_2 X^2 Z^2 + a_{4,N} V X Z^4 + a_{6,2N} V^2 Z^6 & 2 \times (3, 1, 2, \dots, N-1, N) \\ \prod_{i=0}^N v_{2i-1} = D & (0, \dots, 0), \end{cases} \quad (3.16)$$

where on the right the multi-degrees of the two hypersurfaces are displayed. Note that, as was the case for the Calabi-Yau threefold in section 3.1, these degrees are even.

The second Chern class of the resolved fourfold, which determines the quantization condition of the G_4 flux [19], is conveniently split into two pieces

$$c_2(\tilde{Z}_4^{\text{Sp}(N)}) = c_2(Z_4) + \Delta c_2, \quad (3.17)$$

where the first part is the second Chern class of the smooth phase of the elliptically fibered fourfold

$$c_2(Z_4) = c_2(B_3) + 11c_1^2(B_3) - 12F^2, \quad (3.18)$$

while the second depends in an easy way on the N exceptional divisors arising after the resolution

$$\Delta c_2 = - \sum_{i=1}^N i (7c_1(B_3) - i\mathcal{D}) E_{2i-1}. \quad (3.19)$$

In the expressions above F indicates the class of the 0-section $Z = 0$, $\mathcal{D}$ the Poincaré-dual class of $\{D = 0\} \subset B_3$, $E_k : v_k = 0$ are the blow-up classes, and we have everywhere omitted the pull-back map acting on classes of the base. It has been proven in [22] that the class (3.18) is always even. Thus we concentrate on the class (3.19), which, if odd, leads to a “half”-quantized G_4 flux. Let us remark here that eq. (3.19) provides in this context an iteration of Fulton’s formula [27, 28] for resolved manifolds after a single blow-up, thus extending the result for $N = 1$ already discussed in [22].

Analogously to the case of the Calabi-Yau threefold (3.1), a generic equation like (3.16) for the Calabi-Yau fourfold does not allow us to identify the 4-cycles on which the integral of Δc_2 is odd. Therefore, we implement the general logic outlined in the previous subsection by the following ansatz for some of the polynomial coefficients of (3.16)

$$D \equiv P \hat{D} + Q \tilde{D} \quad (3.20)$$

$$a_{2k+2, kN} \equiv P \hat{a}_{2k+2, kN} + Q \tilde{a}_{2k+2, kN} \quad k = 0, 1, 2.$$

This extends conditions (3.3), which are recovered here by taking $k = 0$ only, and implies condition (3.13) since

$$W = a_{4,N}^2 + a_1 a_{3,N} a_{4,N} - a_1^2 a_{6,2N} - a_2 a_{3,N}^2 - 4a_2 a_{6,2N}. \quad (3.21)$$

By looking at the structure of the various exceptional divisors $\{v_{2i-1} = 0\}$ when restricted to the curve $\{P = Q = 0\}$, one easily realizes that their pattern of intersection undergoes exactly the transition displayed in figure 2. In particular, all of them split into two well-defined branches and all the pre-existing monodromies are killed. Indeed, on $\{P = Q = 0\}$ one has

$$E_{2i-1} : \left\{ \begin{array}{l} v_{2i-1} = 0 \\ Y = 0 \end{array} \right. \cup \left\{ \begin{array}{l} v_{2i-1} = 0 \\ Y = -a_1 \end{array} \right. \quad i = 1, \dots, N-1, \quad (3.22)$$

each of which geometrically is a $\mathbb{P}^1$ with coordinates v_{2i-3}, v_{2i+1} fibered over the curve. However, as already stressed, it is the last exceptional divisors which interests us more. Before the transition, E_{2N-1} is a quadratic $\mathbb{P}^1$ embedded in the $\mathbb{P}^2$ with coordinates X, Y, v_{2N-3} . But after imposing (3.20), our Calabi-Yau fourfold gains extra $(2, 2)$ -cycles, due to the splitting of E_{2N-1} , and they look like

$$C_{(4)}^{(1)} : \left\{ \begin{array}{l} P = 0 \\ Q = 0 \\ v_{2N-1} = 0 \\ Y = 0 \end{array} \right., \quad (3.23)$$

$$C_{(4)}^{(2)} : \left\{ \begin{array}{l} P = 0 \\ Q = 0 \\ v_{2N-1} = 0 \\ Y = -(a_1 X + a_{3,N} v_{2N-3}) \end{array} \right., \quad (3.24)$$

where in (3.24) we have performed the N gauge-fixings $Z = v_{2i-1} = 1$ for $i = 0, \dots, N-2$, so that $V = v_{2N-3}$. We are still left with one gauge freedom, which we can use to describe the geometry of this 4-cycles. A closer look shows indeed that they are $\mathbb{P}^1$ -fibrations over the curve $\{Q = 0\} \cap \{P = 0\}$ contained in the surface where the singularity was. The fibers are $E_{2N-1}^{(1)}$ and $E_{2N-1}^{(2)}$ respectively, in the notation of the previous subsection. The last equations of (3.23) and (3.24) tell us that X and v_{2N-3} are exactly the coordinates of the $\mathbb{P}^1$ fibers, due to the presence of the element $v_{2N-3}XY$ in the Stanley-Reisner ideal (3.15). Again these 4-cycles are not complete intersection with the Calabi-Yau fourfold. Rather they are surfaces of the ambient six-fold which are automatically contained in $\tilde{Z}_4^{\text{Sp}(N)}$, namely they automatically satisfy equations (3.16). While integrating Δc_2 over any 4-cycle which is complete intersection with (3.16) gives certainly an even number, due to the even-ness of the fourfold class, the new 4-cycles may in contrast be good candidates to detect the Freed-Witten anomaly of F-theory. In fact, we find the following general formula

$$\begin{aligned} \int_{C_{(4)}^{(1,2)}} \Delta c_2 &= \int_{B_3} [7 c_1(B_3) - (2N-1) \mathcal{D}] \mathcal{P} \mathcal{Q} \\ &= \int_{\{P=Q=0\} \subset S_2} [2(4-N) c_1(B_3)|_{S_2} + (2N-1) c_1(S_2)], \end{aligned} \quad (3.25)$$

where $\mathcal{P}, \mathcal{Q}$ are the Poincaré-dual class of $\{P = 0\}, \{Q = 0\} \subset B_3$ respectively, and in the last step we have used the adjunction formula. Thanks to the simple structure of Δc_2 in (3.19), which does not contain terms quadratic in the exceptional divisors, eq. (3.25) can

be easily derived. To do that, one first of all simplifies the integrand using the elements $v_{2i-1}XY|_{i=0,\dots,N-1}$ of the SR ideal (3.15). They imply on $\{Y = 0\}$ the following relation

$$E_{2N-1} \sum_{i=1}^N iE_{2i-1} = -D \left(2c_1(B_3) + 2F - \sum_{i=1}^N iE_{2i-1} \right) + 2E_{2N-1}c_1(B_3). \quad (3.26)$$

It is easy to see that now one can factorize the class of $\{Y = 0\} \cap \{D = 0\}$, which is half the class of the proper transform (3.16). Therefore one is led to compute an integral over the blown-up elliptic fourfold, where the following formulae hold for any type of singularity [29, 30]

$$\begin{aligned} \int_{\tilde{Z}_4} E_k \mathcal{A} \mathcal{B} \mathcal{C} &= 0 & k = 1, \dots, \text{rank } \mathbb{G}, \\ \int_{\tilde{Z}_4} E_k E_l \mathcal{A} \mathcal{B} &= -C_{kl} \int_{B_3} \mathcal{D} \mathcal{A} \mathcal{B} & k, l = 1, \dots, \text{rank } \mathbb{G}. \end{aligned} \quad (3.27)$$

Here $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are any three divisors of the base and C_{kl} is the *symmetric* Cartan matrix of the Lie algebra $\mathbb{G}$ corresponding to the singularity. For the $\text{Sp}(N)$ case the latter is the $N \times N$ matrix $C_{kl} = 2(\delta_{kl}[2 - \delta_{kN}] - \delta_{k+1,l} - \delta_{k,l+1})$, compatibly with (3.15).

The remarkable fact about formula (3.25), as the last line suggests, is that its even/oddness only depends on the singular locus S_2 being spin or not, while it is independent of whether the class $\mathcal{D}$ which defines it as a subvariety of B_3 is even or odd. In other words, it is only sensitive to an intrinsic property of that surface and not on the details of its embedding. More formally, the modulo two reduction of the expression in square brackets in (3.25) is $w_2(S_2)$, while it does not contain information about the spin properties of the normal bundle of S_2 in B_3 . On the one hand, this provides, as already argued, the mechanism for Freed-Witten anomaly cancellation of the corresponding D7-brane stack. On the other hand, it says that the degree of the polynomial defining S_2 as a divisor of B_3 plays no essential role in the quantization rule of G_4 . This is important because, a polynomial D of odd degree may well lead to a spin, and thus non-anomalous, D7-stack. For this to happen, B_3 itself must be non-spin.

It is not difficult to convince ourselves again the assumption (3.20) we have made does not generate singularities on the Calabi-Yau fourfold (3.16). By computing the gradient of eq. (3.16) subjected to the ansatz (3.20), one can indeed rigorously verify the absence of singularities.

Finally, it is important to note that our four-cycles, although constructed as $\mathbb{P}^1$ -fibrations over curves living at brane intersections, are never actually *entire* matter surfaces. The way we tune our moduli force the matter curves to become reducible, and our four-cycles are fibrations over one such component. In our case, $\{P = Q = 0\}$ is but one component of the whole matter curve. Indeed, the half-integral quantization of the gauge flux, as treated in [20], is *designed* to yield correct indices for bifundamental matter at brane intersections. It would therefore be contradictory to ever find a half-integral period of such a flux over a matter curve, but not over a piece of such a curve. This is consistent with the fact that [23] have not registered any shift in the G_4 quantization on matter surfaces.

4 Detection of the Freed-Witten anomaly in F-theory: SU-series

In this section we address the same problem for F-theory configurations with an I_M^s Kodaira singularity on a generic codimension one locus of the base. We will see that things are more subtle here both in the geometric F-theory picture and in the weakly coupled string description. We will again start with the Sen limit and afterwards study the geometrical structures of these split-type singularities by distinguishing even and odd ranks.

4.1 Sen limit

A crucial feature that distinguishes (and makes it more subtle) the weak coupling limit of F-theory configurations with unitary-type singularities from those with symplectic-type ones is that the formers lead to Calabi-Yau threefolds with conifold singularities. The origin of these singularities is in the Tate coefficient a_2 having for I_M^s a single zero along the singular locus $\{D = 0\}$, instead of being generic as is the case in eq. (3.1). This shape of a_2 is actually already contained in our constrained form (3.3). In other words, to treat Sen's limit of $SU(M)$ F-theory models, we simply have to further require in the moduli space the conditions

$$\begin{aligned}\hat{a}_2 &\equiv \hat{D} a_{2,1} \\ \tilde{a}_2 &\equiv \tilde{D} a_{2,1} .\end{aligned}\tag{4.1}$$

It is easy to see that this restriction is now sufficiently drastic to generate codimension three singularities in the Calabi-Yau threefold given by the following intersection of four divisors in the ambient four-manifold

$$\text{conifold points} : \begin{cases} \xi = 0 \\ a_1 = 0 \\ D = 0 \\ \hat{a}_2 = 0 . \end{cases}\tag{4.2}$$

Indeed, the CY hypersurface equation assumes the shape of a conifold:

$$\xi^2 = a_1^2 + \hat{a}_2 D .\tag{4.3}$$

However this restriction is essential to make visible the $U(M)$ stack⁴ of D7-branes and its image-stack under the orientifold map, as was elucidated in [23]. Indeed, the surface S_2 in (3.2) now clearly factorizes in brane plus image-brane

$$S_2^\pm : \begin{cases} D = 0 \\ \xi = \pm a_1 . \end{cases}\tag{4.4}$$

In contrast, in the “constrained” non-split situation discussed in section 3.1, the stack was invariant and only its curve $\{P = Q = 0\}$ factorized.

⁴We neglect here the issue of the $U(1)$ related to the center of mass and its fate in the F-theory lift.

Therefore we are now able to immediately identify the 2-cycles for detection of Freed-Witten anomalies without any need of further requirements on the moduli space. Given any polynomial Q with the same properties as before, the 2-cycles look like

$$C_{(2)}^{\pm} : \begin{cases} D = 0 \\ Q = 0 \\ \xi = \pm a_1 . \end{cases} \quad (4.5)$$

Moreover, notice that these Riemann surfaces do not generically intersect the conifold singularities of X_3 , because this would require solving five equations in the ambient four-manifold. Hence it is likely that the presence of the conifold points does not affect the evaluation of the integral of $c_1(S_2)$ over $C_{(2)}^{\pm}$. Nevertheless, being the Calabi-Yau threefold singular, the computation of Chern classes itself may no longer be reliable. And on top of that, it is not clear whether in this circumstance $w_2(S_2)$ equals $w_2(N_{X_3} S_2)$, and which one matters for cancellation of Freed-Witten anomalies. A procedure is therefore needed to cure this singularity. To this end, one may immediately think to a couple of possibilities, namely the small resolution of the conifold points and the deformation. However [31], both are ruled out: The former because, in the absence of B -field, it would lead to a Calabi-Yau threefold not invariant under the orientifold involution, the latter because it would break the unitary gauge group to the parent symplectic one. A standard resolution of the conifold points is also not acceptable, because it would be non-crepant. We will not attempt here to solve this problem at weak coupling. Rather, we will focus on finding the 4-cycles in the well-behaved F-theory which are able to detect Freed-Witten anomalies and thus give the right lift of (4.5).

4.2 F-theory lift

Let us now discuss in detail the structure of the F-theory fourfold in the presence of an I_N^s Kodaira singularity and its complete resolution. Once again, our analysis aims at finding suitable integral $(2, 2)$ -cycles for detecting shifted quantization conditions for the G_4 flux. It turns out to be convenient to split the discussion in two sub-cases, namely $SU(\text{even})$ and $SU(\text{odd})$ gauge groups in eight dimensions. In appendix A we have collected general formulae holding for all $SU(M)$ singularities, which we will need throughout this section.

4.2.1 The $SU(2N)$ family

We start from the even case as it is easier to deal with. The reason is that the I_{2N}^s family of Kodaira singularities is very closely related to the I_{2N}^{ns} . By looking at the table of the Tate algorithm in [26], one indeed realizes that the only difference is the one already encountered for the Sen limit in section 4.1, i.e. a_2 is generic in the non-split case, while it has a single zero along $\{D = 0\}$ in the split case. This strongly suggests that we should treat I_{2N}^s singularities in a similar manner as we did for I_{2N}^{ns} in section 3.2.

Since $SU(2) = Sp(1)$ has already been discussed in the previous section, the present analysis concerns $SU(2N)$ gauge groups for $N \geq 2$. Recall, however, that the prescription given in [26] for resolving I_{2N}^s requires to add further toric coordinates v_{2i} for $i = 1, \dots, N -$

Figure 3. This shows the transition from the extended Dynkin diagram of SU(4) (left) to the extended Dynkin diagram of SU(5) (right) happening along the curve $\{P = Q = 0\}$ due to the singularity enhancement. The fifth D-brane of the SU(5) stack is given by the Whitney-type brane. The orange nodes are the fibers of the 4-cycles on which it is possible to detect the Freed-Witten anomaly.

1, with new projective $\mathbb{C}^*$ -actions ρ_{2i} such that σ, X, Y, v_{2i} have weights $(1, i, i + 1, -1)$ respectively under ρ_{2i} . We refer to appendix A for the details of the geometry.

One can now easily seek for the detecting 4-cycles in analogy with the analysis done for the $\text{Sp}(N)$ singularities in section 3.2, by imposing that:

$$D \equiv P \hat{D} + Q \tilde{D} \quad (4.6)$$

$$a_{2k+2, kN} \equiv P \hat{a}_{2k+2, kN} + Q \tilde{a}_{2k+2, kN} \quad k = 1, 2, \quad (4.7)$$

and $a_2 = a_{2,1} D$. The ansatz makes the gauge symmetry enhance on $\{P = Q = 0\}$ from $\text{SU}(2N)$ to $\text{SU}(2N + 1)$ along the whole matter curve. The enhancement manifest itself as the splitting into two of the node $E_{2N-1} \mapsto E_{2N-1}^{(1)} \cup E_{2N-1}^{(2)}$. Such transition is shown in figure 3 for the $N = 2$ case.

Focusing on the first component, we find a 4-cycle $C^{(1)}$ given by the fibration of $E_{2N-1}^{(1)}$ over the curve detecting the Freed-Witten anomaly in type IIB:

$$C_{(4)}^{(1)} : \begin{cases} P = 0 \\ Q = 0 \\ v_{2N-1} = 0 \\ Y = 0 \end{cases}, \quad (4.8)$$

One can prove that the general formula for the integral of $c_2(\tilde{Z}_4^{\text{SU}(M)})$, on the surface (4.8) is the following:

$$\int_{C_{(4)}^{(1)}} c_2 = \int_{B_3} (6 c_1(B_3) - (2N - 1) \cdot \mathcal{D}) \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (4.9)$$

This shows that the even or oddness of $c_2(\tilde{Z}_4^{\text{SU}(M)})$ is correlated with the spin-ness or non spin-ness of the brane stack on $\mathcal{D}$. Again, we emphasize that, as explained in the last paragraph of 3.2.2, this 4-cycle cannot be a matter surface. Those should always yield integral periods for G_4 .

We will outline the arguments behind the proof, which is rather tedious in its detail:

The general elements of the Stanley-Reisner ideal found in appendix A imply that, on the 4-cycle (4.8), the following coordinates can be taken to be different from zero, and ‘gauge-fixed’ to one: $\sigma, v_1, \dots, v_{2N-3}, Z$. This means that the respective divisor classes vanish on the 4-cycle. Only X, v_{2N-2} remain unfixed.

Since our goal is to relate an integral on the ambient sixfold X_6 of the fourfold $\tilde{Z}_4^{\text{SU}(M)}$ to an integral on B_3 , we must identify a form that is Poincaré dual to B_3 in X_6 . In this case, we may choose one of the following two equivalent six-forms:

$$[Y] \cdot [v_{2N-1}] \cdot [X] \quad \text{or} \quad [Y] \cdot [v_{2N-1}] \cdot [v_{2N-2}]. \quad (4.10)$$

One can see that this works, because after setting all of these coordinates to zero, those on B_3 remain unfixed. Let us pick the latter $PD_{B_3} \equiv [Y] \cdot [v_{2N-1}] \cdot [v_{2N-2}]$.

Let us calculate the total Chern class $c(\tilde{Z}_4^{\text{SU}(M)})$ with the adjunction formula:

$$c(\tilde{Z}_4^{\text{SU}(M)}) = \frac{c(B_3) \cdot (1 + [x]) \cdot (1 + [y]) \cdot (1 + [z]) \cdot (1 + [\sigma]) \cdot \prod_{i=1}^{2N-1} (1 + [v_i])}{((1 + \mathcal{D}) \cdot (1 + [x] + [y] + c_1(B_3)))}. \quad (4.11)$$

By restricting onto $C_{(4)}^{(1)}$, the SR ideal allows us to get rid of a lot of terms:

$$c(\tilde{Z}_4^{\text{SU}(M)})|_{C_{(4)}^{(1)}} = \frac{c(B_3) \cdot (1 + x) \cdot (1 + y) \cdot (1 + v_{2N-2}) \cdot (1 + v_{2N-1})}{((1 + \mathcal{D}) \cdot (1 + x + y + c_1(B_3)))}. \quad (4.12)$$

For convenience, we will denote a divisor class of a coordinate p simply as p , and drop the $[]$. We must now exploit all linear relations between divisor classes, in order to express $c_2(\tilde{Z}_4^{\text{SU}(M)})$ in terms of the classes that vanish, and of v_{2N-2} . The general relations, *after restricting onto $C_{(4)}^{(1)}$* , are the following:

$$[v_{2N-1}] = \mathcal{D} - [v_{2N-2}]; \quad [X] = 2c_1(B_3) - N\mathcal{D} + [v_{2N-2}]; \quad [Y] = 3c_1(B_3) - N\mathcal{D}. \quad (4.13)$$

Dropping terms with four indices along B_3 , we can then extract the second Chern class:

$$c_2(\tilde{Z}_4^{\text{SU}(M)})|_{C_{(4)}^{(1)}} = (4c_1(B_3) - 2\mathcal{D}) \cdot [v_{2N-2}] - [v_{2N-2}]^2. \quad (4.14)$$

Finally, from the relation in the SR ideal $[v_{2N-2}] \cdot [X] = 0$, we find that $[v_{2N-2}]^2 = (N\mathcal{D} - 2c_1(B_3)) \cdot [v_{2N-2}]$. From this, we arrive at our general formula (4.9).

4.2.2 The $\text{SU}(2N+1)$ family

We are left to address the odd series of unitary groups, which is more subtle. We will actually develop a method for detecting the shift in quantization of c_2 for *all* unitary groups with $\text{SU}(M \geq 5)$, also including the even ones. The case $\text{SU}(3)$ is special, and is remanded to appendix B

By looking at the Tate table [26], we see that in order to enforce an I_{2N+1}^s singularity on top of an existing I_{2N}^s two more polynomials have to vanish at linear order on $\{D = 0\}$, i.e. $a_{4,N}$ and $a_{6,2N}$. Therefore, the ansatz for *all* the a -coefficients in (3.20) becomes automatically satisfied.

Figure 4. This shows the transition from the extended Dynkin diagram of SU(5) (left) to the extended Dynkin diagram of SO(10) (right) happening along the curve $\{P = Q = 0\}$ due to the singularity enhancement. Nodes connected by arrows are identified. The orange nodes are the fibers of the 4-cycles on which it is possible to detect the Freed-Witten anomaly.

A type IIA argument totally analogous to the one given in section 3.2 tells us that we can no longer use the branches of the last exceptional divisor to identify the detecting 4-cycles. Indeed, they now lift open strings stretching between two ordinary branes of the $SU(2N+1)$ -stack, both of them carrying an half-quantized gauge flux in the non-spin case. Therefore we have to look elsewhere.

The only other matter curve available is the intersection of the non-abelian stack and the orientifold plane, i.e. $\{D = a_1 = 0\} \subset B_3$. The perturbative orientifold does not permit open strings to end on it, and therefore does not carry a flux. However, we could consider closed strings which loop around the O-plane and come back with the opposite orientation, sweeping *non-orientable worldsheets*. If we now consider a S^1 -family of such worldsheets, we obtain a 3-cycle, like for instance a loop of Klein bottles. We take such a 3-cycle to intersect a given D-brane on the 2-cycle we are using to detect the perturbative Freed-Witten anomaly. Finally, this 3-cycle will lift in M-theory to the 4-cycle we are looking for in this new situation.

This argument suggests that we should constrain the complex structure of the blown-up fourfold, which is given in eq. (A.4), in such a way that the curve $\{P = Q = 0\} \subset B_3$ is automatically contained in both the D-brane stack and the orientifold plane. Therefore, we impose the following conditions

$$\begin{aligned} D &\equiv P \hat{D} + Q \tilde{D} \\ a_1 &\equiv P \hat{a}_1 + Q \tilde{a}_1, \end{aligned} \tag{4.15}$$

since the polynomial defining the O7-plane is

$$\text{O7} : h = a_1^2 + 4a_{2,1}D. \tag{4.16}$$

Since the curve $\{P = Q = 0\}$ is a branch of the intersection between the non-abelian stack and the O-plane, we experience on it the gauge symmetry enhancement from $SU(2N+1)$ to $SO(4N+2)$. This transition is pictorially shown for the $N = 2$ case in figure 4 and

it works the same way for $SU(2N)$ too, which enhances to $SO(4N)$. Hence this provides an additional, alternative way of detecting Freed-Witten anomalies for $SU(2N)$ F-theory configurations. The only notable exceptions are $SU(2)$, which enhances to the III-Kodaira singularity and $SU(3)$, which enhances to the IV^s -Kodaira singularity [32]. Since $SU(2) = Sp(1)$, we have already described for it the rules for the quantization of the G_4 flux. In contrast, we have not been able to detect the Freed-Witten anomaly for $SU(3)$ F-theory configurations. In fact, we argue that for those configurations the second Chern class of the fourfold is *always* even and in appendix B we give an argument in favor of this conjecture. Nevertheless, it is not yet clear which is the mechanism for Freed-Witten anomaly cancellation in their weak coupling limit if the $SU(3)$ -stack is non-spin but no half-quantized gauge flux is induced. We hope to come back to this issue in the near future.

The pattern of symmetry enhancement of $SU(M)$ ($M \geq 4$) to orthogonal symmetries on the O-plane can be described as follows, at the level of Dynkin diagrams (see figure 4). The Cartan node of the exceptional divisor E_2 splits into three branches, one of which gets identified with E_1 and therefore cannot be used for detecting Freed-Witten anomalies,⁵ while the other two can be used as fibers for two good detecting 4-cycles.⁶ Moreover, the Cartan nodes of E_i for $i = 3, \dots, M-2$ separately split into two branches, all of which can in principle be used as fibers for two good detecting 4-cycles. Each of those gets identified with one branch belonging to each of the two neighboring nodes, with the only exception of one branch of E_{M-2} . The latter, together with one branch of E_2 , E_0 and the last divisor E_{M-1} , does not undergo any identification. Thus, in total, there are $M-3$ identifications. One can easily see that this pattern exactly reproduces the extended Dynkin diagram of the $SO(2M)$ gauge group.

All of this can be easily verified from the general algebraic equation of the Calabi-Yau hypersurface given in appendix A. In particular, the explicit expressions of some of these new nodes fibered over the curve $\{P = Q = 0\}$ are particularly easy. They are the branches which do undergo identifications

$$C_{(4)j} : \begin{cases} P = 0 \\ Q = 0 \\ v_{j-1} = 0 \\ v_j = 0 \end{cases}, \quad j = 2, \dots, M-2. \quad (4.17)$$

As stressed above, the first of these new 4-cycles ($j = 2$) is not relevant for our purposes of detecting anomalies. The general formula of the integral of $c_2(\tilde{Z}_4)$, given in eq. (A.5), on the surfaces (4.17) with, say, $j = 3$ is given by the following very simple expression:

$$\int_{C_{(4)3}} c_2(\tilde{Z}_4) = \int_{B_3} \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (4.18)$$

The proof proceeds very much analogously to the case covered in the previous section:

⁵Recall that E_1 interpolates between two ordinary D-branes of the stack.

⁶There is a subtlety here. For $SU(4)$ only this is not a priori clear because of the extra requirement of factorization one has to impose for the $SO(8)$ enhancement [26]. Therefore our second method works correctly starting from $SU(5)$.

One realizes that, on $C_{(4)3}$, the following coordinates are fixed to be non-zero: $(X, Y, \sigma, Z, v_5, \dots, v_{M-1})$. Hence, their divisor classes vanish on the 4-cycle. We then identify the following six-form as the Poincaré dual to B_3 in the ambient sixfold X_6 :

$$PD_{B_3} \equiv v_1 \cdot v_2 \cdot v_3. \quad (4.19)$$

We will also need the following linear relations, valid on $C_{(4)3}$:

$$[v_2] = -2c_1(B_3) + 2\mathcal{D} - [v_1]; \quad [v_3] = -c_1(B_3) + \mathcal{D} - [v_1]; \quad [v_4] = 3c_1(B_3) - 2\mathcal{D} + [v_1]. \quad (4.20)$$

Finally, we need the relation $[v_1] \cdot [v_4] = 0$, which gives us $[v_1]^2 = [v_1] \cdot (2\mathcal{D} - 3c_1(B_3))$

Now, we substitute all of these relations into the adjunction formula for the Chern class of $\tilde{Z}_4$, and arrive at the following result:

$$\int_{C_{(4)3}} c_2(\tilde{Z}_4) = \int_{X_6} [v_1] \cdot [v_2] \cdot [v_3] \cdot \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (4.21)$$

$$= \int_{B_3} \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q} = \int_C \mathcal{D}. \quad (4.22)$$

Again, we emphasize that, as explained in the last paragraph of 3.2.2, this 4-cycle cannot be a matter surface. Those should always yield integral periods for G_4 .

Although we provided a perturbative IIA explanation for how to detect the half-integral shift, the four-cycles we use in F-theory suggest something non-perturbative. As we can see for example in figure 4, the nodes of the Dynkin diagram we are using suggest that, at the non-perturbative level, the half-integral shift of fluxes is detected by string-junctions in IIB theory. It would be interesting to flesh out the details of this picture.

4.3 The example of SU(5)

We will now illustrate all of our general formulae in the interesting case of SU(5). The Weierstrass model for the fourfold such that it creates an SU(5) singularity along a divisor given by $D = 0$ in B_3 is given by the following complete intersection:

$$Y^2 + a_1 X Y Z + a_{3,2} \sigma^2 Y Z^3 = \quad (4.23)$$

$$= X^3 + a_{2,1} \sigma X^2 Z^2 + a_{4,3} \sigma^3 X Z^4 + a_{6,5} \sigma^5 Z^6$$

$$D = \sigma, \quad (4.24)$$

where σ is an auxiliary coordinate. This space is singular at $X = Y = \sigma = 0$. In order to resolve it, we introduce four new coordinates $v_1, \dots, v_4$. The full ambient space of the resolved fourfold $\tilde{Z}_4$ is a projective bundle over B_3 . The fiber is a toric space described by the following table:

$$W\mathbb{P}^3_{\sigma XY Z v_1 \dots v_4} : \begin{array}{c|cccccccc} & \sigma & X & Y & Z & v_1 & v_2 & v_3 & v_4 \\ \hline & 0 & 2 & 3 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 1 & 1 & 0 & -1 & 0 & 0 & 0 \\ & 0 & 0 & 1 & 0 & 1 & -1 & 0 & 0 \\ & 0 & 1 & 0 & 0 & 0 & 1 & -1 & 0 \\ & 0 & 0 & 1 & 0 & 0 & 0 & 1 & -1 \end{array} \quad (4.25)$$

The SR ideal for this space is the following:

$$SR_{\text{SU}(5)} : \left\{ XYZ ; \sigma XY ; v_i Z|_{i=1,\dots,4} ; v_1 Y ; v_3 Y ; v_2 X ; \sigma v_3 ; \sigma v_4 ; v_1 v_4 \right\}. \quad (4.26)$$

The resolved fourfold $\tilde{Z}_4$ is then given by the following complete intersection:

$$\tilde{Z}_4^{\text{SU}(M)} : \begin{cases} v_2 v_4 Y^2 + a_1 X Y Z + a_{3,2} \sigma^2 v_1 v_2 Y Z^3 = \\ = X^3 v_1 v_3^2 v_4 + a_{2,1} \sigma v_1 v_3 X^2 Z^2 + a_{4,3} \sigma^3 v_1^2 v_2 v_3 X Z^4 + a_{6,5} \sigma^5 v_1^3 v_2^2 v_3 Z^6 \\ \sigma v_1 v_2 v_3 v_4 = D. \end{cases} \quad (4.27)$$

We now impose the following ansatz on the complex structure moduli of our resolved fourfold:

$$\begin{aligned} D &\equiv P \hat{D} + Q \tilde{D} \\ a_1 &\equiv P \hat{a}_1 + Q \tilde{a}_1, \end{aligned} \quad (4.28)$$

where P and Q are polynomials in B_3 defining a holomorphic curve $\mathcal{C}$ as the complete intersection in B_3 . The first line of this ansatz simply forces $\mathcal{C} \subset \tilde{Z}_4$ to be contained in the 7-brane at $D = 0$, whereas the second enforces that a_1 vanishes along it. This is the putative curve along which one measures the Freed-Witten shift in the gauge flux quantization.

Now, we construct our distinguished 4-cycle as a complete intersection of four equations in X_6 :

$$C_{(4)} : \quad P = 0 \quad \cap \quad Q = 0 \quad \cap \quad v_2 = 0 \quad \cap \quad v_3 = 0. \quad (4.29)$$

This is the 4-cycle along which we will detect the shift in the quantization of G_4 .

We note, that on $C_{(4)}$, the following coordinates are barred from vanishing: Z, X, Y, σ , leaving only v_1 and v_4 unfixed.

We follow the procedure described in 4.2.2. By applying the linear relations, we arrive at the following result:

$$\int_{C_{(4)}} c_2(\tilde{Z}_4) = \int_{X_6} [v_1] \cdot [v_2] \cdot [v_3] \cdot \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (4.30)$$

$$= \int_{\mathcal{C}} \mathcal{D}. \quad (4.31)$$

In the last line, we use the fact that, once we set $v_1 = v_2 = v_3 = 0$, then all coordinates of X_6 are ‘gauge-fixed’ to one, except for those of B_3 . Hence, this locus corresponds to an embedded copy of B_3 in X_6 .

This formula should be interpreted as follows: Suppose we have a IIB setup with a stack of branes along a divisor $\{D = 0\}$, that we suspect is a non-spin four-dimensional manifold. Suppose that its non-spin-ness can be detected on some curve $\mathcal{C}$ given by $\{P = Q = 0\} \subset B_3$. In other words, suppose that $\int_{\mathcal{C}} D$ is odd.⁷ Then $c_2(\tilde{Z}_4)$ will be odd along the so-constructed complex surface $C_{(4)}$ that is a $\mathbb{P}^1$ -fibration over $\mathcal{C}$. Again, we emphasize that, as explained in the last paragraph of 3.2.2, this 4-cycle cannot be a matter surface. Those should always yield integral periods for G_4 .

⁷This assumes that B_3 is itself spin, i.e. $c_1(B_3)$ is even. We hope to address the case of non-spin B_3 in the future.

5 U(1)-restrictions

Having treated situations where the Calabi-Yau fourfold develops symplectic and unitary singularities along a divisor on the base manifold, we would now like to take our analysis one step further, by implementing yet another restriction on the complex structure moduli space: The so-called U(1) – *restriction*. This restriction, introduced in [24], consists in setting the a_6 polynomial in the ‘Tate form’ of the Weierstrass equation to zero: $a_6 \equiv 0$. The effect of this restriction can be understood from the perturbative IIB point of view as follows: The setups we have been considering contain a stack of 7-branes with U(N) or Sp(N) gauge groups along some divisor $\mathcal{D}$. However, the tadpole cancellation of the axio-dilaton implies that there must be another 7-brane that saturates the rest of the available negative charge generated by the O7-plane. This brane is generically invariant under the orientifold involution, and hence carries gauge group $O(1)$.

Imposing the U(1) restriction splits up this remaining brane into a brane/image-brane pair, thereby allowing the pair to enhance to a U(1) gauge group. This situation is more interesting if one wishes to be able to turn on chirality-inducing fluxes.

From the F-theory point of view, a singularity is created along the curve where the brane meets its image. The singularity type is recognizable as a curve of conifolds. To be more explicit, let us take again our example of the SU(5) model. The U(1)-restricted Tate form takes the form

$$v_2 v_4 Y^2 + a_1 X Y Z + a_{3,2} \sigma^2 v_1 v_2 Y Z^3 = X^3 v_1 v_3^2 v_4 + a_{2,1} \sigma v_1 v_3 X^2 Z^2 + a_{4,3} \sigma^3 v_1^2 v_2 v_3 X Z^4$$

which can be rewritten as follows:

$$Y \tilde{Y} = X K, \quad (5.1)$$

where

$$\begin{aligned} \tilde{Y} &\equiv v_2 v_4 Y + a_1 X Z + a_{3,2} \sigma^2 v_1 v_2 Z^3, \\ K &\equiv X^2 v_1 v_3^2 v_4 + a_{2,1} \sigma v_1 v_3 X Z^2 + a_{4,3} \sigma^3 v_1^2 v_2 v_3 Z^4. \end{aligned} \quad (5.2)$$

This is manifestly singular at the locus $Y = \tilde{Y} = X = K = 0$. In order to resolve this, we can apply the method of the small resolution, as was done in [14]. We construct a $\mathbb{P}^1$ -bundle over X_6 with coordinates (s, t) , and define the resolved fourfold as the solution space of the matrix equation:

$$\begin{pmatrix} Y & K \\ X & \tilde{Y} \end{pmatrix} \begin{pmatrix} s \\ t \end{pmatrix} = 0,$$

intersected with the usual equation

$$D = \sigma v_1 \dots v_4. \quad (5.3)$$

We could also pick the transposed equation, but for simplicity we will choose this one. In order for this equation to be consistent, the new coordinates must transform appropriately under the $\mathbb{C}^*$ -actions that act on all other coordinates. It is sufficient to impose the following linear relations among divisor classes:

$$[s] - [t] = [K] - [Y] = [\tilde{Y}] - [X]. \quad (5.4)$$

An additional $\mathbb{C}^*$ -action is introduced such that (s, t) form a $\mathbb{P}^1$. The new ambient space, which is a $\mathbb{P}^1$ -bundle over X_6 , is a *seven-fold* X_7 . Its Stanley-Reisner ideal SR_{X_7} is simply given by all elements of SR_{X_6} , plus the element st .

Let us study the cases $SU(M \geq 5)$ for simplicity. The 4-cycles that we defined in the previous section for detecting shifted quantization conditions are now defined by the same equations as before, supplemented by the equation $s = 0$:

$$C_{(4)} : P = 0 \quad \cap \quad Q = 0 \quad \cap \quad v_2 = 0 \quad \cap \quad v_3 = 0 \quad \cap \quad s = 0. \quad (5.5)$$

Now, we apply similar techniques as before to carry out our calculations, and arrive at the following general formula:

$$\int_{C_{(4)}} c_2(\tilde{Z}_4) = \int_{X_7} [v_1] \cdot [v_2] \cdot [v_3] \cdot [s] \cdot \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (5.6)$$

$$= \int_{B_3} \mathcal{D} \cdot \mathcal{P} \cdot \mathcal{Q}. \quad (5.7)$$

Hence, the $U(1)$ -restriction does not alter the results on quantization for $SU(M \geq 5)$. For the Sp -series we conjecture that odd-rank Sp groups lead to even second Chern classes and vice-versa. This is because in the odd-rank cases the two branches of the Whitney umbrella brane are separately non-spin and the induced flux on them is also half-integral. This effect cancels the analog effect arising on the non-abelian stack. In contrast, in the even-rank cases, the branches of the Whitney umbrella are spin and the induced gauge flux is integrally quantized. For this reason we still find, by explicit computation, an odd second Chern class also for the $U(1)$ -restricted $SU(4)$ -model.

6 Outlook

Our analysis suggests many directions for further investigation. For some gauge groups, we find that it is the (non)-spin-ness of the tangent bundle of the D7-stack that will determine, whether G_4 in the corresponding resolved fourfold will be integrally quantized or not. In others, it is the normal bundle in B_3 that matters. For even $c_1(B_3)$, both have the same implications. The case for non-spin base manifold needs clarification.

The Sen limit for $SU(N)$ gauge groups induces conifold singularities in the CY double-cover of B_3 that do not seem to admit any viable resolutions. In order to decide on topological matters such as the spin-ness of a divisor, we need to know the appropriate treatment of such singularities.

The elusive case of $SU(3)$ needs elucidation. Its peculiarity probably stems from the fact that its enhancement along the orientifold does not yield an $SO(6)$ group.

Acknowledgments

A. C. is a Research Associate of the Fonds de la Recherche Scientifique F.N.R.S. (Belgium). R. S. wants to thank the Theory Division of CERN for hospitality at some stage of this work. We would like to thank A. Braun, I. Garcia-Etxebarria, T. Grimm, W. Lerche for discussions and especially R. Valandro for initial collaboration to the project.

A Geometry of SU(N) F-theory configurations

We collect here some general formulae valid for the blown-up elliptic fourfold describing an SU(M) F-theory configuration. Let us define for convenience the operations $\bar{x} := x \bmod 2$, $\hat{x} := [(x-1)/2]$, with $\hat{0} = 0$, and $\tilde{x} := [x/2]$, where $[\cdot]$ extracts the integer part.

Our Calabi-Yau fourfolds $\tilde{Z}_4^{\text{SU}(M)}$ are defined as complete intersections in an ambient sixfold X_6 , constructed as projective bundles over B_3 . The table of projective weights defining the ambient fiber over B_3 , in a slightly different basis with respect to (3.14) looks like

$$W\mathbb{P}_{\sigma XYZ v_1 v_2 \dots v_{M-2} v_{M-1}}^3 : \begin{array}{c|cccccccccc} \sigma & X & Y & Z & v_1 & v_2 & v_3 & \cdots & v_{M-2} & v_{M-1} \\ \hline 0 & 2 & 3 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & 1 & 0 & -1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & -1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & -1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \overline{(M-1)} & \overline{M} & 0 & 0 & 0 & 0 & \cdots & 1 & -1 \end{array} \quad (\text{A.1})$$

From the table (A.1) one can easily infer the Stanley-Reisner ideal of the ambient variety. However, one encounters a subtlety for SU(3), which has to be treated separately. Therefore, for $M \neq 3$, the SR ideal is made of $2 + M(M-1)/2$ elements and it looks like

$$SR_{\text{SU}(M)} : \left\{ XYZ ; v_0 XY ; v_i Z|_{i=1, \dots, M-1} ; v_{2i-1} Y|_{i=1, \dots, \widehat{M}} ; v_{2i} X|_{i=1, \dots, \widetilde{M-1}} ; v_i v_j \left| \begin{array}{l} i, j=0, \dots, M-1 \\ j-i > 2 \end{array} \right. \right\}, \quad (\text{A.2})$$

where we have defined $v_0 := \sigma$.

The Stanley-Reisner ideal for SU(3) has instead 6 elements

$$SR_{\text{SU}(3)} : \left\{ XYZ ; v_0 XY ; v_1 Z ; v_2 Z ; v_1 Y ; v_0 X v_2 \right\}. \quad (\text{A.3})$$

The general formula for the proper transform, valid for every M , is

$$\tilde{Z}_4^{\text{SU}(M)} : \left\{ \begin{array}{l} \prod_{i=1}^{\widehat{M}} v_{2i} Y^2 + a_1 XYZ + a_{3, \widetilde{M}} v_0 \prod_{i=0}^{M-1} v_i^{\widehat{M-i-1}} Y Z^3 = \\ = X^3 \prod_{i=1}^{\widetilde{M}} v_{2i-1}^i \prod_{j=1}^{\widehat{M}} v_{2j}^{j-1} + a_{2,1} v_0 \prod_{i=1}^{\widetilde{M}} v_{2i-1} X^2 Z^2 + \\ + a_{4, \widehat{M+1}} v_0 \prod_{i=0}^{M-1} v_i^{\widehat{M-i}} X Z^4 + a_{6, M} v_0 \prod_{i=0}^{M-1} v_i^{M-i-1} Z^6 \\ \prod_{i=0}^{M-1} v_i = D, \end{array} \right. \quad (\text{A.4})$$

where the multi-degree of the first equation is $(6, 2, 3, 4, \dots, M-2, M-1, M)$.

The second Chern class of the blown-up fourfold (A.4) is again conveniently split in pieces

$$c_2 \left(\tilde{Z}_4^{\text{SU}(M)} \right) = c_2 \left(\tilde{Z}_4^{\text{Sp}(\widetilde{M})} \right) + \Delta c'_2, \quad (\text{A.5})$$

where the first term is the c_2 of the fourfold corresponding to the parent $\mathrm{Sp}(\widetilde{M})$ singularity and it is given by formula (3.17). The second term clearly vanishes for $M = 2$, for the special case $M = 3$ is

$$\Delta c_2'^{\mathrm{SU}(3)} = E_2 \left[-10 c_1(B_3) + \mathcal{D} + 2 E_1 + E_2 \right]. \quad (\text{A.6})$$

and, finally, has the following form for $M \geq 4$

$$\begin{aligned} \Delta c_2' = E_2 \left[-8 c_1(B_3) + \mathcal{D} + E_1 \right] + \\ - \sum_{i=2}^{\widehat{M}} E_{2i} \left[2 E_{2i-2} + E_{2i} + (7i+3) c_1(B_3) - (i^2 + i + 1) \mathcal{D} \right], \end{aligned} \quad (\text{A.7})$$

B The $\mathrm{SU}(3)$ case

We have stressed several times that F-theory configurations with $\mathrm{SU}(3)$ gauge group are somewhat special and have to be treated separately. Moreover, we have not been able to find for them any non-trivial shift to the quantization rule of the G_4 flux. In this appendix we want to give an hint that, in fact, such a shift is unlikely arising for a Calabi-Yau fourfold with an $\mathrm{SU}(3)$ singularity blown-up in the usual way [26], i.e. by introducing the two toric vectors v_1, v_2 .

By inspecting the expression of $c_2(\tilde{Z}_4^{\mathrm{SU}(3)})$ (see (A.5) and (A.6)), one realizes that this quantity is even if and only if the class $c_1 E_1 + D(E_1 + E_2) + E_2^2$ is even. Therefore

$$c_2(\tilde{Z}_4^{\mathrm{SU}(3)}) \sim c_1 E_1 + D(E_1 + E_2) + E_2^2 \quad \text{mod } H^4(\tilde{Z}_4^{\mathrm{SU}(3)}, 2\mathbb{Z}). \quad (\text{B.1})$$

This is because all the other summand appear with an even numerical factor in front and they are Poincaré dual to effective 4-cycles of the Calabi-Yau fourfold. Moreover, as one can see from eq. (A.4), the equation of the $\mathrm{SU}(3)$ Calabi-Yau hypersurface contains the term $v_2 Y^2$. This term is necessary, in the sense that we cannot deform it away without making the hypersurface singular.⁸ But now the two elements of the Stanley-Reisner ideal of the ambient variety (A.3) $v_0 XY$ and $v_0 X v_2$ tell us that, on the blown-up fourfold

$$D(E_1 + E_2) + E_2^2 \sim E_1^2 \quad \text{mod } H^4(\tilde{Z}_4^{\mathrm{SU}(3)}, 2\mathbb{Z}). \quad (\text{B.2})$$

Finally, the element $v_1 Y$ implies that

$$E_1^2 \sim c_1 E_1 \quad \text{mod } H^4(\tilde{Z}_4^{\mathrm{SU}(3)}, 2\mathbb{Z}). \quad (\text{B.3})$$

While odd values of c_2 could still be found in principle on non-holomorphic 4-cycles, this argument nevertheless points towards the conjecture that in fact $c_2(\tilde{Z}_4^{\mathrm{SU}(3)})$ is always an even class.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

⁸One can indeed check that, for instance, the codimension two locus $\{\sigma = X = Z = 0\}$ would become singular.

References

- [1] A. Collinucci, *New F-theory lifts*, *JHEP* **08** (2009) 076 [[arXiv:0812.0175](#)] [[INSPIRE](#)].
- [2] A. Collinucci, *New F-theory lifts. II. Permutation orientifolds and enhanced singularities*, *JHEP* **04** (2010) 076 [[arXiv:0906.0003](#)] [[INSPIRE](#)].
- [3] R. Blumenhagen, T.W. Grimm, B. Jurke and T. Weigand, *F-theory uplifts and GUTs*, *JHEP* **09** (2009) 053 [[arXiv:0906.0013](#)] [[INSPIRE](#)].
- [4] R. Blumenhagen, T.W. Grimm, B. Jurke and T. Weigand, *Global F-theory GUTs*, *Nucl. Phys. B* **829** (2010) 325 [[arXiv:0908.1784](#)] [[INSPIRE](#)].
- [5] T.W. Grimm, S. Krause and T. Weigand, *F-theory GUT vacua on compact Calabi-Yau fourfolds*, *JHEP* **07** (2010) 037 [[arXiv:0912.3524](#)] [[INSPIRE](#)].
- [6] J. Marsano, N. Saulina and S. Schäfer-Nameki, *F-theory compactifications for supersymmetric GUTs*, *JHEP* **08** (2009) 030 [[arXiv:0904.3932](#)] [[INSPIRE](#)].
- [7] C.-M. Chen, J. Knapp, M. Kreuzer and C. Mayrhofer, *Global SO(10) F-theory GUTs*, *JHEP* **10** (2010) 057 [[arXiv:1005.5735](#)] [[INSPIRE](#)].
- [8] M. Esole and S.-T. Yau, *Small resolutions of SU(5)-models in F-theory*, [arXiv:1107.0733](#) [[INSPIRE](#)].
- [9] M. Esole, J. Fullwood and S.-T. Yau, *D₅ elliptic fibrations: non-Kodaira fibers and new orientifold limits of F-theory*, [arXiv:1110.6177](#) [[INSPIRE](#)].
- [10] J. Marsano, N. Saulina and S. Schäfer-Nameki, *On G-flux, M5 instantons and U(1)s in F-theory*, [arXiv:1107.1718](#) [[INSPIRE](#)].
- [11] J. Marsano, N. Saulina and S. Schäfer-Nameki, *A note on G-fluxes for F-theory model building*, *JHEP* **11** (2010) 088 [[arXiv:1006.0483](#)] [[INSPIRE](#)].
- [12] T.W. Grimm, T.-W. Ha, A. Klemm and D. Klevers, *Computing brane and flux superpotentials in F-theory compactifications*, *JHEP* **04** (2010) 015 [[arXiv:0909.2025](#)] [[INSPIRE](#)].
- [13] T.W. Grimm, M. Kerstan, E. Palti and T. Weigand, *Massive Abelian gauge symmetries and fluxes in F-theory*, *JHEP* **12** (2011) 004 [[arXiv:1107.3842](#)] [[INSPIRE](#)].
- [14] A.P. Braun, A. Collinucci and R. Valandro, *G-flux in F-theory and algebraic cycles*, *Nucl. Phys. B* **856** (2012) 129 [[arXiv:1107.5337](#)] [[INSPIRE](#)].
- [15] S. Krause, C. Mayrhofer and T. Weigand, *G₄ flux, chiral matter and singularity resolution in F-theory compactifications*, *Nucl. Phys. B* **858** (2012) 1 [[arXiv:1109.3454](#)] [[INSPIRE](#)].
- [16] C. Vafa, *Evidence for F-theory*, *Nucl. Phys. B* **469** (1996) 403 [[hep-th/9602022](#)] [[INSPIRE](#)].
- [17] F. Denef, *Les Houches lectures on constructing string vacua*, [arXiv:0803.1194](#) [[INSPIRE](#)].
- [18] T. Weigand, *Lectures on F-theory compactifications and model building*, *Class. Quant. Grav.* **27** (2010) 214004 [[arXiv:1009.3497](#)] [[INSPIRE](#)].
- [19] E. Witten, *On flux quantization in M-theory and the effective action*, *J. Geom. Phys.* **22** (1997) 1 [[hep-th/9609122](#)] [[INSPIRE](#)].
- [20] R. Minasian and G.W. Moore, *K-theory and Ramond-Ramond charge*, *JHEP* **11** (1997) 002 [[hep-th/9710230](#)] [[INSPIRE](#)].

- [21] D.S. Freed and E. Witten, *Anomalies in string theory with D-branes*, *Asian J. Math* **3** (1999) 819 [[hep-th/9907189](#)] [[INSPIRE](#)].
- [22] A. Collinucci and R. Savelli, *On flux quantization in F-theory*, *JHEP* **02** (2012) 015 [[arXiv:1011.6388](#)] [[INSPIRE](#)].
- [23] S. Krause, C. Mayrhofer and T. Weigand, *Gauge fluxes in F-theory and type IIB orientifolds*, [arXiv:1202.3138](#) [[INSPIRE](#)].
- [24] T.W. Grimm and T. Weigand, *On Abelian gauge symmetries and proton decay in global F-theory GUTs*, *Phys. Rev. D* **82** (2010) 086009 [[arXiv:1006.0226](#)] [[INSPIRE](#)].
- [25] A. Collinucci, F. Denef and M. Esole, *D-brane deconstructions in IIB orientifolds*, *JHEP* **02** (2009) 005 [[arXiv:0805.1573](#)] [[INSPIRE](#)].
- [26] M. Bershadsky et al., *Geometric singularities and enhanced gauge symmetries*, *Nucl. Phys. B* **481** (1996) 215 [[hep-th/9605200](#)] [[INSPIRE](#)].
- [27] W. Fulton, *Intersection theory*, Springer-Verlag, Berlin Germany and New York U.S.A. (1998).
- [28] B. Andreas and G. Curio, *On discrete twist and four flux in $N = 1$ heterotic/F theory compactifications*, *Adv. Theor. Math. Phys.* **3** (1999) 1325 [[hep-th/9908193](#)] [[INSPIRE](#)].
- [29] T.W. Grimm, *The $N = 1$ effective action of F-theory compactifications*, *Nucl. Phys. B* **845** (2011) 48 [[arXiv:1008.4133](#)] [[INSPIRE](#)].
- [30] T.W. Grimm and R. Savelli, *Gravitational instantons and fluxes from M/F-theory on Calabi-Yau fourfolds*, *Phys. Rev. D* **85** (2012) 026003 [[arXiv:1109.3191](#)] [[INSPIRE](#)].
- [31] R. Donagi and M. Wijnholt, *Higgs bundles and UV completion in F-theory*, [arXiv:0904.1218](#) [[INSPIRE](#)].
- [32] M. Cvetič, I. Garcia-Etxebarria and J. Halverson, *Global F-theory models: instantons and gauge dynamics*, *JHEP* **01** (2011) 073 [[arXiv:1003.5337](#)] [[INSPIRE](#)].