RECEIVED: May 22, 2014

REVISED: July 14, 2014

ACCEPTED: July 24, 2014

PUBLISHED: August 11, 2014

Low tension strings on a cosmological singularity

Ben Craps,^{a,b} Chethan Krishnan^c and Ayush Saurabh^{c,d,e}

^a*Theoretische Natuurkunde, Vrije Universiteit Brussel, and International Solvay Institutes,
Pleinlaan 2, B-1050 Brussels, Belgium*

^b*Laboratoire de Physique Théorique, École Normale Supérieure,
24 rue Lhomond, F-75231 Paris Cedex 05, France*

^c*Center for High Energy Physics, Indian Institute of Science,
Bangalore – 560012, India*

^d*International Center for Theoretical Sciences, Indian Institute of Science Campus,
Bangalore – 560012, India*

^e*School of M.A.C.E., The University of Manchester,
Manchester – M13 9PL, United Kingdom*

E-mail: Ben.Craps@vub.ac.be, chethan.krishnan@gmail.com,
ayushsaurabh@hotmail.com

ABSTRACT: It has recently been argued that the singularity of the Milne orbifold can be resolved in higher spin theories. In string theory scattering amplitudes, however, the Milne singularity gives rise to ultraviolet divergences that signal uncontrolled backreaction. Since string theory in the low tension limit is expected to be a higher spin theory (although precise proposals only exist in special cases), we investigate what happens to these scattering amplitudes in the low tension limit. We point out that the known problematic ultraviolet divergences disappear in this limit. In addition we systematically identify all divergences of the simplest 2-to-2 tree-level string scattering amplitude on the Milne orbifold, and argue that the divergences that survive in the low tension limit have sensible infrared interpretations.

KEYWORDS: Higher Spin Gravity, Scattering Amplitudes, Bosonic Strings, Spacetime Singularities

ARXIV EPRINT: [1405.3935](https://arxiv.org/abs/1405.3935)

Contents

1	Introduction	1
2	Review: the 4-point string amplitude on Milne	2
3	Structure of the integral	4
3.1	Case-1: $m_4 > m_1$ and $m_3 > m_2$	5
3.2	Case-2: $m_4 > m_1$ and $m_2 > m_3$	7
4	Divergences of the four-point function	8
4.1	Type-1: divergences from boundaries, unrelated to poles	10
4.2	Type-2: poles at boundary	11
4.3	Type-3: poles at Mandelstam maxima/minima	14
4.4	Type-4: IR divergences identified in [1]	16
5	Summary of divergences	16

1 Introduction

Singularities in time-dependent backgrounds are not well-understood in string theory. One context where some effort in this direction has been expended is in the case of time-dependent orbifolds of flat space; see [2–6] for reviews. Being descendants of flat space makes these geometries amenable to string perturbation theory, but it has been found that, already at tree level, string amplitudes in them have UV divergences arising from uncontrolled backreaction at the singularity [1, 7–9]. In this paper, we will revisit the string amplitudes in [1] on the Milne geometry, which has a spacelike orbifold singularity.

Recently, it was observed in [10, 11] that certain cosmological singularities arising as quotients of dS_3 or flat space can in fact be resolved, if we embed these singularities into higher spin theories.¹ In particular, [11] (see also [14]) shows that the Milne singularity can be resolved in the context of flat space higher spin theories [15, 16]. The idea is that one can get rid of the metric singularity via higher spin gauge transformations, while preserving the holonomy in the Chern-Simons language.

In the tensionless $\alpha' \rightarrow \infty$ limit of string theory, the tower of higher spin string states becomes massless. It is expected that this limit of string theory is captured by a massless, interacting higher spin theory: in AdS various arguments have been presented to make this correspondence more concrete [17–21]. In the flat space case, it is unclear what the precise statement is, but one expects that the 2+1 D higher spin theory of [15, 16] should morally capture some aspects of string theory in the tensionless limit, even in flat space.

¹The latter were formulated as Chern-Simons theories in 2+1 dimensions [12, 13].

If this belief is correct, one would expect that the string scattering amplitudes in Milne should be well-defined in the large α' limit. This is because the scattering amplitude is gauge invariant, so in the $\alpha' \rightarrow \infty$ limit it should be well-defined if the singularity is a gauge artifact in the higher spin picture. In this paper, we will do a scan of the divergences of the Milne 2-to-2 tree-level string scattering amplitude and show that it is indeed UV finite when the (dimensionless) α' is large enough. (We will be more precise about this in the next section.) Our analysis is exhaustive, and we will find infinite classes of divergences. However, we will argue that the divergences that survive in the large α' limit are all IR divergences that either (a) have been previously noticed in [1], or (b) can be understood to be arising from the tower of intermediate string states going on-shell and therefore are expected on physical grounds.

The purpose of our analysis is to show that the problematic tree-level divergences identified in [1] disappear in the large α' limit, which resolves the apparent tension between the pathological behavior of tree-level string scattering amplitudes on the Milne orbifold [1] and the recent results suggesting that higher spin theory is well-behaved on the same space [11]. The possible connection between string theory and higher spin theory suggests that string loop corrections should also be well-behaved in the large α' limit, but this would be much harder to verify directly, and we will not attempt to do so here.

The paper is organized as follows. We first review the 2-to-2 tree-level string scattering amplitude on the Milne geometry (section 2) to fix our notations and to lay the groundwork for the discussions in the following sections. Section 3 describes various relevant features of the integrand and section 4 undertakes a careful scan of the divergences that can arise in this integral. This is the main technical part of the paper. Section 5 summarizes the various divergences and categorizes them as UV or IR.

2 Review: the 4-point string amplitude on Milne

The Milne orbifold is obtained from Minkowski space $ds^2 = -2dX^+dX^- + d\vec{X}^2$ by the boost identification $X^\pm \rightarrow \exp(\pm 2\pi)X^\pm$. In [1], the four-point function $\langle \psi_3^* \psi_4^* \psi_1 \psi_2 \rangle$ of tachyon vertex operators

$$\psi_{m_j, l_j, \vec{p}_j} = \frac{e^{i\vec{p}_j \cdot \vec{X}}}{2\sqrt{2\pi i}} \int_{-\infty}^{\infty} dw e^{\frac{i}{\sqrt{2}}(m_j X^- e^{-w} + m_j X^+ e^w)} e^{il_j w} \quad (2.1)$$

in tree-level bosonic string theory was computed to be

$$\sum \frac{(2\pi)^{24}}{4} \delta^{24} \left(\sum \epsilon_i \vec{p}_i \right) \delta \left(\sum \epsilon_i l_i \right) \int_0^\infty dv_4 G(s) G(t) G(u) \frac{v_2^{il_2+1} v_3^{-il_3+1} v_4^{-il_4-1}}{|m_2 m_3 (v_2^2 - v_3^2)|}. \quad (2.2)$$

The ϵ_i are +1 for the incoming particles (1 and 2), and -1 for the outgoing particles (3 and 4). The momenta l_i along the Milne circle are integers and we will set them to zero in what follows because they are phases, and not crucial for the divergence/convergence discussion we undertake. Working with momenta measured in string units (which amounts to setting $\alpha' = 1$), the mass shell condition reads

$$m^2 = -4 + \vec{p}^2. \quad (2.3)$$

The parameter m^2 ($m > 0$) is the effective 2-D mass squared. We have defined

$$G(x) = \frac{\Gamma(-1 - \frac{x}{4})}{\Gamma(2 + \frac{x}{4})} \quad (2.4)$$

and the Mandelstam variables are

$$s = -(p_1 + p_2)^2 = -8 + m_1 m_2 \left(v_2 + \frac{1}{v_2} \right) - 2\vec{p}_1 \cdot \vec{p}_2, \quad (2.5)$$

$$t = -(p_1 - p_3)^2 = -8 - m_1 m_3 \left(v_3 + \frac{1}{v_3} \right) + 2\vec{p}_1 \cdot \vec{p}_3, \quad (2.6)$$

$$u = -(p_1 - p_4)^2 = -8 - m_1 m_4 \left(v_4 + \frac{1}{v_4} \right) + 2\vec{p}_1 \cdot \vec{p}_4. \quad (2.7)$$

A standard constraint is

$$s + t + u = -16. \quad (2.8)$$

Also v_2 and v_3 are defined by²

$$v_2 = \frac{AB + m_2^2 - m_3^2 \mp \sqrt{\Delta}}{2m_2 B}, \quad (2.9)$$

$$v_3 = \frac{-AB + m_2^2 - m_3^2 \mp \sqrt{\Delta}}{2m_3 B}, \quad (2.10)$$

where

$$A = -m_1 + m_4 v_4, \quad (2.11)$$

$$B = -m_1 + \frac{m_4}{v_4}, \quad (2.12)$$

$$\Delta = (m_2^2 - m_3^2)^2 - 2AB(m_2^2 + m_3^2) + A^2 B^2. \quad (2.13)$$

This specific form of v_2 and v_3 arises from delta functions that enforce

$$m_1 + m_2 v_2 - m_3 v_3 - m_4 v_4 = 0, \quad (2.14)$$

$$m_1 + \frac{m_2}{v_2} - \frac{m_3}{v_3} - \frac{m_4}{v_4} = 0. \quad (2.15)$$

It is also important that v_2, v_3 need to be positive, so only the positive solutions need to be retained (and summed over) in the integral. The outermost summation in the amplitude (2.2) refers to this summation over the positive branches of v_2 and v_3 .

The problematic divergence identified in [1] comes from the large v_4 region of the integral (2.2), in which the integrand is in the Regge regime, $s \rightarrow m_1 m_4 v_4 \rightarrow \infty$ with finite $t \rightarrow -(\vec{p}_1 - \vec{p}_3)^2$. This integration region gives a contribution of the form

$$\int^\infty dv_4 v_4^{-\frac{(\vec{p}_1 - \vec{p}_3)^2}{2}}, \quad (2.16)$$

which diverges whenever $\alpha'(\vec{p}_1 - \vec{p}_3)^2 < 2$, where we reinstated α' . This makes it clear that the problematic divergence disappears whenever the momentum transfer is large enough

²The upper and lower signs in the following expressions are correlated.

in string units, and therefore in the large α' limit. The physical intuition is that for large momentum transfer, string amplitudes are very soft in the Regge regime.

Our goal in the rest of the paper is to study the convergence properties of (2.2) thoroughly, in order to make sure that no problematic divergences remain in the tree level amplitude. The integral has a fair amount of structure and the problem is fairly detail oriented, so in the next section we proceed systematically to characterize the integral.

As mentioned in the introduction, a study of string loop corrections is beyond the scope of the present paper. In Minkowski space, arguments have been presented that string loop diagrams are also soft in the Regge regime for sufficiently large momentum transfer [22] (although the perturbation series could not be summed). In the Milne orbifold, one would also have to include contributions from the exchange of twisted sector strings, which we will not attempt.

3 Structure of the integral

As stated earlier, we need pairs of positive v_2 and positive v_3 among (2.9)–(2.10), out of the four possible pairs of combinations. This means we have to find out the regions of integration in (2.2) where the positivity properties of v_2 and v_3 change.

It turns out that there are three qualitatively different regions in the v_4 -half line. The defining features of these regions are governed by the parameters m_1 , m_2 , m_3 and m_4 . These parameters control where v_2 and v_3 change signs, or become complex, or are indeterminate. As we find out from (2.9)–(2.10), constituents of one of the pairs of v_2 , v_3 change signs whenever $A = 0$, and both v_2 , v_3 become indeterminate whenever $B = 0$. This defines the formal boundaries of the three adjoining regions of integration.

To understand these regions in more detail, we first introduce

$$v_{2u} = \frac{AB + m_2^2 - m_3^2 - \sqrt{\Delta}}{2m_2B}, \quad (3.1)$$

$$v_{2d} = \frac{AB + m_2^2 - m_3^2 + \sqrt{\Delta}}{2m_2B}, \quad (3.2)$$

$$v_{3u} = \frac{-AB + m_2^2 - m_3^2 - \sqrt{\Delta}}{2m_3B}, \quad (3.3)$$

$$v_{3d} = \frac{-AB + m_2^2 - m_3^2 + \sqrt{\Delta}}{2m_3B}. \quad (3.4)$$

We also define some new parameters:

$$P_{12} = \vec{p}_1 \cdot \vec{p}_2, \quad P_{13} = \vec{p}_1 \cdot \vec{p}_3, \quad P_{14} = \vec{p}_1 \cdot \vec{p}_4, \quad (3.5)$$

$$v_{4A} = \frac{m_1}{m_4}, \quad v_{4B} = \frac{m_4}{m_1}. \quad (3.6)$$

Now, it is easily inferred from (2.11)–(2.12) that A changes sign at some $v_4 = v_{4A}$ and $A > 0$ for $v_4 > v_{4A}$. Similarly, B changes sign at some $v_4 = v_{4B}$ and $B < 0$ for $v_4 > v_{4B}$.

Depending on the actual values of m_1 , m_2 , m_3 , m_4 , P_{12} , and P_{13} , we will have specific kinematic conditions defining the regions of integration and the properties of the four-point

function. These are what we call the kinematic parameters, and they fully describe the amplitude.³

For all these cases, we adopt a global convention that the m_1 would always stand for the lighter of the two incoming particles (i.e. $m_2 \geq m_1$) and the m_3 would always stand for the lighter of the two outgoing particles (i.e. $m_4 \geq m_3$). Even with this assumption, we still have four separate cases of orderings between the mass parameters that we can consider. Of these we call $m_4 > m_1$ and $m_3 > m_2$ Case-1, and $m_4 > m_1$ and $m_2 > m_3$ Case-2 and discuss them in detail. The other two cases (e.g., $m_1 > m_4$ and $m_2 > m_3$) are either kinematically impossible or have analogous divergence structures, so we can omit them.

It is also worth noticing that the parameters that we choose need to satisfy certain constraints due to the rules of vector algebra and momentum conservation:

$$\vec{p}_1^2 + \vec{p}_2^2 \geq 2|P_{12}| \iff |P_{12}| \leq \frac{m_1^2 + m_2^2}{2} + 4, \quad (3.7)$$

$$\vec{p}_1^2 + \vec{p}_3^2 \geq 2|P_{13}| \iff |P_{13}| \leq \frac{m_1^2 + m_3^2}{2} + 4, \quad (3.8)$$

$$\vec{p}_1^2 + \vec{p}_4^2 \geq 2|P_{14}| \iff |P_{14}| \leq \frac{m_1^2 + m_4^2}{2} + 4, \quad (3.9)$$

$$\vec{p}_1 + \vec{p}_2 = \vec{p}_3 + \vec{p}_4 \iff P_{14} = m_1^2 + 4 - P_{13} + P_{12}. \quad (3.10)$$

Note that P_{14} is not an independent parameter.

Now we turn to a discussion of the two cases.

3.1 Case-1: $m_4 > m_1$ and $m_3 > m_2$

In this case, $v_{4A} < v_{4B}$ and so we could define our three regions of integration as follows,

Region-I: $v_4 \in (0, v_{4A})$

Region-II: $v_4 \in (v_{4A}, v_{4B})$

Region-III: $v_4 \in (v_{4B}, \infty)$

In Region-I, $A < 0$ and $B > 0$. So, through inspection of the expressions (3.1)–(3.4), we find that $\sqrt{\Delta} > (|AB| + |m_2^2 - m_3^2|)$. Therefore in Region-I, v_{2u} and v_{3u} would always be negative, and v_{2d} and v_{3d} would always be positive. Conversely, in Region-III, through similar analysis we find that v_{2u} and v_{3u} would always be positive, and v_{2d} and v_{3d} would always be negative. In Region-II, though, we find that we do not have any pair of v_2 and v_3 in (3.1)–(3.4) in which both constituents of the pair are positive at the same time. Hence, in the present case, the integrand does not exist in Region-II.

For notational convenience, we will define the integrands in Region-I, III as follows:

$$i_1(v_{2d}(v_4), v_{3d}(v_4), v_4) = G(s(v_{2d}))G(t(v_{3d}))G(u(v_4)) \frac{v_{2d}v_{3d}v_4^{-1}}{|m_2m_3(v_{2d}^2 - v_{3d}^2)|}, \quad (3.11)$$

$$i_3(v_{2u}(v_4), v_{3u}(v_4), v_4) = G(s(v_{2u}))G(t(v_{3u}))G(u(v_4)) \frac{v_{2u}v_{3u}v_4^{-1}}{|m_2m_3(v_{2u}^2 - v_{3u}^2)|}. \quad (3.12)$$

³Note that we have set the l_i to zero as discussed earlier.

Figure 1. Plots of s, t , and u for $m_4 > m_1$, $m_3 > m_2$ for positive v_4 . The values of the kinematic parameters used for making the plot are $m_1 = 5.0$, $m_2 = 5.8$, $m_3 = 6.8$, $m_4 = 8.0$, $P_{12} = 10.5$, $P_{13} = 33.5$.

Now, the integral in Region-I (we call it I_1) can be easily shown to be the same as the integral in Region-III by an appropriate renaming of $v_4 \rightarrow \frac{1}{x}$. Specifically, the contribution to the four-point function integral from Region-III (we call it I_3) transforms as

$$\begin{aligned} I_3 &= \sum \frac{(2\pi)^{24}}{4} \delta^{24} \left(\sum \epsilon_i \vec{p}_i \right) \delta \left(\sum \epsilon_i l_i \right) \int_{v_{4B}}^{\infty} dv_4 i_3(v_{2u}(v_4), v_{3u}(v_4), v_4) \\ &\rightarrow \sum \frac{(2\pi)^{24}}{4} \delta^{24} \left(\sum \epsilon_i \vec{p}_i \right) \delta \left(\sum \epsilon_i l_i \right) \int_{v_{4A}}^0 dx i_3(v_{2u}(x^{-1}), v_{3u}(x^{-1}), x^{-1}) \times - \left(\frac{1}{x^2} \right). \end{aligned}$$

Now if after simplifications we replace the x with v_4 , we find that the transformed integral above becomes

$$I_3 \rightarrow \sum \frac{(2\pi)^{24}}{4} \delta^{24} \left(\sum \epsilon_i \vec{p}_i \right) \delta \left(\sum \epsilon_i l_i \right) \int_0^{v_{4A}} dv_4 i_1(v_{2d}(v_4), v_{3d}(v_4), v_4) = I_1.$$

Thus the net scattering amplitude in this case is given by

$$I = I_1 + I_3 = 2 I_3.$$

In figure 1 we have transformed away Region-I into Region-III and only show Region-III.

Figure 2. Plots of s , t and u for $m_4 > m_1$, $m_2 > m_3$, $(m_3 + m_4) > (m_1 + m_2)$ for positive v_4 , where the subscript u stands for functions of v_{2u} , v_{3u} and subscript d stands for functions of v_{2d} , v_{3d} . The values of the kinematic parameters used for making the plot are $m_1 = 5.0$, $m_2 = 9.5$, $m_3 = 5.5$, $m_4 = 10.0$, $P_{12} = 39.5$, $P_{13} = 4.0$.

3.2 Case-2: $m_4 > m_1$ and $m_2 > m_3$

In this case, we can choose the various regions as in Case-1. The integrand behaves very similar to the integrand in Case-1 in the $v_4 \rightarrow \infty$ limit but the difference now is that $s(v_{2u})$, $t(v_{3u})$, and u are now finite in the $v_4 \rightarrow v_{4B}$ limit and the integrand now exists in Region-II as well.

One crucial observation is that there are two possibilities to be distinguished. These two possibilities⁴ are $(m_3 + m_4) \geq (m_1 + m_2)$ and $(m_1 + m_2) \geq (m_3 + m_4)$. If the former condition is satisfied, Δ can change sign in Region-II and values become complex. So v_4 corresponding to $\Delta = 0$ defines two boundaries (we call them v_{4+} and v_{4-})⁵ for the integral within Region-II as can be seen in the figure 2. We also find that all the solutions for v_{2u} , v_{2d} , v_{3u} , and v_{3d} are either positive or complex, all at the same time for any

⁴Throughout the paper, we will be making comments about the case $(m_1 + m_2) = (m_3 + m_4)$, which can be thought of as a special case of either of the possibilities.

⁵There are four roots for the locations of $\Delta = 0$ points in v_4 space. Out of these four roots, only two lie in $v_4 > 0$. We quote them here for completeness:

$$v_{4\pm} = \frac{m_1^2 - m_2^2 - m_3^2 + m_4^2 + 2m_2m_3 \pm \sqrt{(m_1^2 - m_2^2 - m_3^2 + m_4^2 + 2m_2m_3)^2 - 4m_1^2m_4^2}}{2m_1m_4}.$$

Figure 3. Plots of s , t and u for $m_4 > m_1$, $m_2 > m_3$, $(m_1 + m_2) > (m_3 + m_4)$ for positive v_4 , where the subscript u stands for functions of v_{2u} , v_{3u} and subscript d stands for functions of v_{2d} , v_{3d} . The values of the kinematic parameters used for making the plot are $m_1 = 10.0$, $m_2 = 40.0$, $m_3 = 20.0$, $m_4 = 25.0$, $P_{12} = 20.0$, $P_{13} = 40.0$.

specific v_4 -location in Region-II. Therefore, since $i_1(v_{2d}, v_{3d}, v_4)$ and $i_3(v_{2u}, v_{3u}, v_4)$ both exist at the same time in Region-II and could also be transformed into each other using the transformation process presented in Case-1, we would essentially have to worry only about integration in the interval (v_{4+}, ∞) , which takes the form

$$I = 2 \int_{v_{4+}}^{v_{4B}} dv_4 i_1(v_{2d}(v_4), v_{3d}(v_4), v_4) + 2 \int_{v_{4+}}^{\infty} dv_4 i_3(v_{2u}(v_4), v_{3u}(v_4), v_4). \quad (3.13)$$

When $(m_1 + m_2) > (m_3 + m_4)$, however, Δ stays positive throughout Region-II. This case is shown in figure 3. Now we always have $\Delta \geq 0$ and so the Mandelstam invariants are real in whole of Region-II. This means that the net integral simply becomes

$$I = 2 \int_1^{v_{4B}} dv_4 i_1(v_{2d}(v_4), v_{3d}(v_4), v_4) + 2 \int_1^{\infty} dv_4 i_3(v_{2u}(v_4), v_{3u}(v_4), v_4). \quad (3.14)$$

4 Divergences of the four-point function

In this section we will present a systematic scan of the divergences in the four-point function. This section is technical, but our conclusions are summarized in the next section for the reader's convenience.

In the four-point function, we have multiple gamma functions which give rise to poles in the integrand. These poles are avoided by the $i\epsilon$ prescription of the Feynman propagator. This prescription differs from the Cauchy Principal Value (CPV) prescription by a delta function term, which does not give rise to a divergence. Therefore, since we are only interested in determining whether the integral is finite, it suffices to do so using a CPV prescription.

For a generic gamma function, $\Gamma(x)$, around the pole $x = -n$ ($n > 0$), we have

$$\lim_{\delta x \rightarrow 0} (\Gamma(-n - \delta x) + \Gamma(-n + \delta x)) = \frac{2(-1)^{-n}\psi(n+1)}{n!}, \quad (4.1)$$

where ψ is the digamma function. The last expression is completely finite and would imply that the integration through a gamma function pole would be finite in the sense of CPV.

Now suppose the argument of the gamma function is a function $f(x)$, with $f(x) = -n$ for some $x = x_n$. If $f'(x_n) \neq 0$, a very similar argument shows that the integral around the pole is finite. However, if x_n is a local minimum or a local maximum of f , then the pole is approached from the “same side” and the integral diverges.

To use these observations in the analysis of our four-point function, we first write the four-point function integral around a pole (at say $v_4 = v_{4n}$) in the following form:

$$\int_{v_{4n}-\delta v_4}^{v_{4n}+\delta v_4} dv_4 F(v_4) \Gamma(M(v_4)), \quad (4.2)$$

where $M(v_4)$ is a function of one of the Mandelstam invariants, and $F(v_4)$ is assumed to be continuous in the concerned interval, which would be the generic situation.⁶ By the above reasoning, poles at generic values of v_4 do not make the integral diverge. However for some specific kinematic configurations we would have special gamma function poles which cause the four-point function to diverge. This can happen when

- there is a pole at the boundary of integration so that there are no two “sides” for the poles to cancel against each other in the CPV, or
- there is a pole at a maximum/minimum of the Mandelstam variable that falls within the integration range, as we discussed above.

Now, we classify and explain all the divergences of the four-point function as follows:

Type-1: divergences from boundaries of the integral not related to poles.

Type-2: divergences from gamma function poles at the boundary of the integral.

⁶If $F(v_4)$ passes through zero at v_{4n} , one might worry that the sign change would cause a divergence. But the pole of the gamma function goes as $1/(v_4 - v_{4n})$ and therefore any fractional or integer power law approach to zero of $F(v_4)$ (which are the only cases relevant for us), will result in a converging integral. A more tricky situation arises when $F(v_4)$, instead of being continuous, has a discontinuity at v_{4n} and has opposite signs on either side of the pole. This can happen when more than one gamma function has a pole at the same location. But one can show that because of the $s+t+u = -16$ constraint, the integral is again finite: this is tied crucially to the Veneziano (or Virasoro-Shapiro) structure of the Gamma functions.

Type-3: divergences from gamma function poles occurring at the maxima/minima of Mandelstam invariants.

Type-4: IR type divergences in specific kinematic configurations.

4.1 Type-1: divergences from boundaries, unrelated to poles

For the kinematic configuration in Case-1, presented in the previous section, the two boundaries of the integral are $v_4 = v_{4B}$ and $v_4 = +\infty$. In the $v_4 \rightarrow \infty$ limit, we find that

$$\begin{aligned} v_{2u} &\rightarrow \frac{(-m_1 + m_4 v_4)}{m_2} \rightarrow \infty, & v_{3u} &\rightarrow \frac{m_3}{m_1}, \\ s &\rightarrow m_1 m_4 v_4 \rightarrow \infty, & t &\rightarrow -(\vec{p}_1 - \vec{p}_3)^2, & u &\rightarrow -m_1 m_4 v_4 \rightarrow -\infty. \end{aligned}$$

This is the Regge limit condition, $s \rightarrow \infty$ and t fixed. Under this condition the term $G(s)G(t)G(u)$ simplifies as

$$G(s)G(t)G(u) \rightarrow -\left(\frac{s}{4}\right)^{2+\frac{t}{2}} \frac{\Gamma(-1-\frac{t}{4})}{\Gamma(2+\frac{t}{4})}.$$

Using the limiting expressions above, the four-point function simplifies as

$$C \times \int^\infty dv_4 v_4^{-\frac{(\vec{p}_1 - \vec{p}_3)^2}{2}}, \quad (4.3)$$

where C is some constant. So in the $v_4 \rightarrow \infty$ limit, the four-point function will diverge whenever the exponent in the integrand is such that [1]

$$\frac{(\vec{p}_1 - \vec{p}_3)^2}{2} \leq 1 \iff (\vec{p}_1 - \vec{p}_3)^2 \leq 2.$$

Similarly in $v_4 \rightarrow v_{4B}$ limit, we have

$$\begin{aligned} v_{2u} &\rightarrow \frac{(m_2^2 - m_3^2)v_4}{m_2(m_4 - m_1 v_4)} \rightarrow \infty, & v_{3u} &\rightarrow \frac{(m_2^2 - m_3^2)v_4}{m_3(m_4 - m_1 v_4)} \rightarrow \infty, \\ s &\rightarrow \frac{m_1(m_2^2 - m_3^2)v_4}{(m_4 - m_1 v_4)} \rightarrow \infty, & t &\rightarrow -\frac{m_1(m_2^2 - m_3^2)v_4}{(m_4 - m_1 v_4)} \rightarrow \infty, \\ u &\rightarrow -(\vec{p}_1 - \vec{p}_4)^2. \end{aligned}$$

Again we have the Regge limit condition, $s \rightarrow \infty$ and u fixed, which leads to the simplification

$$G(s)G(t)G(u) \rightarrow -\left(\frac{s}{4}\right)^{2+\frac{u}{2}} \frac{\Gamma(-1-\frac{u}{4})}{\Gamma(2+\frac{u}{4})}.$$

Using the limiting expressions above, the four-point function simplifies as

$$C' \times \int_{v_{4B}} dv_4 \left(\frac{1}{m_4 - m_1 v_4}\right)^{2-\frac{(\vec{p}_1 - \vec{p}_4)^2}{2}}, \quad (4.4)$$

where again C' is a constant. Thus, in $v_4 \rightarrow v_{4B}$ limit, four-point function will diverge whenever the exponent satisfies the condition

$$2 - \frac{(\vec{p}_1 - \vec{p}_4)^2}{2} \geq 1 \iff (\vec{p}_1 - \vec{p}_4)^2 \leq 2.$$

Now for the kinematic configuration in Case-2, we have three boundaries,⁷ which are $v_4 = v_{4+}$, $v_4 = v_{4B}$, and $v_4 = +\infty$. The integral I_3 exists in the interval $(v_{4+}, +\infty)$, and the integral I_1 exists in the interval (v_{4+}, v_{4B}) . Using similar analysis as for Case-1, we find that I_3 diverges in the same way as for the Case-1 in $v_4 \rightarrow \infty$ limit, that is, whenever $(\vec{p}_1 - \vec{p}_3)^2 \leq 2$. For *generic* values of the kinematic parameters in Case-2 configuration, the Mandelstam invariants are finite in the $v_4 \rightarrow v_{4+}$ limit and so both I_1 and I_3 do not diverge. Though, I_1 diverges as $v_4 \rightarrow v_{4B}$ whenever $(\vec{p}_1 - \vec{p}_4)^2 \leq 2$.

4.2 Type-2: poles at boundary

In the previous subsection, we showed that some Mandelstam invariants approach finite values near the boundaries of the integral, for instance, $t \rightarrow -(\vec{p}_1 - \vec{p}_3)^2$ as $v_4 \rightarrow \infty$, and $u \rightarrow -(\vec{p}_1 - \vec{p}_4)^2$ as $v_4 \rightarrow v_{4B}$ for the kinematic configuration in Case-1. If corresponding to these finite values, the arguments of gamma functions are non-negative integers, we would have poles at the boundary of the integral. Integration to such poles would not be finite as they are only being approached from one direction in the sense of CPV.

For both Case-1 & 2 at $v_4 = v_{4B}$ we have $u = -(\vec{p}_1 - \vec{p}_4)^2$ and as $v_4 \rightarrow \infty$ we have $t \rightarrow -(\vec{p}_1 - \vec{p}_3)^2$, respectively. Corresponding to these values of t and u , we would have a boundary pole at $v_4 = v_{4B}$ whenever $(\vec{p}_1 - \vec{p}_4)^2 = 4$ or at $v_4 \rightarrow \infty$ whenever $(\vec{p}_1 - \vec{p}_3)^2 = 4$.⁸

One potential divergence that can arise in the scattering amplitude comes from the

$$\frac{1}{|v_2^2 - v_3^2|}$$

factor in the integrand at the lower limit of integration, v_{4+} . But the leading behavior of this factor goes as $\sim \frac{1}{(v_4 - v_{4+})^{1/2}}$ for *generic* v_{4+} and so the integral is finite at the lower limit of integration. The rest of the amplitude integrand goes to a constant as $v_4 \rightarrow v_{4+}$ (except at isolated values of the kinematic parameters which we will discuss momentarily). But when $v_{4+} = 1$, which corresponds to special choice of parameters, we will see in the next subsection that there is an extra divergence that emerges, which was noted in [1].

In Case-2, the Mandelstam invariants are finite at $v_4 = v_{4+}$ and can be calculated to be

$$s(v_{4+}) \equiv -\frac{m_2(m_1^2 + m_2^2 + m_3^2 - m_4^2 - 2m_2m_3)}{m_2 - m_3} - 2P_{12} - 8, \quad (4.5)$$

$$t(v_{4+}) \equiv -\frac{m_3(m_1^2 + m_2^2 + m_3^2 - m_4^2 - 2m_2m_3)}{m_3 - m_2} + 2P_{13} - 8, \quad (4.6)$$

$$u(v_{4+}) \equiv -m_1^2 + m_2^2 + m_3^2 - m_4^2 - 2m_2m_3 + 2P_{14} - 8. \quad (4.7)$$

⁷The following discussion assumes $m_1 + m_2 < m_3 + m_4$. When $m_1 + m_2 > m_3 + m_4$, essentially the same discussion goes through once we replace v_{4+} with 1.

⁸This boundary pole for $v_4 \rightarrow \infty$ could also be thought of as a boundary pole at $v_4 = 0$ if we think of it as a divergence in the integral I_1 instead of I_3 . Such translations about the origin of the divergence exists in other cases as well, but we have chosen not to emphasize them.

Figure 4. The plots of the branches of v_2 and v_3 . Only positive values are relevant, so we plot only them. The values of the kinematic parameters used for making the plot are $m_1 = 5.0$, $m_2 = 9.5$, $m_3 = 5.5$, $m_4 = 10.0$, $P_{12} = 39.5$, $P_{13} = 4.0$.

If these values correspond to poles of the gamma function (i.e., when these expressions equal $4(n - 1)$ for some non-negative integer n), naively we could expect boundary divergences at $v_4 = v_{4+}$. But there is a subtlety involved here. This is because the integral in Case-2 involves two pieces (3.13) and they contribute destructively at the boundary. One way to see this is to note that the two integrals can locally (around v_{4+}) be written as integrals over v_2 instead of as integrals over v_4 , by a change of variables.⁹ The point v_{4+} is a branch point for v_2 where the two branches v_{2u} and v_{2d} meet, as can be seen from figure 4. A basic observation is that the integrands i_1 and i_3 are the same function, but over the two branches.

Now consider a situation when there is a pole at the boundary arising from (4.5), i.e., in the gamma function associated to s (or equivalently, v_2). The situation when there is a pole in t , namely (4.6), is entirely analogous.¹⁰ The potential divergences in i_1 and i_3 together can be written as

$$\int_{v_{4+}} \frac{dv_4}{\sqrt{v_4 - v_{4+}}} \tilde{\Gamma}(s(v_{2u}(v_4))) + \int_{v_{4+}} \frac{dv_4}{\sqrt{v_4 - v_{4+}}} \tilde{\Gamma}(s(v_{2d}(v_4))). \quad (4.8)$$

⁹A related observation is that the tangent of s (or v_2) at v_{4+} is vertical. This means that finding analytic estimates of the integral via Taylor expansion in v_4 around v_{4+} is not feasible, even if the integral were perfectly well-defined and finite. So to apply the type of logic that we used at the beginning of section 3.2 based on the gamma function and its poles, one needs to first go over to a more convenient variable around v_{4+} . This is precisely what v_2 is.

¹⁰The u -channel case (4.7) is different, and we will discuss it towards the end of this subsection.

We have suppressed the rest of the functions in the integrand because they are well-defined and continuous across the branches and just go along for the ride, and we are using the notation $\tilde{\Gamma}(s)$ to mean $\Gamma(-1 - s/4)$ for brevity. As discussed, v_2 is a better variable to work with around v_{4+} so the integral becomes

$$\int_{v_2(v_{4+})} \frac{dv_{2u}}{\sqrt{v_4 - v_{4+}}} \frac{\tilde{\Gamma}(s(v_{2u}))}{dv_{2u}/dv_4} + \int_{v_2(v_{4+})} \frac{dv_{2d}}{\sqrt{v_4 - v_{4+}}} \frac{\tilde{\Gamma}(s(v_{2d}))}{dv_{2d}/dv_4}. \quad (4.9)$$

Now the crucial observation is that $1/(dv_2/dv_4)$ is continuous across the branches, but goes *through* zero at v_{4+} and cancels the factor $1/\sqrt{v_4 - v_{4+}}$ (up to a minus sign on the v_{2u} branch). The whole expression can be written as an integral along the $v_2(v_4)$ curve stretching across both branches, the portion near v_{4+} being proportional to

$$\int dv_2 \tilde{\Gamma}(s(v_2)), \quad (4.10)$$

where now the integral is in a small neighborhood around the point $v_2(v_{4+})$. There is a pole at that point, but this expression is bound to be finite in the sense of CPV by the discussions of the previous sections. That s as a function of v_2 does not have a maximum/minimum at $v_2(v_{4+})$ is readily checked.¹¹ The conclusion of all this is that the potential divergences that could have arisen from (4.5) and (4.6) when they correspond to gamma function poles are actually spurious, and in fact there are no divergences from them (unless the Mandelstam invariants have a maximum/minimum at v_{4+} as discussed in the footnote).

The conclusion of all this is that the potential divergences that could have arisen from (4.5) and (4.6) when they correspond to gamma function poles are actually spurious, and in fact there are no divergences from them.

The same is not however true about the poles corresponding to (4.7). One can check that there are indeed divergences in the integral that arise when (4.7) is equal to $4(n-1)$, with n a non-negative integer. This condition can in fact be written in a suggestive form as

$$(m_2 - m_3)^2 - (\vec{p}_2 - \vec{p}_3)^2 = 4(n-1). \quad (4.13)$$

This suggests that this divergence can be understood in terms of intermediate tower of string states going on-shell and is therefore a “good” divergence unrelated to the singu-

¹¹An exception to this is when we have a maximum/minimum of s or t at v_{4+} . This corresponds to the condition $(m_3 + m_4) = (m_1 + m_2)$ which coincides with $v_{4+} = 1$ and gives rise to divergences. In such a situation, the expression (4.5) for gamma function poles for a non-negative integer argument n happens when

$$s = (m_1 + m_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 = 4(n-1). \quad (4.11)$$

Similar arguments apply for t which reduces (4.6) to

$$t = (m_1 - m_3)^2 - (\vec{p}_1 - \vec{p}_3)^2 = 4(n-1). \quad (4.12)$$

Satisfyingly, both these expressions can be understood as special cases (for s and t) of the extrema poles we discuss in the next subsection when the extremum happens at the boundary.

larity.¹² To argue further for the infrared character of these divergences, one can replace the vertex operators (2.1) by their asymptotic behavior for large X^+X^- (far away from the singularity), given above (A.6) of [1], and act with $\square$ on $\psi_3^*\psi_2$. The result for large X^+X^- is that $\square = -(m_2 - m_3)^2 + (\vec{p}_2 - \vec{p}_3)^2$, in agreement with (4.13).

A thing to note about these boundary pole divergences is that the conditions above for them to arise are constraints on the kinematic parameters. This means that they are not generic.

4.3 Type-3: poles at Mandelstam maxima/minima

As we discussed earlier, whenever there is a gamma function pole in v_4 space at a maximum/minimum of a Mandelstam invariant, the integration through the pole would cause a divergence in the four-point function.

The discussion of the extrema splits into two possibilities depending on whether $(m_3 + m_4) > (m_1 + m_2)$ or $(m_3 + m_4) < (m_1 + m_2)$. For extrema of s with $(m_3 + m_4) > (m_1 + m_2)$, the divergences appear from poles in the integration range when $dv_2/dv_4 = 0$. In this situation, the expression $-1 - \frac{s}{4} = -n$ for non-negative integer n , reduces to the suggestive form

$$s = (m_4 + m_3)^2 - (\vec{p}_4 + \vec{p}_3)^2 = 4(n - 1). \quad (4.16)$$

On the other hand when $(m_3 + m_4) < (m_1 + m_2)$, the relevant maxima/minima of s arise from $ds/dv_2 = 0$ (i.e. $v_2 = 1$), which again reduces to a simple form

$$s = (m_1 + m_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 = 4(n - 1) \quad (4.17)$$

where n is some non-negative integer. It should be noted that the two conditions above are same whenever $(m_3 + m_4) = (m_1 + m_2)$, occurring at $v_4 = 1$.

A similar conclusion holds for t . Here also, condition $(m_3 + m_4) > (m_1 + m_2)$ means poles come from $dv_3/dv_4 = 0$. The fact that these extrema are at non-negative integers n gives rise to

$$t = (m_2 - m_4)^2 - (\vec{p}_2 - \vec{p}_4)^2 = 4(n - 1). \quad (4.18)$$

An extra caveat is that these poles lie in the integration range only when $m_2 > m_4$ (which automatically forces $m_3 > m_1$). For $(m_3 + m_4) < (m_1 + m_2)$, we have extrema of t when $dt/dv_3 = 0$ (i.e. $v_3 = 1$) giving us the condition on t ,

$$t = (m_1 - m_3)^2 - (\vec{p}_1 - \vec{p}_3)^2 = 4(n - 1). \quad (4.19)$$

¹²It is possible to show that (4.13) has solutions in kinematic parameters that satisfy all the constraints for arbitrarily large n . An explicit way to see this is to choose an α (where $0 < \alpha < 1/\sqrt{2}$) that solves

$$n = \frac{(1 - 8\alpha^2 + 2\alpha^3 + 4\alpha^4 + 4\alpha^5 + \alpha^6)}{4\alpha^2} \quad (4.14)$$

and set

$$m_1 = \alpha, \quad m_4 = \frac{1}{\alpha}, \quad m_2 = \frac{1}{\alpha} - \alpha, \quad m_3 = \alpha(1 + \alpha), \quad P_{12} = \frac{1}{2\alpha^2}, \quad P_{13} = \frac{\alpha^2}{2} + 4 \quad (4.15)$$

It is easily checked that all the consistency conditions between the kinematic parameters are satisfied.

These t -poles are in the integration range only when $m_3 > m_1$ (which automatically forces $m_2 > m_4$). Again, both the former conditions are equivalent for $(m_3 + m_4) = (m_1 + m_2)$.

It is worth remarking that the extrema of u occur only at $v_4 = 1$, and this point is part of the integration range when $(m_3 + m_4) \leq (m_1 + m_2)$. Poles at these extrema will occur whenever u at $v_4 = 1$ satisfies

$$u = (m_1 - m_4)^2 - (\vec{p}_1 - \vec{p}_4)^2 = 4(n - 1) \quad (4.20)$$

for non-negative integer n . For convenience of comparison of the various towers of divergent poles that we have uncovered, we repeat also (4.13) here:

$$u = (m_2 - m_3)^2 - (\vec{p}_2 - \vec{p}_3)^2 = 4(n - 1). \quad (4.21)$$

Note that these poles are not extrema poles, and exist only when $(m_3 + m_4) \geq (m_1 + m_2)$ which is the necessary condition for boundary poles to exist.

Now we see that we have towers of divergent poles for each channel. The arguments for the existence of the various poles is not manifestly symmetric under interchange of labels because we chose to do our integration along v_4 , which breaks the symmetry between v_3 and v_4 , and also because we are looking at various cases separately (in particular, note that the cases we are explicitly considering have $m_4 \geq m_3$ and exchanging t and u corresponds to exchanging m_4 and m_3). The origin of a divergence as a boundary/extremum divergence is not independent of these choices and is merely an artifact. Another sanity check is that when the parameter condition $(m_3 + m_4) = (m_1 + m_2)$ is satisfied, it is straightforward to see that the extrema divergences become the boundary divergences of the last section because $v_{4+} = 1$ in this case.

These divergences that we have identified are expected IR divergences corresponding to intermediate string states going on-shell. We note that these divergences are non-generic (as in, they appear only when the parameters satisfy the conditions listed in this subsection). But it is possible to show that solutions exist for all n (non-negative). This is somewhat non-trivial to demonstrate in general because the kinematic parameters have to satisfy various constraints as well as consistency conditions between them. For the case of the s -poles (4.16), for example, once one finds a solution (by trial and error - this can be easily accomplished with Mathematica) for a small integer n that satisfies the constraints, one can increase m_4 as one wishes while holding the other parameters fixed in order to satisfy (4.16): it is easy to check that all the kinematic constraints will still be satisfied. Various arguments of a similar flavor can also be used for (4.17)–(4.21) as well¹³ to show that these expressions correspond to infinite towers of string states going on-shell.

¹³A specific ansatz for u -poles was given in a previous footnote. The t -poles (4.18) are also somewhat subtle, so we present an explicit ansatz here that finds solutions of (4.18) for all non-negative integers. Choose an ϵ (where $0 < \epsilon \leq \frac{1}{4}$) that solves

$$n = -1 + \frac{1}{16\epsilon^2} \quad (4.22)$$

and set

$$m_1 = \epsilon, \quad m_4 = \frac{1}{\epsilon^2}, \quad m_2 = \frac{1}{\epsilon^2} + \frac{1}{2\epsilon}, \quad m_3 = \frac{1}{\epsilon}, \quad P_{12} = \frac{1}{2\epsilon^2}, \quad P_{13} = \frac{\epsilon^2}{2} + \frac{1}{2\epsilon^2} \quad (4.23)$$

It is easily checked that all the consistency conditions for the kinematic parameters are satisfied.

4.4 Type-4: IR divergences identified in [1]

For the Case-2 configuration, we find that $v_{4+} = v_{4-} = 1$ whenever $(m_1 + m_2) = (m_3 + m_4)$ and this gives rise to one more divergence.¹⁴ As $v_4 \rightarrow 1$, both $v_{2u}, v_{3u} \rightarrow 1$ and all the Mandelstam invariants are finite. The four-point function integral takes the form

$$C'' \times \int_{v_4=1} dv_4 \frac{1}{(v_{2u} - v_{3u})}, \quad (4.24)$$

where C'' is a constant. This integral above diverges logarithmically in the $v_4 \rightarrow 1$ limit. This divergence is not associated with the singularity but is an infra-red divergence [1].

It is important to note that the condition $v_{4+} = 1$ is crucial for the existence of this divergence: $v_{4+} = 1$ forces $v_{4+} = v_{4-}$ and this results in Δ in the definition of v_2 and v_3 contributing to the leading behavior. This makes the integrand near the lower limit to behave as $1/(v_4 - v_{4+})$ instead of $1/(v_4 - v_{4+})^{1/2}$. Crucially, this means that the divergence again has an interpretation as an isolated IR divergence [1].

An integral can diverge only from one of its boundaries or from a region in the integration range where the integrand blows up badly enough. We have checked every such possibility by looking at the boundaries, the poles of the gamma functions and the divergence due to $1/(v_{2u,d} - v_{3u,d})$. This means that our scan of the divergences is an exhaustive one.

5 Summary of divergences

In this section, we summarize the divergences in the tree-level 2-to-2 string scattering amplitude. We will only list the divergences in the context of the two cases we have considered. There exist analogous divergences that arise when the ordering of the mass parameters are changed, but these can all be obtained from the cases we list here by appropriate permutations of the labels.

- UV divergences that arise when $(\vec{p}_1 - \vec{p}_3)^2 \leq 2$ or when $(\vec{p}_1 - \vec{p}_4)^2 \leq 2$.¹⁵
- IR divergences that arise when one of the conditions (4.16)–(4.21) is satisfied, arising from the tower of string states going on-shell.
- IR divergences arising from tachyons going on-shell when $(\vec{p}_1 - \vec{p}_3)^2 = 4$ or when $(\vec{p}_1 - \vec{p}_4)^2 = 4$.
- The logarithmic IR divergence that arises when $(m_1 + m_2) = (m_3 + m_4)$.

The first and last kinds of divergences were already noticed in [1]. Our claim here is that all the other IR divergences are physical divergences expected to be present in healthy scattering amplitudes. The UV divergences disappear in the large (dimensionless) α' region of the parameter space, as already belabored in the introduction.

¹⁴This divergence was noticed in [1]. See their discussion right before section 4. The various sign choices arising in their $(m_4 - m_1)^2 = (m_2 - m_3)^2$ condition are distributed over the various cases here, which are analogous to this one. So we do not discuss them.

¹⁵When one re-instates the α' these will read $\alpha'(\vec{p}_1 - \vec{p}_3)^2 \leq 2$ and $\alpha'(\vec{p}_1 - \vec{p}_4)^2 \leq 2$.

Acknowledgments

We thank Pallab Basu for managing to feed one of the authors for the last few months, despite Indian bureaucracy. CK thanks Joan Simon for discussions on a related collaboration. This work was supported in part by the Belgian Federal Science Policy Office through the Interuniversity Attraction Pole P7/37, by FWO-Vlaanderen through project G020714N, and by the Vrije Universiteit Brussel through the Strategic Research Program “High-Energy Physics”.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M. Berkooz, B. Craps, D. Kutasov and G. Rajesh, *Comments on cosmological singularities in string theory*, *JHEP* **03** (2003) 031 [[hep-th/0212215](#)] [[INSPIRE](#)].
- [2] H. Liu, G.W. Moore and N. Seiberg, *The challenging cosmic singularity*, [gr-qc/0301001](#) [[INSPIRE](#)].
- [3] L. Cornalba and M.S. Costa, *Time dependent orbifolds and string cosmology*, *Fortschr. Phys.* **52** (2004) 145 [[hep-th/0310099](#)] [[INSPIRE](#)].
- [4] B. Durin and B. Pioline, *Closed strings in Misner space: a toy model for a big bounce?*, [hep-th/0501145](#) [[INSPIRE](#)].
- [5] B. Craps, *Big bang models in string theory*, *Class. Quant. Grav.* **23** (2006) S849 [[hep-th/0605199](#)] [[INSPIRE](#)].
- [6] M. Berkooz and D. Reichmann, *A short review of time dependent solutions and space-like singularities in string theory*, *Nucl. Phys. Proc. Suppl.* **171** (2007) 69 [[arXiv:0705.2146](#)] [[INSPIRE](#)].
- [7] H. Liu, G.W. Moore and N. Seiberg, *Strings in a time dependent orbifold*, *JHEP* **06** (2002) 045 [[hep-th/0204168](#)] [[INSPIRE](#)].
- [8] A. Lawrence, *On the instability of 3-D null singularities*, *JHEP* **11** (2002) 019 [[hep-th/0205288](#)] [[INSPIRE](#)].
- [9] G.T. Horowitz and J. Polchinski, *Instability of space-like and null orbifold singularities*, *Phys. Rev. D* **66** (2002) 103512 [[hep-th/0206228](#)] [[INSPIRE](#)].
- [10] C. Krishnan and S. Roy, *Higher spin resolution of a toy big bang*, *Phys. Rev. D* **88** (2013) 044049 [[arXiv:1305.1277](#)] [[INSPIRE](#)].
- [11] C. Krishnan and S. Roy, *Desingularization of the Milne universe*, [arXiv:1311.7315](#) [[INSPIRE](#)].
- [12] M.R. Gaberdiel and R. Gopakumar, *An AdS_3 dual for minimal model CFTs*, *Phys. Rev. D* **83** (2011) 066007 [[arXiv:1011.2986](#)] [[INSPIRE](#)].
- [13] A. Campoleoni, S. Fredenhagen, S. Pfenninger and S. Theisen, *Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields*, *JHEP* **11** (2010) 007 [[arXiv:1008.4744](#)] [[INSPIRE](#)].

- [14] C. Krishnan, A. Raju and S. Roy, *A Grassmann path from AdS_3 to flat space*, *JHEP* **03** (2014) 036 [[arXiv:1312.2941](#)] [[INSPIRE](#)].
- [15] H. Afshar, A. Bagchi, R. Fareghbal, D. Grumiller and J. Rosseel, *Spin-3 gravity in three-dimensional flat space*, *Phys. Rev. Lett.* **111** (2013) 121603 [[arXiv:1307.4768](#)] [[INSPIRE](#)].
- [16] H.A. Gonzalez, J. Matulich, M. Pino and R. Troncoso, *Asymptotically flat spacetimes in three-dimensional higher spin gravity*, *JHEP* **09** (2013) 016 [[arXiv:1307.5651](#)] [[INSPIRE](#)].
- [17] X. Bekaert, S. Cnockaert, C. Iazeolla and M.A. Vasiliev, *Nonlinear higher spin theories in various dimensions*, [hep-th/0503128](#) [[INSPIRE](#)].
- [18] B. Sundborg, *Stringy gravity, interacting tensionless strings and massless higher spins*, *Nucl. Phys. Proc. Suppl.* **102** (2001) 113 [[hep-th/0103247](#)] [[INSPIRE](#)].
- [19] E. Witten, *Spacetime reconstruction*, talk given at the *J.H. Schwarz 60th Birthday Conference*, California Institute of Technology, Pasadena U.S.A., 3–4 Nov. 2001.
- [20] S. Giombi et al., *Chern-Simons theory with vector fermion matter*, *Eur. Phys. J. C* **72** (2012) 2112 [[arXiv:1110.4386](#)] [[INSPIRE](#)].
- [21] C.-M. Chang, S. Minwalla, T. Sharma and X. Yin, *ABJ triality: from higher spin fields to strings*, *J. Phys. A* **46** (2013) 214009 [[arXiv:1207.4485](#)] [[INSPIRE](#)].
- [22] D.J. Gross and P.F. Mende, *The high-energy behavior of string scattering amplitudes*, *Phys. Lett. B* **197** (1987) 129 [[INSPIRE](#)].