RECEIVED: October 4, 2017

ACCEPTED: October 5, 2017

PUBLISHED: October 16, 2017

Erratum: Non-global structure of the $\mathcal{O}(\alpha_s^2)$ dijet soft function

Andrew Hornig,^a Christopher Lee,^b Iain W. Stewart,^{b,c} Jonathan R. Walsh^{d,e}
and Saba Zuberi^{d,e}

^aDepartment of Physics, University of Washington,
Box 351560, Seattle, WA 98195, U.S.A.

^bCenter for Theoretical Physics, Massachusetts Institute of Technology, 77 Massachusetts Ave.,
Cambridge, MA 02139, U.S.A.

^cCenter for the Fundamental Laws of Nature, Harvard University, 17 Oxford St.,
Cambridge, MA 02138, U.S.A.

^dTheoretical Physics Group, Ernest Orlando Lawrence Berkeley National Laboratory,
1 Cyclotron Rd., 366 LeConte Hall #7300, Berkeley, CA 94720, U.S.A.

^eCenter for Theoretical Physics, University of California,
366 LeConte Hall #7300, Berkeley, CA 94720, U.S.A.

E-mail: ahornig@uw.edu, cleel37@mit.edu, iains@mit.edu, jwalsh@lbl.gov,
szuberi@lbl.gov

ERRATUM TO: [JHEP08\(2011\)054](#)

ARXIV EPRINT: [1105.4628](#)

The position space results in eqs. (3.30)–(3.32) of ref. [1] are correct, but there are typos in the coefficients extracted in the $x_1 \gg x_2$ limit of $t_2(x_1/x_2)$, so eq. (3.33) should read:

$$s_2^{[2]} = -\frac{2\pi^2}{3}C_F C_A, \quad s_2^{[1]} = 2 \left[C_F C_A \frac{(11\pi^2 - 3 - 18\zeta_3)}{9} + C_F T_R n_f \left(\frac{6 - 4\pi^2}{9} \right) \right], \quad (3.33)$$

$$s_2^{[0]} = -s_2^{[1]} \ln 2 - 4C_F C_A F_N(1) - 4C_F T_R n_f F_Q(1) + C_F C_A s_2^{[C_F C_A]} + C_F T_R n_f s_2^{[n_f]}.$$

Similarly, from taking the large $\ell_1 \gg \ell_2$ limit of the momentum space result in eq. (3.36), the coefficients in eq. (3.39) should read:

$$s_{2c}^{[0]} = -s_{2c}^{[1]} \ln 2 - 4C_F C_A f_N(1) - 4C_F T_R n_f f_Q(1) + C_F C_A s_{2\rho}^{[C_F C_A]} + C_F T_R n_f s_{2\rho}^{[n_f]}. \quad (3.39)$$

Finally, there are constant terms that should be added to the μ -dependent part of the momentum-space soft function, which appear from the conversion of logarithms from position to momentum space, so eq. (3.43) should read:

$$\begin{aligned}
\mathcal{R}_c(\ell_1^c, \ell_2^c, \mu) = & -\frac{\alpha_s(\mu)C_F}{\pi} \left(L_1^2 + L_2^2 - \frac{\pi^2}{3} \right) \\
& + \frac{\alpha_s^2(\mu)}{(4\pi)^2} \left\{ 8C_F^2 (L_1^2 + L_2^2)^2 + \left(\frac{88}{9}C_FC_A - \frac{32}{9}C_FT_Rn_f \right) (L_1^3 + L_2^3) \right. \\
& + \left[-\frac{20\pi^2}{3}C_F^2 + C_FC_A \left(\frac{4\pi^2}{3} - \frac{268}{9} \right) + \frac{80}{9}C_FT_Rn_f \right] (L_1^2 + L_2^2) \\
& + \left[64\zeta_3 C_F^2 + C_FC_A \left(\frac{808}{27} - \frac{22\pi^2}{9} - 28\zeta_3 \right) \right. \\
& \left. \left. - C_FT_Rn_f \left(\frac{224}{27} - \frac{8\pi^2}{9} \right) \right] (L_1 + L_2) - C_F^2 \frac{28\pi^4}{45} \right. \\
& \left. + C_FC_A \left(\frac{352\zeta_3}{9} + \frac{268\pi^2}{27} - \frac{4\pi^4}{9} \right) - C_FT_Rn_f \left(\frac{128\zeta_3}{9} + \frac{80\pi^2}{27} \right) \right\}. \quad (3.43)
\end{aligned}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Hornig, C. Lee, I. W. Stewart, J. R. Walsh and S. Zuberi, *Non-global structure of the $O(\alpha_s^2)$ dijet soft function*, *JHEP* **08** (2011) 054 [[arXiv:1105.4628](https://arxiv.org/abs/1105.4628)].