

RECEIVED: August 26, 2016

ACCEPTED: November 4, 2016

PUBLISHED: November 11, 2016

# Bethe ansatz solutions of the $\tau_2$ -model with arbitrary boundary fields

Xiaotian Xu,<sup>a,b</sup> Kun Hao,<sup>a,b</sup> Tao Yang,<sup>a,b,1</sup> Junpeng Cao,<sup>c,d,e</sup> Wen-Li Yang<sup>a,b,f</sup> and Kangjie Shi<sup>a,b</sup>

<sup>a</sup>*Institute of Modern Physics, Northwest University,  
Xian 710069, China*

<sup>b</sup>*Shaanxi Key Laboratory for Theoretical Physics Frontiers,  
Xian 710069, China*

<sup>c</sup>*Beijing National Laboratory for Condensed Matter Physics,  
Institute of Physics, Chinese Academy of Sciences,  
Beijing 100190, China*

<sup>d</sup>*Collaborative Innovation Center of Quantum Matter,  
Beijing, China*

<sup>e</sup>*School of Physical Sciences, University of Chinese Academy of Sciences,  
Beijing, China*

<sup>f</sup>*Beijing Center for Mathematics and Information Interdisciplinary Sciences,  
Beijing, 100048, China*

E-mail: [xxt0206@stumail.nwu.edu.cn](mailto:xxt0206@stumail.nwu.edu.cn), [haoke72@163.com](mailto:haoke72@163.com), [yangt@nwu.edu.cn](mailto:yangt@nwu.edu.cn),  
[junpengcao@iphy.ac.cn](mailto:junpengcao@iphy.ac.cn), [wlyang@nwu.edu.cn](mailto:wlyang@nwu.edu.cn), [kjshi@nwu.edu.cn](mailto:kjshi@nwu.edu.cn)

**ABSTRACT:** The quantum  $\tau_2$ -model with generic site-dependent inhomogeneity and arbitrary boundary fields is studied via the off-diagonal Bethe Ansatz method. The eigenvalues of the corresponding transfer matrix are given in terms of an inhomogeneous  $T - Q$  relation, which is based on the operator product identities among the fused transfer matrices and the asymptotic behavior of the transfer matrices. Moreover, the associated Bethe Ansatz equations are also obtained.

**KEYWORDS:** Bethe Ansatz, Lattice Integrable Models

ARXIV EPRINT: [1608.06209](https://arxiv.org/abs/1608.06209)

<sup>1</sup>Corresponding author.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Transfer matrix</b>	<b>2</b>
<b>3</b>	<b>Properties of the transfer matrix</b>	<b>5</b>
3.1	Asymptotic behaviors and average values	5
3.2	Fusion hierarchy	7
<b>4</b>	<b>Truncation identity</b>	<b>10</b>
<b>5</b>	<b>Eigenvalues of the fundamental transfer matrix</b>	<b>12</b>
5.1	Functional relations of eigenvalues	12
5.2	T-Q relation	14
5.2.1	Generic case	14
5.2.2	Degenerate case	15
<b>6</b>	<b>Conclusion</b>	<b>16</b>
<b>A</b>	<b>Specific cases of the fused <math>K</math>-matrices</b>	<b>17</b>
<b>B</b>	<b>Explicit expression of the average value functions</b>	<b>17</b>

---

## 1 Introduction

The finite-size inhomogeneous  $\tau_2$ -model also known as the Baxter-Bazhanov-Stroganov model (BBS model) [1–4] is a  $N$ -state spin lattice model, which is intimately related to some other integrable models under certain parameter constraints such as the chiral Potts model [5–10] and the relativistic quantum Toda chain model [11]. Lots of papers have appeared to explain such connections and many efforts have been made to calculate the eigenvalues of the chiral Potts model by solving the  $\tau_2$ -model with a recursive functional relation [4, 12–14]. The  $\tau_2$ -model is a simple quantum integrable models associated with cyclic representation of the Wely algebra. Although its integrability has been proven [3] for decades, there is still no effective method to solve the model completely due to lack of a simple  $Q$ -operator solution in terms of Baxter’s  $T - Q$  relation. In fact, the  $Q$ -operator is a very complicated function defined in high genus space and its concrete expression is hard to be derived. Very recently, Paul Fendley had found a “parafermionic” way to diagonalise a simple solvable Hamiltonian associated with the chiral Potts model [15]. Subsequently, this method was generalised to solve the  $\tau_2$ -model with particular open boundaries [16–18].

Until very recently, the Bethe Ansatz solution of the periodic  $\tau_2$ -model with generic site-dependent inhomogeneity was obtained by constructing an inhomogeneous  $T - Q$  relation with polynomial  $Q$ -functions (i.e. off-diagonal Bethe Ansatz method (ODBA)) [19–24]), which provides a perspective to investigate the  $\tau_2$ -model with generic open boundary condition. Moreover, such a solution allows the authors [25] further to retrieve the corresponding Bethe states.

The aim of this paper is to explicitly construct the eigenvalues of the transfer matrix for the open  $\tau_2$ -model with the most generic inhomogeneity, where we generalise the ODBA method to solve the open inhomogeneous  $\tau_2$ -model with arbitrary integrable open boundary condition in combination with the fusion technique. By introducing an off-diagonal term in the conventional  $T - Q$  relation (i.e., the inhomogeneous  $T - Q$  relation), we obtain the spectrum of the generic open  $\tau_2$ -model and the associated Bethe Ansatz equations.

The outline of this paper is as follows. In section 2, we begin with a brief introduction of the fundamental transfer matrix. In section 3, we study the properties of the transfer matrix and employing the so-called fusion procedure [26–28] to construct the higher-spin transfer matrices, which obey an infinite fusion hierarchy. In section 4, we obtain the truncation identity for the fused transfer matrices when the bulk anisotropy value takes the special case  $\eta = \frac{2i\pi}{p}$  and the exact functional relations of the fundamental transfer matrix. In section 5, we give the eigenvalues of the transfer matrix in terms of some inhomogeneous  $T - Q$  relation and the associated Bethe Ansatz equation. In the last section, we summarize our results and give some discussions. Some detailed technical calculations are given in appendices A and B.

## 2 Transfer matrix

Let us fix an odd integer  $p$  such that  $p \geq 3$ , and let  $\mathbf{V}$  be a  $p$ -dimensional vector space (i.e. the local Hilbert space) with an orthonormal basis  $\{|m\rangle |m \in \mathbb{Z}_p\}$ . Define two  $p \times p$  matrices  $X$  and  $Z$  which act on the basis as

$$X|m\rangle = q^m|m\rangle, \quad Z|m\rangle = |m+1\rangle, \quad m \in \mathbb{Z}_p, \quad (2.1)$$

where  $q \equiv e^{-\eta}$  is a  $p$ -root of unity (i.e.,  $q^p = 1$ ). The embedding operators  $\{X_n, Z_n | n = 1, \dots, N\}$  denote the generators of the ultra-local Weyl algebra:

$$X_n Z_m = q^{\delta_{nm}} Z_m X_n, \quad X_n^p = Z_n^p = 1, \quad \forall n, m \in \{1, \dots, N\}. \quad (2.2)$$

It has been shown that the  $\tau_2$ -model can be described by a quantum integrable spin chain [3]. In order to construct the monodromy matrix, one need to introduce the  $L$ -operators for each site of the quantum chain. The associated  $L$ -operator  $L_n(u) \in \text{End}(\mathbf{C}^2 \otimes \mathbf{V})$  defined in the most general cyclic representation of  $U_q(sl_2)$ , is given by [3]

$$\begin{aligned} L_n(u) &= \begin{pmatrix} e^u d_n^{(+)} X_n + e^{-u} d_n^{(-)} X_n^{-1} & (g_n^{(+)} X_n^{-1} + g_n^{(-)} X_n) Z_n \\ (h_n^{(+)} X_n^{-1} + h_n^{(-)} X_n) Z_n^{-1} & e^u f_n^{(+)} X_n^{-1} + e^{-u} f_n^{(-)} X_n \end{pmatrix} \\ &= \begin{pmatrix} A_n(u) & B_n(u) \\ C_n(u) & D_n(u) \end{pmatrix}, \quad n = 1, \dots, N, \end{aligned} \quad (2.3)$$

where  $d_n^{(+)}, d_n^{(-)}, g_n^{(+)}, g_n^{(-)}, h_n^{(+)}, h_n^{(-)}, f_n^{(+)}$  and  $f_n^{(-)}$  are some parameters associated with each site. These parameters are subjected to two constraints,

$$g_n^{(-)}h_n^{(-)} = f_n^{(-)}d_n^{(+)}, \quad g_n^{(+)}h_n^{(+)} = f_n^{(+)}d_n^{(-)}, \quad n = 1, \dots, N, \quad (2.4)$$

which ensure that the above L-operator  $L_n$  satisfies the Yang-Baxter algebra [3],

$$R(u-v)(L_n(u) \otimes 1)(1 \otimes L_n(v)) = (1 \otimes L_n(v))(L_n(u) \otimes 1)R(u-v), \quad n = 1, \dots, N. \quad (2.5)$$

The associated  $R$ -matrix  $R(u) \in \text{End}(\mathbf{C}^2 \otimes \mathbf{C}^2)$  is the well-known six-vertex  $R$ -matrix given by

$$R(u) = \begin{pmatrix} \sinh(u+\eta) & 0 & 0 & 0 \\ 0 & \sinh u \sinh \eta & 0 & 0 \\ 0 & \sinh \eta \sinh u & 0 & 0 \\ 0 & 0 & 0 & \sinh(u+\eta) \end{pmatrix}, \quad (2.6)$$

with the crossing parameter  $\eta$  taking the special values

$$\eta = 2i\pi/p, \quad p = 2l+1, \quad l = 1, 2, \dots. \quad (2.7)$$

The R-matrix satisfies the quantum Yang-Baxter equation (QYBE) [29, 30] and becomes some projectors when the spectral parameter  $u$  takes some special values as

$$\text{Antisymmetric-fusion conditions : } R(-\eta) = -2 \sinh \eta P^{(-)}, \quad (2.8)$$

$$\text{Symmetric-fusion conditions : } R(\eta) = 2 \sinh \eta \text{Diag}(\cosh \eta, 1, 1, \cosh \eta) P^{(+)}, \quad (2.9)$$

where  $P^{(+)}$  ( $P^{(-)}$ ) is the symmetric (anti-symmetric) projector of the tensor space  $\mathbf{C}^2 \otimes \mathbf{C}^2$ . Associated with the local  $L$ -operators  $\{L_n(u) | n = 1, \dots, N\}$  given by (2.3), let us introduce the one-row monodromy matrix  $T(u)$

$$T(u) = \begin{pmatrix} \mathbf{A}(u) & \mathbf{B}(u) \\ \mathbf{C}(u) & \mathbf{D}(u) \end{pmatrix} = L_N(u) L_{N-1}(u) \cdots L_1(u). \quad (2.10)$$

The local relations (2.5) imply that the monodromy matrix  $T(u)$  also satisfies the Yang-Baxter algebra

$$R(u-v)(T(u) \otimes 1)(1 \otimes T(v)) = (1 \otimes T(v))(T(u) \otimes 1)R(u-v), \quad (2.11)$$

which ensures the integrability of the  $\tau_2$ -model with the periodic boundary condition [3].

Integrable open chain can be constructed as follows [31]. Let us introduce a pair of  $K$ -matrices  $K^-(u)$  and  $K^+(u)$ . The former satisfies the reflection equation (RE) [32]

$$\begin{aligned} & R_{12}(u_1 - u_2)K_1^-(u_1)R_{21}(u_1 + u_2)K_2^-(u_2) \\ &= K_2^-(u_2)R_{12}(u_1 + u_2)K_1^-(u_1)R_{21}(u_1 - u_2), \end{aligned} \quad (2.12)$$

and the latter satisfies the dual RE

$$\begin{aligned} & R_{12}(u_2 - u_1)K_1^+(u_1)R_{21}(-u_1 - u_2 - 2\eta)K_2^+(u_2) \\ &= K_2^+(u_2)R_{12}(-u_1 - u_2 - 2\eta)K_1^+(u_1)R_{21}(u_2 - u_1). \end{aligned} \quad (2.13)$$

In order to construct the associated open spin chain, let us introduce the  $\hat{L}(u)$  in the form of

$$\begin{aligned}\hat{L}_n(u) &= \begin{pmatrix} e^{-u-\eta} f_n^{(+)} X_n^{-1} + e^{u+\eta} f_n^{(-)} X_n & -(g_n^{(+)} X_n^{-1} + g_n^{(-)} X_n) Z_n \\ -(h_n^{(+)} X_n^{-1} + h_n^{(-)} X_n) Z_n^{-1} & e^{-u-\eta} d_n^{(+)} X_n + e^{u+\eta} d_n^{(-)} X_n^{-1} \end{pmatrix} \\ &= \begin{pmatrix} D_n(-u-\eta) & -B_n(-u-\eta) \\ -C_n(-u-\eta) & A_n(-u-\eta) \end{pmatrix}. \end{aligned}\quad (2.14)$$

It is easy to check that  $L_n(u)$  enjoys the crossing property

$$L_n(u) = \sigma^y \hat{L}_n^t(-u-\eta) \sigma^y, \quad n = 1, \dots, N, \quad (2.15)$$

and the inverse relation

$$L_n(u) \hat{L}_n(-u) = \text{Det}_q\{L_n(u)\} \times \text{id}, \quad n = 1, \dots, N,$$

where the function  $\text{Det}_q\{L_n(u)\}$  is the quantum determinant (which will be given by below (3.32)). Associated with the local  $L$ -operators  $\{\hat{L}_n(u)|n = 1, \dots, N\}$  given by (2.14), let us introduce another one-row monodromy matrix  $\hat{T}(u)$  (cf., (2.10))

$$\hat{T}(u) = \begin{pmatrix} \hat{\mathbf{A}}(u) & \hat{\mathbf{B}}(u) \\ \hat{\mathbf{C}}(u) & \hat{\mathbf{D}}(u) \end{pmatrix} = \hat{L}_1(u) \hat{L}_2(u) \cdots \hat{L}_N(u). \quad (2.16)$$

For the system with integrable open boundaries, the transfer matrix  $t(u)$  of the  $\tau_2$ -model with open boundaries can be constructed as [31]

$$t(u) = \text{tr}\{K^+(u)T(u)K^-(u)\hat{T}(u)\}, \quad (2.17)$$

where  $\text{tr}$  denotes trace over ‘auxiliary space’. The quadratic relation (2.11) and (dual) reflection equations (2.12) and (2.13) lead to the fact that the transfer matrix  $t(u)$  of the  $\tau_2$ -model with different spectral parameters are mutually commutative [31], i.e.,  $[t(u), t(v)] = 0$ , which ensures the integrability of the model by treating  $t(u)$  as the generating functional of the conserved quantities.

In this paper, we consider the most generic non-diagonal  $K^-(u)$  matrix found in refs. [33, 34], which is in the form of

$$K^-(u) = \begin{pmatrix} K_{11}^-(u) & K_{12}^-(u) \\ K_{21}^-(u) & K_{22}^-(u) \end{pmatrix}, \quad (2.18)$$

with

$$\begin{aligned}K_{11}^-(u) &= 2[\sinh(\alpha_-) \cosh(\beta_-) \cosh(u) + \cosh(\alpha_-) \sinh(\beta_-) \sinh(u)], \\ K_{22}^-(u) &= 2[\sinh(\alpha_-) \cosh(\beta_-) \cosh(u) - \cosh(\alpha_-) \sinh(\beta_-) \sinh(u)], \\ K_{12}^-(u) &= e^{\theta_-} \sinh(2u), \quad K_{21}^-(u) = e^{-\theta_-} \sinh(2u), \end{aligned}\quad (2.19)$$

where  $\alpha_-$ ,  $\beta_-$ , and  $\theta_-$  are three free boundary parameters. The most generic non-diagonal  $K$ -matrix  $K^+(u)$  is given by

$$K^+(u) = K^-(-u-\eta)|_{(\alpha_-, \beta_-, \theta_-) \rightarrow (-\alpha_+, -\beta_+, \theta_+)}.\quad (2.20)$$

We note that the two  $K$ -matrices possess the following properties

$$K^\mp(u + i\pi) = -\sigma^z K^\mp(u) \sigma^z, \quad (2.21)$$

$$K^-(0) = \frac{1}{2} \text{tr}(K^-(0)) \times \text{id}, \quad K^-\left(\frac{i\pi}{2}\right) = \frac{1}{2} \text{tr}\left(K^-\left(\frac{i\pi}{2}\right) \sigma^z\right) \times \sigma^z, \quad (2.22)$$

where  $\sigma^\alpha$  with  $\alpha = x, y, z$  are the Pauli matrices.

### 3 Properties of the transfer matrix

#### 3.1 Asymptotic behaviors and average values

Based on the explicit expressions (2.3) and (2.14) of the L-operators, the generic boundary matrices (2.19)–(2.20), and the definition (2.10)–(2.16) of the monodromy matrices, we note that the transfer matrix  $t(u)$  given by (2.17) has the asymptotic behavior,

$$\lim_{u \rightarrow \pm\infty} t(u) = -\frac{1}{4} e^{\pm\{(2N+4)u+(N+2)\eta\}} \left\{ e^{\theta_+ - \theta_-} F^{(+)} F^{(-)} + e^{-\theta_+ + \theta_-} D^{(+)} D^{(-)} \right\} \times \text{id}, \quad (3.1)$$

where  $D^{(\pm)}$  and  $F^{(\pm)}$  are four constants related to the inhomogeneous parameters as follows,

$$D^{(\pm)} = \prod_{n=1}^N d_n^{(\pm)}, \quad F^{(\pm)} = \prod_{n=1}^N f_n^{(\pm)}. \quad (3.2)$$

Moreover, we can calculate the special values of the associated transfer matrix at  $u = 0, \frac{i\pi}{2}$  with the help of the relations (2.22), namely,

$$\begin{aligned} t(0) &= -2^3 \sinh \alpha_- \cosh \beta_- \sinh \alpha_+ \cosh \beta_+ \cosh \eta \\ &\times \prod_{n=1}^N \left( e^{-\eta} d_n^{(+)} f_n^{(+)} + e^\eta d_n^{(-)} f_n^{(-)} - e^\eta g_n^{(+)} h_n^{(-)} - e^{-\eta} g_n^{(-)} h_n^{(+)} \right) \times \text{id}, \end{aligned} \quad (3.3)$$

$$\begin{aligned} t\left(\frac{i\pi}{2}\right) &= -2^3 \cosh \alpha_- \sinh \beta_- \cosh \alpha_+ \sinh \beta_+ \cosh \eta \\ &\times \prod_{l=1}^N \left( e^{-\eta} d_l^{(+)} f_l^{(+)} + e^\eta d_l^{(-)} f_l^{(-)} + e^\eta g_l^{(+)} h_l^{(-)} + e^{-\eta} g_l^{(-)} h_l^{(+)} \right) \times \text{id}. \end{aligned} \quad (3.4)$$

The expressions of the  $L$ -operators (2.3) and (2.14) allows us to derive their quasi-periodicities

$$L_n(u + i\pi) = -\sigma^z L_n(u) \sigma^z, \quad (3.5)$$

$$\hat{L}_n(u + i\pi) = -\sigma^z \hat{L}_n(u) \sigma^z. \quad (3.6)$$

The quasi-periodicity of  $K$ -matrices (2.21) enables us to obtain the associated periodicity property of the transfer matrix  $t(u)$

$$t(u + i\pi) = t(u). \quad (3.7)$$

The above relation implies that the transfer matrix  $t(u)$  can be expressed in terms of  $e^{2u}$  as a Laurent polynomial of the form

$$t(u) = e^{2(N+2)u} t_{N+2} + e^{2(N+1)u} t_{N+1} + \cdots + e^{-2(N+2)u} t_{-(N+2)}, \quad (3.8)$$

where  $\{t_n | n = N+2, N+1, \dots, -(N+2)\}$  form the  $2N+5$  conserved charges. In particular,  $t_{N+2}$  and  $t_{-(N+2)}$  are given by

$$\begin{aligned} t_{N+2} &= -\frac{1}{4} e^{(N+2)\eta} \left\{ e^{\theta_+ - \theta_-} F^{(+)} F^{(-)} + e^{-\theta_+ + \theta_-} D^{(+)} D^{(-)} \right\}, \\ t_{-(N+2)} &= -\frac{1}{4} e^{-(N+2)\eta} \left\{ e^{\theta_+ - \theta_-} F^{(+)} F^{(-)} + e^{-\theta_+ + \theta_-} D^{(+)} D^{(-)} \right\}, \end{aligned}$$

where the constants  $D^{(\pm)}$  and  $F^{(\pm)}$  are determined by (3.2).

Following the method in [35–37] and using the crossing relation of the L-matrix (2.15) and the explicit expressions of the K-matrices (2.19) and (2.20), we verify that the corresponding transfer matrix  $t(u)$  satisfies the following crossing relation

$$t(-u - \eta) = t(u). \quad (3.9)$$

We can define the average value  $\mathcal{O}(u)$  of the matrix elements of the monodromy matrices  $T(u)$  and  $\hat{T}(u)$  (or the  $L$ -operators  $L_n(u)$  and the  $\hat{L}$ -operators  $\hat{L}_n(u)$ ) by using the averaging procedure [38]:

$$\mathcal{O}(u) = \prod_{m=1}^p O(u - m\eta),$$

where the operator  $O(u)$  can be either  $\{\mathbf{A}(u), \mathbf{B}(u), \mathbf{C}(u), \mathbf{D}(u), \hat{\mathbf{A}}(u), \hat{\mathbf{B}}(u), \hat{\mathbf{C}}(u), \hat{\mathbf{D}}(u)\}$  or  $\{A_n(u), B_n(u), C_n(u), D_n(u), \hat{A}_n(u), \hat{B}_n(u), \hat{C}_n(u), \hat{D}_n(u) | n = 1, \dots, N\}$ . It was shown in ref. [38] that

$$\mathcal{T}(u) = \begin{pmatrix} \mathcal{A}(u) & \mathcal{B}(u) \\ \mathcal{C}(u) & \mathcal{D}(u) \end{pmatrix} = \mathcal{L}_N(u) \mathcal{L}_{N-1}(u) \cdots \mathcal{L}_1(u), \quad (3.10)$$

$$\hat{\mathcal{T}}(u) = \begin{pmatrix} \hat{\mathcal{A}}(u) & \hat{\mathcal{B}}(u) \\ \hat{\mathcal{C}}(u) & \hat{\mathcal{D}}(u) \end{pmatrix} = \hat{\mathcal{L}}_1(u) \hat{\mathcal{L}}_2(u) \cdots \hat{\mathcal{L}}_N(u), \quad (3.11)$$

and the average values of each  $L$ -operator and  $\hat{L}$ -operator are given by

$$\begin{aligned} \mathcal{L}_n(u) &= \begin{pmatrix} \mathcal{A}_n(u) & \mathcal{B}_n(u) \\ \mathcal{C}_n(u) & \mathcal{D}_n(u) \end{pmatrix} \\ &= \begin{pmatrix} e^{pu} \{d_n^{(+)}\}^p + e^{-pu} \{d_n^{(-)}\}^p & \{g_n^{(+)}\}^p + \{g_n^{(-)}\}^p \\ \{h_n^{(+)}\}^p + \{h_n^{(-)}\}^p & e^{pu} \{f_n^{(+)}\}^p + e^{-pu} \{f_n^{(-)}\}^p \end{pmatrix}, \end{aligned} \quad (3.12)$$

$$\begin{aligned} \hat{\mathcal{L}}_n(u) &= \begin{pmatrix} \hat{\mathcal{A}}_n(u) & \hat{\mathcal{B}}_n(u) \\ \hat{\mathcal{C}}_n(u) & \hat{\mathcal{D}}_n(u) \end{pmatrix} \\ &= \begin{pmatrix} e^{pu} \{f_n^{(-)}\}^p + e^{-pu} \{f_n^{(+)}\}^p & -\{g_n^{(+)}\}^p - \{g_n^{(-)}\}^p \\ -\{h_n^{(+)}\}^p - \{h_n^{(-)}\}^p & e^{pu} \{d_n^{(-)}\}^p + e^{-pu} \{d_n^{(+)}\}^p \end{pmatrix}, \end{aligned} \quad (3.13)$$

with  $n = 1, \dots, N$ . Note that the average values of the matrix elements are Laurent polynomials of  $e^{pu}$ , which implies

$$\mathcal{T}(u + \eta) = \mathcal{T}(u), \quad \mathcal{L}_n(u + \eta) = \mathcal{L}_n(u), \quad n = 1, \dots, N, \quad (3.14)$$

$$\hat{\mathcal{T}}(u + \eta) = \hat{\mathcal{T}}(u), \quad \hat{\mathcal{L}}_n(u + \eta) = \hat{\mathcal{L}}_n(u), \quad n = 1, \dots, N, \quad (3.15)$$

$$\lim_{u \rightarrow \pm\infty} \mathcal{A}(u) = e^{\pm pNu} \left\{ D^{(\pm)} \right\}^p, \quad (3.16)$$

$$\lim_{u \rightarrow \pm\infty} \hat{\mathcal{A}}(u) = e^{\pm pNu} \left\{ F^{(\mp)} \right\}^p, \quad (3.17)$$

$$\lim_{u \rightarrow \pm\infty} \mathcal{D}(u) = e^{\pm pNu} \left\{ F^{(\pm)} \right\}^p, \quad (3.18)$$

$$\lim_{u \rightarrow \pm\infty} \hat{\mathcal{D}}(u) = e^{\pm pNu} \left\{ D^{(\mp)} \right\}^p. \quad (3.19)$$

### 3.2 Fusion hierarchy

The main tool adopted in this paper to solve the open  $\tau_2$ -model is the so-called fusion technique, by which high-dimensional representations can be obtained from the low-dimensional ones. The fusion technique was first developed in refs. [26–28] for  $R$ -matrices, and then generalised for  $K$ -matrices in refs. [39–42]. In recent years, this technique has been extensively used in solving lots of integrable models [43, 44]. Following the procedure in ref. [27], we introduce the projectors

$$P_{1\dots m}^{(+)} = \frac{1}{m!} \sum_{\sigma \in S_m} \mathcal{P}_\sigma, \quad (3.20)$$

where  $S_m$  is the permutation group of  $m$  indices, and  $\mathcal{P}_\sigma$  is the permutation operator in the tensor space  $\otimes_{k=1}^m \mathcal{C}^2$ . For instance,

$$\begin{aligned} P_{12}^{(+)} &= \frac{1}{2}(1 + \mathcal{P}_{12}), \\ P_{123}^{(+)} &= \frac{1}{6}(1 + \mathcal{P}_{23}\mathcal{P}_{12} + \mathcal{P}_{12}\mathcal{P}_{23} + \mathcal{P}_{12} + \mathcal{P}_{23} + \mathcal{P}_{13}). \end{aligned}$$

The fused spin- $j$   $K^-$ -matrix is given by [40, 41]

$$\begin{aligned} K_{\{a\}}^{-(j)}(u) &= P_{a_1, \dots, a_{2j}}^{(+)} \prod_{k=1}^{2j} \left\{ \left[ \prod_{l=1}^{k-1} R_{a_l, a_k} (2u + (k+l-2j-1)\eta) \right] \right. \\ &\quad \left. \times K_{a_k}^{-\left(\frac{1}{2}\right)} \left( u + \left( k - j - \frac{1}{2} \right) \eta \right) \right\} P_{a_1, \dots, a_{2j}}^{(+)}, \end{aligned} \quad (3.21)$$

where  $\{a\} \equiv \{a_1, \dots, a_{2j}\}$  and  $K^{-\left(\frac{1}{2}\right)}(u) = K^-(u)$ . The fused spin- $j$   $K^+$ -matrix is given by duality

$$K_{\{a\}}^{+(j)}(u) = \frac{1}{f^{(j)}(u)} K_{\{a\}}^{-(j)}(-u - \eta) \Big|_{(\alpha_-, \beta_-, \theta_-) \rightarrow (-\alpha_+, -\beta_+, \theta_+)}, \quad (3.22)$$

where the normalization factor  $f^{(j)}(u)$  is

$$f^{(j)}(u) = \prod_{l=1}^{2j-1} \prod_{k=1}^l [-\rho(2u + (l+k+1-2j)\eta)],$$

with

$$\rho(u) = \sinh(u - \eta) \sinh(u + \eta). \quad (3.23)$$

The fused (boundary) matrices satisfy the generalized (boundary) Yang-Baxter equations [40, 41].

We introduce further the fused spin- $j$  monodromy matrices  $T_{\{a\}}^{(j)}(u)$  and  $\hat{T}_{\{a\}}^{(j)}(u)$  in terms of the fundamental monodromy matrices  $T^{(\frac{1}{2})}(u) = T(u)$  and  $\hat{T}^{(\frac{1}{2})}(u) = \hat{T}(u)$  as follows:

$$\begin{aligned} T_{\{a\}}^{(j)}(u) &= P_{1,\dots,2j}^{(+)} T_1^{(\frac{1}{2})} \left( u - \left( j - \frac{1}{2} \right) \eta \right) T_2^{(\frac{1}{2})} \left( u - \left( j - \frac{1}{2} \right) \eta + \eta \right) \\ &\quad \times \cdots T_{2j}^{(\frac{1}{2})} \left( u + \left( j - \frac{1}{2} \right) \eta \right) P_{1,\dots,2j}^{(+)}, \end{aligned} \quad (3.24)$$

$$\begin{aligned} \hat{T}_{\{a\}}^{(j)}(u) &= P_{1,\dots,2j}^{(+)} \hat{T}_1^{(\frac{1}{2})} \left( u - \left( j - \frac{1}{2} \right) \eta \right) \hat{T}_2^{(\frac{1}{2})} \left( u - \left( j - \frac{1}{2} \right) \eta + \eta \right) \\ &\quad \times \cdots \hat{T}_{2j}^{(\frac{1}{2})} \left( u + \left( j - \frac{1}{2} \right) \eta \right) P_{1,\dots,2j}^{(+)}. \end{aligned} \quad (3.25)$$

The fused transfer matrices  $t^{(j)}(u)$  which correspond to a spin- $j$  auxiliary space can be constructed by the fused monodromy matrices and  $K$ -matrices as

$$t^{(j)}(u) = \text{tr}_{\{a\}} \left\{ K_{\{a\}}^{+(j)}(u) T_{\{a\}}^{(j)}(u) K_{\{a\}}^{-(j)}(u) \hat{T}_{\{a\}}^{(j)}(u) \right\}. \quad (3.26)$$

The double-row transfer matrix  $t(u)$  given by (2.17) corresponds to the fundamental case  $j = \frac{1}{2}$ ; that is  $t^{(\frac{1}{2})}(u) = t(u)$ . Also, the fused transfer matrices constitute commutative families

$$[t^{(j)}(u), t^{(j')}(v)] = 0, \quad j, j' \in \frac{1}{2}, 1, \frac{3}{2}, \dots. \quad (3.27)$$

These transfer matrices also satisfy the so-called fusion hierarchy [39–41, 45]

$$\begin{aligned} t^{(\frac{1}{2})}(u) t^{(j-\frac{1}{2})}(u - j\eta) &= t^{(j)} \left( u - \left( j - \frac{1}{2} \right) \eta \right) + \delta(u) t^{(j-1)} \left( u - (j + \frac{1}{2})\eta \right), \\ j &= \frac{1}{2}, 1, \frac{3}{2}, \dots, \end{aligned} \quad (3.28)$$

with the conventions  $t^{(-\frac{1}{2})}(u) = 0$  and  $t^{(0)} = \text{id}$ . The coefficient  $\delta(u)$ , the so-called quantum determinant [26, 46–48], is given by

$$\delta(u) = -\nu(u) \text{Det}_q \{K^+(u)\} \text{Det}_q \{T(u)\} \text{Det}_q \{K^-(u)\} \text{Det}_q \{\hat{T}(u)\}, \quad (3.29)$$

where

$$\nu(u) = \frac{1}{\sinh(2u + \eta) \sinh(2u - \eta)}, \quad (3.30)$$

$$\text{Det}_q\{T(u)\} = \text{tr}_{12}\{P_{12}^- T_1(u) T_2(u + \eta)\} = \prod_{n=1}^N \text{Det}_q\{L_n(u)\}, \quad (3.31)$$

$$\text{Det}_q\{L_n(u)\} = e^{2u-\eta} d_n^{(+)} f_n^{(+)} + e^{-2u+\eta} d_n^{(-)} f_n^{(-)} - e^\eta g_n^{(+)} h_n^{(-)} - e^{-\eta} g_n^{(-)} h_n^{(+)}, \quad (3.32)$$

$$\text{Det}_q\{\hat{T}(u)\} = \text{tr}\{P_{12}^- \hat{T}_1(u) \hat{T}_2(u + \eta)\} = \prod_{n=1}^N \text{Det}_q\{\hat{L}_n(u)\}, \quad (3.33)$$

$$\text{Det}_q\{\hat{L}_n(u)\} = e^{-2u-\eta} d_n^{(+)} f_n^{(+)} + e^{2u+\eta} d_n^{(-)} f_n^{(-)} - e^\eta g_n^{(+)} h_n^{(-)} - e^{-\eta} g_n^{(-)} h_n^{(+)}, \quad (3.34)$$

$$\begin{aligned} \text{Det}_q\{K^-(u)\} &= \text{tr}_{12}\{P_{12} K_1^-(u) R_{12}(2u + \eta) K_2^-(u + \eta)\} \\ &= -2^2 \sinh(2u - 2\eta) \sinh(u + \alpha_-) \sinh(u - \alpha_-) \\ &\quad \times \cosh(u + \beta_-) \cosh(u - \beta_-), \end{aligned} \quad (3.35)$$

$$\begin{aligned} \text{Det}_q\{K^+(u)\} &= \text{tr}_{12}\{P_{12} K_2^+(u + \eta) R_{12}(-2u - 3\eta) K_2^+(u)\} \\ &= \text{Det}_q\{K^-(u - 2\eta)\}|_{(\alpha_-, \beta_-, \theta_-) \rightarrow (-\alpha_+, -\beta_+, \theta_+)} \\ &= 2^2 \sinh(2u + 2\eta) \sinh(u + \alpha_+) \sinh(u - \alpha_+) \\ &\quad \times \cosh(u + \beta_+) \cosh(u - \beta_+). \end{aligned} \quad (3.36)$$

Let us introduce the functions  $a(u)$  and  $d(u)$  as follows:

$$\begin{aligned} a(u) &= -2^2 \frac{\sinh(2u + 2\eta)}{\sinh(2u + \eta)} \sinh(u - \alpha_-) \sinh(u - \alpha_+) \\ &\quad \times \cosh(u - \beta_-) \cosh(u - \beta_+) \bar{A}(u), \end{aligned} \quad (3.37)$$

$$d(u) = a(-u - \eta), \quad (3.38)$$

where

$$\bar{A}(u) = e^{-N\eta} G^{(-)} H^{(+)} \prod_{n=1}^N \left( e^u - e^{-u+2\eta} \frac{d_n^{(-)} f_n^{(-)}}{g_n^{(-)} h_n^{(+)}} \right) \left( e^{-u} \frac{d_n^{(+)} f_n^{(+)}}{g_n^{(-)} h_n^{(+)}} - e^u \right). \quad (3.39)$$

Similar as the definitions (3.2), the constants  $G^{(\pm)}$  and  $H^{(\pm)}$  are

$$G^{(\pm)} = \prod_{n=1}^N g_n^{(\pm)}, \quad H^{(\pm)} = \prod_{n=1}^N h_n^{(\pm)}. \quad (3.40)$$

Then it is easy to check that the quantum determinant (3.29) can be expressed in terms of the above functions as

$$\delta(u) = a(u)d(u - \eta). \quad (3.41)$$

Similar to the  $\tau_2$ -model with the periodic condition [24], we can also use the recursive relation (3.28) and the coefficient function (3.41) to express the fused transfer matrix  $t^{(j)}(u)$  in terms of the fundamental one  $t^{(\frac{1}{2})}(u)$  with a  $2j$ -order functional relation which can be

expressed as the determinant of some  $2j \times 2j$  matrix [49], namely,

$$t^{(j)}(u) = \begin{vmatrix} t(u+(j-\frac{1}{2})\eta) & -a(u+(j-\frac{1}{2})\eta) \\ -d(u+(j-\frac{3}{2})\eta) & t(u+(j-\frac{3}{2})\eta) & -a(u+(j-\frac{3}{2})\eta) \\ & \ddots & & \\ & & \dots & \\ & & & \ddots \\ & & -d(u-(j+\frac{1}{2})\eta) & t(u-(j+\frac{1}{2})\eta) & -a(u-(j+\frac{1}{2})\eta) \\ & & & -d(u-(j-\frac{1}{2})\eta) & t(u-(j-\frac{1}{2})\eta) \end{vmatrix}, \quad (3.42)$$

$$j = \frac{1}{2}, 1, \frac{3}{2}, \dots.$$

#### 4 Truncation identity

We now proceed to formulate the desired operator identities to determine the spectrum of the transfer matrix  $t(u)$  given by (2.17). For this purpose, we first derive separate truncation identities for the monodromy matrices and  $K$ -matrices. We recall that the fusion approach described in the previous section. When the crossing parameters  $\eta$  takes the special values  $\eta = \frac{2i\pi}{p}$ , one can find that the spin- $\frac{p}{2}$  fused monodromy matrices mentioned in (3.24), (3.25), all take the block-lower triangular forms [38]

$$T^{(\frac{p}{2})}(u) = \begin{pmatrix} \mathcal{A}(u) & \mathcal{B}(u) & 0 \\ \mathcal{C}(u) & \mathcal{D}(u) & 0 \\ g(u) & h(u) & \text{Det}_q\{T(u - (\frac{p-1}{2})\eta)\}FT^{(\frac{p}{2}-1)}(u)F^{-1} \end{pmatrix}, \quad (4.1)$$

$$\hat{T}^{(\frac{p}{2})}(u) = \begin{pmatrix} \hat{\mathcal{A}}(u) & \hat{\mathcal{B}}(u) & 0 \\ \hat{\mathcal{C}}(u) & \hat{\mathcal{D}}(u) & 0 \\ \hat{g}(u) & \hat{h}(u) & \text{Det}_q\{\hat{T}(u - (\frac{p-1}{2})\eta)\}F\hat{T}^{(\frac{p}{2}-1)}(u)F^{-1} \end{pmatrix}, \quad (4.2)$$

where

$$F = M \otimes \sigma_z, \quad (4.3)$$

with

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & [2]_q & 0 & 0 & 0 \\ 0 & 0 & [3]_q & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & 0 & [\frac{p-1}{2}]_q \end{pmatrix}, \quad (4.4)$$

and

$$[x] = \frac{q^x - q^{-x}}{q - q^{-1}}, \quad q = e^{-\eta}. \quad (4.5)$$

Here  $\{g(u), h(u), \hat{g}(u), \hat{h}(u)\}$  are some operators, which are irrelevant to the transfer matrices. Moreover, through tedious calculations, it is found that the general non-diagonal boundary fused  $K^-$ -matrices (3.21) for  $\eta = \frac{2i\pi}{p}$  and  $j = \frac{p}{2}$  take the following form like (4.1)

$$K^{-\left(\frac{p}{2}\right)}(u) = \mu^{\left(\frac{p}{2}\right)}(u) \begin{pmatrix} \mathcal{K}_{11}^{-\left(\frac{p}{2}\right)}(u) & \mathcal{K}_{12}^{-\left(\frac{p}{2}\right)}(u) & 0 \\ \mathcal{K}_{21}^{-\left(\frac{p}{2}\right)}(u) & \mathcal{K}_{22}^{-\left(\frac{p}{2}\right)}(u) & 0 \\ k_3(u) & k_4(u) & K_{33}^{-\left(\frac{p}{2}\right)}(u) \end{pmatrix}, \quad (4.6)$$

where the functions

$$\mu^{\left(\frac{p}{2}\right)}(u) = \prod_{l=1}^{p-1} \prod_{k=1}^l \sinh(2u + (l+k-p+1)\eta), \quad (4.7)$$

$$\begin{aligned} \mathcal{K}_{11}^{-\left(\frac{p}{2}\right)}(u) &= \sum_{l=0}^{\left[\frac{p}{2}\right]} c_p^{2l} (\sinh \alpha_- \cosh \beta_-)^{p-2l} (\cosh \alpha_- \sinh \beta_-)^{2l} \cosh(pu) \\ &\quad + \sum_{l=0}^{\left[\frac{p}{2}\right]} c_p^{2l+1} (\cosh \alpha_- \sinh \beta_-)^{p-2l-1} (\sinh \alpha_- \cosh \beta_-)^{2l+1} \sinh(pu) \\ &\quad + p(\sinh \alpha_- \cosh \beta_- \cosh(pu) + \cosh \alpha_- \sinh \beta_- \sinh(pu)), \end{aligned} \quad (4.8)$$

$$\begin{aligned} \mathcal{K}_{22}^{-\left(\frac{p}{2}\right)}(u) &= \sum_{l=0}^{\left[\frac{p}{2}\right]} c_p^{2l} (\sinh \alpha_- \cosh \beta_-)^{p-2l} (\cosh \alpha_- \sinh \beta_-)^{2l} \cosh(pu) \\ &\quad - \sum_{l=0}^{\left[\frac{p}{2}\right]} c_p^{2l+1} (\cosh \alpha_- \sinh \beta_-)^{p-2l-1} (\sinh \alpha_- \cosh \beta_-)^{2l+1} \sinh(pu) \\ &\quad + p(\sinh \alpha_- \cosh \beta_- \cosh(pu) - \cosh \alpha_- \sinh \beta_- \sinh(pu)), \end{aligned} \quad (4.9)$$

and the  $(p-1) \times (p-1)$  matrix  $K_{33}^{-\left(\frac{p}{2}\right)}(u)$  is related to the spin- $\frac{p-2}{2}$   $K^-$ -matrix by

$$\mathcal{K}_{33}^{-\left(\frac{p}{2}\right)}(u) = \frac{\text{Det}_q \left\{ K^- \left( u - \left( \frac{p-1}{2} \right) \eta \right) \right\}}{\sinh \left[ 2 \left( u - \left( \frac{p-1}{2} \right) \eta \right) + \eta \right]} F K^{-\left(\frac{p-2}{2}\right)}(u) F^{-1}. \quad (4.10)$$

The functions  $\mathcal{K}_{12}^{-\left(\frac{p}{2}\right)}(u)$  and  $\mathcal{K}_{21}^{-\left(\frac{p}{2}\right)}(u)$  give contributions to calculating the asymptotic behavior of the eigenvalue of  $t(u)$  (2.17) and possess the following forms respectively

$$\mathcal{K}_{12}^{-\left(\frac{p}{2}\right)}(u) = \left( \frac{1}{2} \right)^{p-1} e^{p\theta_-} \sinh(2pu), \quad (4.11)$$

$$\mathcal{K}_{21}^{-\left(\frac{p}{2}\right)}(u) = \left( \frac{1}{2} \right)^{p-1} e^{-p\theta_-} \sinh(2pu), \quad (4.12)$$

with the matrix  $F$  being given by (4.3). Moreover,  $k_3(u)$  and  $k_4(u)$  are two  $p \times 1$  matrices which are irrelevant to the associated transfer matrix.

At the same time, the fused  $K^+$ -matrices are given, in view of eq. (3.22) with  $\eta = \frac{2i\pi}{p}$  and  $j = \frac{p}{2}$ , by

$$K_{\{a\}}^{+\left(\frac{p}{2}\right)}(u) = \frac{1}{f^{\left(\frac{p}{2}\right)}(u)} K_{\{a\}}^{-\left(\frac{p}{2}\right)}(-u - \eta) \Big|_{(\alpha_-, \beta_-, \theta_-) \rightarrow (-\alpha_+, -\beta_+, \theta_+)}. \quad (4.13)$$

The explicit expressions of the elements of the fused monodromy matrices and the  $K$ -matrices for the cases  $p = 3$  are given in appendix A.

Hence, we are finally in position to formulate the truncation identity for the fused transfer matrices  $t^{(j)}(u)$  defined in (3.26). Based on the results of (4.1) and (4.2) for the fused monodromy matrices and those of (4.6) and (4.13) for the fused  $K$ -matrices, we obtain

$$t^{(\frac{p}{2})}(u) = \left( \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) \right) \times \text{id} + \delta \left( u - \left( \frac{p-1}{2} \right) \eta \right) t^{(\frac{p-2}{2})}(u), \quad (4.14)$$

where the coefficients  $\tilde{\mathcal{A}}(u)$  and  $\tilde{\mathcal{D}}(u)$  are given by.<sup>1</sup>

$$\begin{aligned} \tilde{\mathcal{A}}(u) &= \left[ \mathcal{K}_{11}^{+(\frac{p}{2})}(u) \mathcal{A}(u) + \mathcal{K}_{12}^{+(\frac{p}{2})}(u) \mathcal{C}(u) \right] \left[ \mathcal{K}_{11}^{-(\frac{p}{2})}(u) \hat{\mathcal{A}}(u) + \mathcal{K}_{12}^{-(\frac{p}{2})}(u) \hat{\mathcal{C}}(u) \right] \\ &\quad + \left[ \mathcal{K}_{11}^{+(\frac{p}{2})}(u) \mathcal{B}(u) + \mathcal{K}_{12}^{+(\frac{p}{2})}(u) \mathcal{D}(u) \right] \left[ \mathcal{K}_{21}^{-(\frac{p}{2})}(u) \hat{\mathcal{A}}(u) + \mathcal{K}_{22}^{-(\frac{p}{2})}(u) \hat{\mathcal{C}}(u) \right], \end{aligned} \quad (4.15)$$

$$\begin{aligned} \tilde{\mathcal{D}}(u) &= \left[ \mathcal{K}_{21}^{+(\frac{p}{2})}(u) \mathcal{A}(u) + \mathcal{K}_{22}^{+(\frac{p}{2})}(u) \mathcal{C}(u) \right] \left[ \mathcal{K}_{11}^{-(\frac{p}{2})}(u) \hat{\mathcal{B}}(u) + \mathcal{K}_{12}^{-(\frac{p}{2})}(u) \hat{\mathcal{D}}(u) \right] \\ &\quad + \left[ \mathcal{K}_{21}^{+(\frac{p}{2})}(u) \mathcal{B}(u) + \mathcal{K}_{22}^{+(\frac{p}{2})}(u) \mathcal{D}(u) \right] \left[ \mathcal{K}_{21}^{-(\frac{p}{2})}(u) \hat{\mathcal{B}}(u) + \mathcal{K}_{22}^{-(\frac{p}{2})}(u) \hat{\mathcal{D}}(u) \right]. \end{aligned} \quad (4.16)$$

It is remarked that the functions  $\mathcal{K}_{11}^{\pm(\frac{p}{2})}(u), \mathcal{K}_{12}^{\pm(\frac{p}{2})}(u), \mathcal{K}_{21}^{\pm(\frac{p}{2})}(u), \mathcal{K}_{22}^{\pm(\frac{p}{2})}(u)$  and the average values of each monodromy matrices are invariant under shifting with  $\eta$ .

Combining the fusion hierarchy (3.28) and the closing relation (4.14) for  $\eta = \frac{2i\pi}{p}$ , we arrive at the functional relation for the fundamental transfer matrix straightforward. Here we give an example of the functional relations for  $p = 3$ :

$$t(u + \eta) t(u) t(u - \eta) - \delta(u + \eta) t(u - \eta) - \delta(u) t(u + \eta) = \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) + \delta(u - \eta) t(u).$$

## 5 Eigenvalues of the fundamental transfer matrix

### 5.1 Functional relations of eigenvalues

The commutativity (3.27) of the fused transfer matrices  $\{t^{(j)}(u)\}$  with different spectral parameters implies that they have common eigenstates. One can set  $|\Psi\rangle$  to be a common eigenstate of these fused transfer matrices with eigenvalues  $\Lambda^{(j)}(u)$ , i.e.,

$$t^{(j)}(u) |\Psi\rangle = \Lambda^{(j)}(u) |\Psi\rangle.$$

The quasi-periodicity (3.7) and the cross relation (3.9) of the transfer matrix  $t(u)$  implies that the corresponding eigenvalue  $\Lambda(u)$  satisfies the properties

$$\Lambda(u + i\pi) = \Lambda(u), \quad (5.1)$$

$$\Lambda(-u - \eta) = \Lambda(u). \quad (5.2)$$

---

<sup>1</sup>The average values  $\{\mathcal{A}(u), \mathcal{B}(u), \mathcal{C}(u), \mathcal{D}(u)\}$  and  $\{\hat{\mathcal{A}}(u), \hat{\mathcal{B}}(u), \hat{\mathcal{C}}(u), \hat{\mathcal{D}}(u)\}$  of the matrix elements of the monodromy matrices  $T(u)$  and  $\hat{T}(u)$  in (4.15) and (4.16) can be obtained from the relations in (3.10) and (3.11) which are based on the average values of each  $L$ -operator and  $\hat{L}$ -operator given in (3.12) and (3.13). For some examples of the small sites such as  $N = 1, 2$ , we give the explicit expressions of the average value functions and also discuss some special constraints that allow one to calculate these average value functions for an arbitrary number of the sites in appendix B.

The asymptotic behavior (3.1) and the special values at  $u = 0, \frac{i\pi}{2}$  of the transfer matrix  $t(u)$  enables us to derive that the corresponding eigenvalue  $\Lambda(u)$  have the following functional relations

$$\lim_{u \rightarrow \pm\infty} \Lambda(u) = -\frac{1}{4} e^{\pm\{(2N+4)u+(N+2)\eta\}} \left\{ e^{\theta_+-\theta_-} F^{(+)}F^{(-)} + e^{-\theta_++\theta_-} D^{(+)}D^{(-)} \right\}, \quad (5.3)$$

$$\begin{aligned} \Lambda(0) &= -2^3 \sinh \alpha_- \cosh \beta_- \sinh \alpha_+ \cosh \beta_+ \cosh \eta \\ &\times \prod_{n=1}^N \left( e^{-\eta} d_n^{(+)} f_n^{(+)} + e^\eta d_n^{(-)} f_n^{(-)} - e^\eta g_n^{(+)} h_n^{(-)} - e^{-\eta} g_n^{(-)} h_n^{(+)} \right), \end{aligned} \quad (5.4)$$

$$\begin{aligned} \Lambda\left(\frac{i\pi}{2}\right) &= -2^3 \cosh \alpha_- \sinh \beta_- \cosh \alpha_+ \sinh \beta_+ \cosh \eta \\ &\times \prod_{n=1}^N \left( e^{-\eta} d_n^{(+)} f_n^{(+)} + e^\eta d_n^{(-)} f_n^{(-)} + e^\eta g_n^{(+)} h_n^{(-)} + e^{-\eta} g_n^{(-)} h_n^{(+)} \right). \end{aligned} \quad (5.5)$$

The analyticity of the  $L$ -operator (2.3), the quasi-periodicity (5.1) and the asymptotic behavior (5.3) of the eigenvalue give rise to that  $\Lambda(u)$  possesses the following analytical property

$$\Lambda(u), \text{ as a function of } e^u, \text{ is a Laurent polynomial of degree } 2N+4 \text{ like (3.8).} \quad (5.6)$$

According to the fusion hierarchy relation (3.28) and the determinant representation (3.42) of the fused transfer matrices, the eigenvalues  $\Lambda^{(j)}(u)$  give some similar representation in terms of the fundamental one  $\Lambda(u) = \Lambda^{(\frac{1}{2})}(u)$

$$\Lambda^{(j)}(u) = \begin{vmatrix} \Lambda(u+(j-\frac{1}{2})\eta) & -a(u+(j-\frac{1}{2})\eta) & & & \\ -d(u+(j-\frac{3}{2})\eta) & \Lambda(u+(j-\frac{3}{2})\eta) & -a(u+(j-\frac{3}{2})\eta) & & \\ & \ddots & & \ddots & \\ & & \cdots & & \\ & & & -d(u-(j+\frac{1}{2})\eta) & \Lambda(u-(j+\frac{1}{2})\eta) & -a(u-(j+\frac{1}{2})\eta) \\ & & & & -d(u-(j-\frac{1}{2})\eta) & \Lambda(u-(j-\frac{1}{2})\eta) \end{vmatrix}, \quad (5.7)$$

$$j = \frac{1}{2}, 1, \frac{3}{2}, \dots,$$

where the functions  $a(u)$  and  $d(u)$  are given by (3.37) and (3.38). The truncation identity (4.14) of the spin- $\frac{p}{2}$  transfer matrix leads to the fact that the corresponding eigenvalue  $\Lambda^{(\frac{p}{2})}(u)$  satisfies the relation

$$\Lambda^{(\frac{p}{2})}(u) = \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) + \delta \left( u - \left( \frac{p-1}{2} \right) \eta \right) \Lambda^{(\frac{p-2}{2})}(u), \quad (5.8)$$

where the functions  $\tilde{\mathcal{A}}(u)$  and  $\tilde{\mathcal{D}}(u)$  are given by (4.15)–(4.16). For example, the functional relation of the eigenvalue for  $p = 3$  is

$$\Lambda(u+\eta) \Lambda(u) \Lambda(u-\eta) - \delta(u+\eta) \Lambda(u-\eta) - \delta(u) \Lambda(u+\eta) = \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) + \delta(u-\eta) \Lambda(u).$$

It is believed [1, 3, 50] that the relations (5.1)–(5.5), the analytic property (5.6) and the truncation identity (5.8) allow us to completely determine the eigenvalues  $\Lambda(u)$  of the fundamental transfer matrix  $t(u)$  given by (2.17).

## 5.2 T-Q relation

### 5.2.1 Generic case

Following the ODBA method [23] and the method developed in [24], we can express eigenvalues of  $t(u)$  in terms of some inhomogeneous  $T - Q$  relation

$$\Lambda(u) = a(u) \frac{Q(u - \eta)}{Q(u)} + d(u) \frac{Q(u + \eta)}{Q(u)} + 2^{(1-p)2N-4p+2} c \frac{\sinh(2u) \sinh(2u + 2\eta) F(u)}{Q(u)}, \quad (5.9)$$

where the functions  $a(u)$  and  $d(u)$  are given by (3.37) and (3.38), and the constant  $c$  is uniquely determined by the inhomogeneous parameters and boundary parameters,

$$\begin{aligned} & \left( \frac{1}{2} \right)^{2p} c \left\{ e^{p(\theta_+ - \theta_-)} \{F^{(+)} F^{(-)}\}^p + e^{-p(\theta_+ - \theta_-)} \{D^{(+)} D^{(-)}\}^p - (-1)^N e^{-p(\alpha_+ + \beta_+ + \alpha_- + \beta_-)} \right. \\ & \times \{G^{(-)} H^{(+)}\}^p - (-1)^N e^{p(\alpha_+ + \beta_+ + \alpha_- + \beta_-)} \{G^{(+)} H^{(-)}\}^p \Big\} = \frac{1}{4} \left\{ e^{\theta_+ - \theta_-} F^{(+)} F^{(-)} \right. \\ & + e^{-(\theta_+ - \theta_-)} D^{(+)} D^{(-)} - (-1)^N e^{-(\alpha_+ + \beta_+ + \alpha_- + \beta_-)} \{G^{(-)} H^{(+)}\} e^{-\eta} \\ & \left. \left. - (-1)^N e^{\alpha_+ + \beta_+ + \alpha_- + \beta_-} \{G^{(+)} H^{(-)}\} e^\eta \right\}. \right. \end{aligned} \quad (5.10)$$

The trigonometric polynomial  $Q(u)$  is parameterized by  $(p-1)N + 2p$  Bethe roots  $\{\lambda_j\}$

$$Q(u) = \prod_{j=1}^{(p-1)N+2p} \sinh(u - \lambda_j) \sinh(u + \lambda_j + \eta), \quad (5.11)$$

which will be specified by the associated BAEs (5.16) below. The function  $F(u)$  which is a Laurent polynomial of degree  $p(2N + 4)$  is given by

$$F(u) = \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) - \bar{\mathcal{A}}(u) - \bar{\mathcal{D}}(u), \quad (5.12)$$

$$\bar{\mathcal{A}}(u) = \prod_{m=1}^p a(u - m\eta), \quad \bar{\mathcal{D}}(u) = \prod_{m=1}^p d(u - m\eta), \quad (5.13)$$

where the functions  $\tilde{\mathcal{A}}(u)$  and  $\tilde{\mathcal{D}}(u)$  are given in (4.15) and (4.16) respectively.

According to the relations (3.14)–(3.15), the definitions (5.13), (4.15) and (4.16) and the explicit expression (4.8)–(4.13) of the elements of the  $K$ -matrices, the function  $F(u)$  can be reduced as a Laurent polynomial of  $e^{pu}$  with a degree  $2N + 4$  (i.e. there are only  $2N + 5$  non-vanishing coefficients), namely,

$$F(u) = \sum_{l=0}^{2N+4} F^{(2p(N+2-l))} (\{d_n^\pm, f_n^\pm, g_n^\pm, h_n^\pm, d_n^\pm, \alpha_\pm, \beta_\pm, \theta_\pm\}) e^{p(2N+4-2l)u}, \quad (5.14)$$

where the  $2N + 5$  coefficients  $\{F^{(2p(N+2-l))}|l = 0, 1, \dots, 2N + 4\}$  are polynomial of the inhomogeneity parameters  $\{d_n^\pm, f_n^\pm, g_n^\pm, h_n^\pm|n = 1, 2, \dots, N\}$  and the boundary parameters  $\{\alpha_\pm, \beta_\pm, \theta_\pm\}$ , and that we can easily prove that the function  $F(u)$  holds the crossing property

$$F(-u - \eta) = F(u). \quad (5.15)$$

The fact that the constant  $c$  satisfies the relation (5.10) ensures that  $\Lambda(u)$  given by (5.9) matches the asymptotic behavior (5.3). The  $(p-1)N+2p$  parameters  $\{\lambda_j|j=1,\dots,(p-1)N+2p\}$  satisfy the associated Bethe Ansatz equations (BAEs)

$$\begin{aligned} a(\lambda_j)Q(\lambda_j - \eta) + d(\lambda_j)Q(\lambda_j + \eta) + 2^{2(1-p)N-4p+2}c \sinh(2\lambda_j) \\ \times \sinh(2\lambda_j + 2\eta)F(\lambda_j) = 0, \quad j = 1, \dots, (p-1)N+2p, \end{aligned} \quad (5.16)$$

which assure that  $\Lambda(u)$  given by (5.9) is indeed a trigonometric polynomial of  $u$ . It is easy to check that  $\Lambda(u)$  given by the inhomogeneous  $T-Q$  relation (5.9) satisfies the properties (5.1)–(5.2) and the functional relations (5.3)–(5.5). Using the method in the appendix A of [24], we have checked that the  $T-Q$  relation (5.9) also make the functional relation (5.8) fulfilled. Therefore, the resulting expression of  $\Lambda(u)$  constructed by the inhomogeneous  $T-Q$  relation (5.9) is the eigenvalue of the fundamental transfer matrix  $t(u)$  of the  $\tau_2$ -model with generic boundary condition.

### 5.2.2 Degenerate case

The third term of the  $T-Q$  relation (5.9) does not vanish when the generic inhomogeneous parameters  $\{d_n^{(\pm)}, f_n^{(\pm)}, g_n^{(\pm)}, h_n^{(\pm)}|n=1,\dots,N\}$  (only obey the constraint (2.4) which ensure the integrability of the model) and the boundary parameters are absence of restriction. Here we consider some special case making the inhomogeneous term vanishes. When the inhomogeneous parameters  $\{d_n^{(\pm)}, f_n^{(\pm)}, g_n^{(\pm)}, h_n^{(\pm)}|n=1,\dots,N\}$  and the boundary parameters  $\{\alpha_{\pm}, \beta_{\pm}, \theta_{\pm}\}$  obey the following extra constraints besides (2.4):

$$\begin{aligned} e^{\theta_+-\theta_-}F^{(+)}F^{(-)} + e^{-(\theta_+-\theta_-)}D^{(+)}D^{(-)} - (-1)^N e^{-(\alpha_++\beta_++\alpha_-+\beta_-)}\{G^{(-)}H^{(+)}\}e^{-\eta} \\ - (-1)^N e^{\alpha_++\beta_++\alpha_-+\beta_-}\{G^{(+)}H^{(-)}\}e^\eta = 0, \end{aligned} \quad (5.17)$$

$$F^{(2(N+2-l)p)}(\{d_n^{(\pm)}, f_n^{(\pm)}, g_n^{(\pm)}, h_n^{(\pm)}, \alpha_{\pm}, \beta_{\pm}, \theta_{\pm}\}) = 0, \quad l = 1, \dots, N+2, \quad (5.18)$$

where  $D^{(\pm)}$ ,  $F^{(\pm)}$ ,  $G^{(\pm)}$  and  $H^{(\pm)}$  are given by (3.2) and (3.40), and each  $F^{(2(N+2-l)p)}$  given in (5.14). The corresponding inhomogeneous  $T-Q$  relation (5.9) reduces to the conventional one [29]:

$$\Lambda(u) = a(u)\frac{\bar{Q}(u-\eta)}{\bar{Q}(u)} + d(u)\frac{\bar{Q}(u+\eta)}{\bar{Q}(u)}, \quad (5.19)$$

where the function  $\bar{Q}(u)$  becomes [19–23, 45, 51–57]

$$\bar{Q}(u) = \prod_{j=1}^M \sinh(u - \lambda_j) \sinh(u + \lambda_j + \eta). \quad (5.20)$$

Here the positive integer  $M$  has to satisfy the following constraint in order to match the the asymptotic behavior (5.3) of  $\Lambda(u)$ , namely,

$$\begin{aligned} \{e^{\theta_+-\theta_-}F^{(+)}F^{(-)} + e^{-(\theta_+-\theta_-)}D^{(+)}D^{(-)}\} - (-1)^N e^{-(\alpha_++\beta_++\alpha_-+\beta_-)}G^{(-)}H^{(+)}e^{-(2N+2M+1)\eta} \\ - (-1)^N e^{(\alpha_++\beta_++\alpha_-+\beta_-)}G^{(+)}H^{(-)}e^{(2N+2M+1)\eta} = 0. \end{aligned} \quad (5.21)$$

Moreover, the  $M$  parameters  $\{\lambda_j|j=1,\dots,M\}$  need to satisfy the associated BAEs

$$\frac{a(\lambda_j)}{d(\lambda_j)} = -\frac{\bar{Q}(\lambda_j + \eta)}{\bar{Q}(\lambda_j - \eta)}, \quad j = 1, \dots, M. \quad (5.22)$$

It is easy to check that the relation (5.17) give rise to  $F^{(p(2N+4))} = F^{(-p(2N+4))} = 0$ . Together with (5.18) and the constrained case in (5.17), we get that the function  $F(u)$  indeed vanishes, namely,  $F(u) = 0$ . Substituting (5.19) into (5.7), we have

$$\begin{aligned}\Lambda^{\left(\frac{p}{2}\right)}(u) &= \bar{\mathcal{A}}(u) + \bar{\mathcal{D}}(u) + \delta\left(u - \left(\frac{p-1}{2}\right)\eta\right)\Lambda^{\left(\frac{p-2}{2}\right)}(u) \\ &= \tilde{\mathcal{A}}(u) + \tilde{\mathcal{D}}(u) + \delta\left(u - \left(\frac{p-1}{2}\right)\eta\right)\Lambda^{\left(\frac{p-2}{2}\right)}(u).\end{aligned}\quad (5.23)$$

Similar with the periodic case, we can also prove that the reduced  $T - Q$  relation (5.19) satisfies the functional relations (5.3)–(5.5) of the transfer matrix and the truncation identity of the fused transfer matrices (5.8) when the inhomogeneity parameters and the boundary parameters  $\{d_n^{(\pm)}, f_n^{(\pm)}, g_n^{(\pm)}, h_n^{(\pm)}, \alpha_{\pm}, \beta_{\pm}, \theta_{\pm} | n = 1, \dots, N\}$  satisfy the constraints (2.4), (5.17), (5.18) and (5.21).

## 6 Conclusion

In this paper, we have studied the most general cyclic representation of the quantum  $\tau_2$ -model (also known as Baxter-Bazhanov-Stroganov (BBS) model) with generic integrable boundary conditions via the ODBA method [23]. Based on the truncation identity (4.14) of the fused transfer matrices obtained from the fusion technique, we construct the corresponding inhomogeneous  $T - Q$  relation (5.9) and the associated BAEs (5.16) for the eigenvalue of the fundamental transfer matrix  $t(u)$ .

It is remarked that if the generic inhomogeneity parameters  $\{d_n^{(\pm)}, f_n^{(\pm)}, g_n^{(\pm)}, h_n^{(\pm)} | n = 1, \dots, N\}$  (only obey the constraint (2.4) which ensures the integrability of the model) and the boundary parameters take the generic values, the inhomogeneous term (i.e., the third term) in the  $T - Q$  relation (5.9) *does not* vanish, as long as one requires a polynomial  $Q$ -function. However, if these inhomogeneity parameters and the boundary parameters satisfy the extra constraints (5.17), (5.18) and (5.21), the resulting inhomogeneous  $T - Q$  relation (5.9) reduces to the conventional one (5.19).

**Note added.** After this paper was completed we became aware of the recent results reported in [58]. The authors use the Sklyanin's separation of variables (SoV) method [59–61] to study the spectral problem for the open  $\tau_2$ -model with some constrains on inhomogeneous parameters and also on the boundary parameters.

## Acknowledgments

We would like to thank Prof. Y. Wang for his valuable discussions and continuous encouragement. The financial supports from the National Natural Science Foundation of China (Grant Nos. 11375141, 11374334, 11347025, 11434013, 11425522, and 11547045), BCMIIS, 2016YFA0300603, and the Strategic Priority Research Program of the Chinese Academy of Sciences are gratefully acknowledged. X. Xu also acknowledges the support by the NWU graduate student innovation fund No. YZZ14102.

## A Specific cases of the fused $K$ -matrices

In this appendix we present the explicit expressions of the matrix elements of the fused  $K$ -matrices  $K^{\pm(\frac{p}{2})}(u)$  given in (4.6) and (4.13) for the  $p = 3$  case as an example. In this case, the corresponding matrix elements are

$$\begin{aligned} \mathcal{K}_{11}^{-(\frac{3}{2})}(u) &= \left(\frac{1}{2}\right)^2 \left\{ (\sinh \alpha_- \cosh \beta_-)^3 \cosh(3u) + 3(\sinh \alpha_- \cosh \beta_-)^2 \cosh \alpha_- \sinh \beta_- \sinh(3u) \right. \\ &\quad + 3 \sinh \alpha_- \cosh \beta_- (\cosh \alpha_- \sinh \beta_-)^2 \cosh(3u) + (\cosh \alpha_- \sinh \beta_-)^3 \sinh(3u) \\ &\quad \left. + 3 \sinh \alpha_- \cosh \beta_- \cosh(3u) + 3 \cosh \alpha_- \sinh \beta_- \sinh(3u) \right\}, \end{aligned} \quad (\text{A.1})$$

$$\begin{aligned} \mathcal{K}_{22}^{-(\frac{3}{2})}(u) &= \left(\frac{1}{2}\right)^2 \left\{ (\sinh \alpha_- \cosh \beta_-)^3 \cosh(3u) - 3(\sinh \alpha_- \cosh \beta_-)^2 \cosh \alpha_- \sinh \beta_- \sinh(3u) \right. \\ &\quad + 3 \sinh \alpha_- \cosh \beta_- (\cosh \alpha_- \sinh \beta_-)^2 \cosh(3u) - (\cosh \alpha_- \sinh \beta_-)^3 \sinh(3u) \\ &\quad \left. + 3 \sinh \alpha_- \cosh \beta_- \cosh(3u) - 3 \cosh \alpha_- \sinh \beta_- \sinh(3u) \right\}, \end{aligned} \quad (\text{A.2})$$

$$\mathcal{K}_{12}^{-(\frac{3}{2})}(u) = \left(\frac{1}{2}\right)^2 e^{3\theta_-} \sinh(6u), \quad \mathcal{K}_{21}^{-(\frac{3}{2})}(u) = \left(\frac{1}{2}\right)^2 e^{-3\theta_-} \sinh(6u), \quad (\text{A.3})$$

$$K_{33}^{-(\frac{3}{2})}(u) = \frac{\text{Det}_q\{K^-(u-\eta)\}}{\sinh(2u-\eta)} \sigma^z K^-(u) \sigma^z, \quad (\text{A.4})$$

and

$$\begin{aligned} \mathcal{K}_{11}^{+(\frac{3}{2})}(u) &= \left(\frac{1}{2}\right)^2 \left\{ -(\sinh \alpha_+ \cosh \beta_+)^3 \cosh(3u) + 3(\sinh \alpha_+ \cosh \beta_+)^2 \cosh \alpha_+ \sinh \beta_+ \sinh(3u) \right. \\ &\quad - 3 \sinh \alpha_+ \cosh \beta_+ (\cosh \alpha_+ \sinh \beta_+)^2 \cosh(3u) + (\cosh \alpha_+ \sinh \beta_+)^3 \sinh(3u) \\ &\quad \left. - 3 \sinh \alpha_+ \cosh \beta_+ \cosh(3u) + 3 \cosh \alpha_+ \sinh \beta_+ \sinh(3u) \right\}, \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned} \mathcal{K}_{22}^{+(\frac{3}{2})}(u) &= -\left(\frac{1}{2}\right)^2 \left\{ (\sinh \alpha_+ \cosh \beta_+)^3 \cosh(3u) + 3(\sinh \alpha_+ \cosh \beta_+)^2 \cosh \alpha_+ \sinh \beta_+ \sinh(3u) \right. \\ &\quad + 3 \sinh \alpha_+ \cosh \beta_+ (\cosh \alpha_+ \sinh \beta_+)^2 \cosh(3u) + (\cosh \alpha_+ \sinh \beta_+)^3 \sinh(3u) \\ &\quad \left. + 3 \sinh \alpha_+ \cosh \beta_+ \cosh(3u) + 3 \cosh \alpha_+ \sinh \beta_+ \sinh(3u) \right\}, \end{aligned} \quad (\text{A.6})$$

$$\mathcal{K}_{12}^{+(\frac{3}{2})}(u) = -\left(\frac{1}{2}\right)^2 e^{3\theta_+} \sinh(6u), \quad \mathcal{K}_{21}^{+(\frac{3}{2})}(u) = -\left(\frac{1}{2}\right)^2 e^{-3\theta_+} \sinh(6u), \quad (\text{A.7})$$

$$K_{33}^{+(\frac{3}{2})}(u) = -\frac{\text{Det}_q\{K^+(u-\eta)\}}{\sinh 2u} \sigma^z K^+(u) \sigma^z. \quad (\text{A.8})$$

## B Explicit expression of the average value functions

In this appendix we discuss certain properties of the average values of the matrix elements of the monodromy matrices  $T(u)$  and  $\hat{T}(u)$  given by (3.10) and (3.11) respectively. Here we present the explicit expressions of these average value functions  $\mathcal{A}(u), \mathcal{B}(u), \mathcal{C}(u), \mathcal{D}(u)$  and  $\hat{\mathcal{A}}(u), \hat{\mathcal{B}}(u), \hat{\mathcal{C}}(u), \hat{\mathcal{D}}(u)$  for some small sites cases (namely,  $N = 1, 2$ ). For  $N = 1$ , they are given by

$$\mathcal{A}(u) = e^{pu} \{d_1^{(+)}\}^p + e^{-pu} \{d_1^{(-)}\}^p, \quad (\text{B.1})$$

$$\mathcal{D}(u) = e^{pu} \{f_1^{(+)}\}^p + e^{-pu} \{f_1^{(-)}\}^p, \quad (\text{B.2})$$

$$\mathcal{B}(u) = \{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p, \quad \mathcal{C}(u) = \{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p, \quad (\text{B.3})$$

$$\hat{\mathcal{A}}(u) = e^{pu}\{f_1^{(-)}\}^p + e^{-pu}\{f_1^{(+)}\}^p, \quad (\text{B.4})$$

$$\hat{\mathcal{D}}(u) = e^{pu}\{d_1^{(-)}\}^p + e^{-pu}\{d_1^{(+)}\}^p, \quad (\text{B.5})$$

$$\hat{\mathcal{B}}(u) = -\{g_1^{(+)}\}^p - \{g_1^{(-)}\}^p, \quad \hat{\mathcal{C}}(u) = -\{h_1^{(+)}\}^p - \{h_1^{(-)}\}^p. \quad (\text{B.6})$$

For  $N = 2$ , they are

$$\begin{aligned} \mathcal{A}(u) &= e^{2pu}\{d_1^{(+)}d_2^{(+)}\}^p + e^{-2pu}\{d_1^{(-)}d_2^{(-)}\}^p + \{d_1^{(-)}d_2^{(+)}\}^p + \{d_1^{(+)}d_2^{(-)}\}^p \\ &\quad + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right), \end{aligned} \quad (\text{B.7})$$

$$\begin{aligned} \mathcal{D}(u) &= e^{2pu}\{f_1^{(+)}f_2^{(+)}\}^p + e^{-2pu}\{f_1^{(-)}f_2^{(-)}\}^p + \{f_1^{(-)}f_2^{(+)}\}^p + \{f_1^{(+)}f_2^{(-)}\}^p \\ &\quad + \left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right), \end{aligned} \quad (\text{B.8})$$

$$\begin{aligned} \mathcal{B}(u) &= e^{pu}\left\{\left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\{d_2^{(+)}\}^p + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\{f_1^{(+)}\}^p\right\} \\ &\quad + e^{-pu}\left\{\left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\{d_2^{(-)}\}^p + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\{f_1^{(-)}\}^p\right\}, \end{aligned} \quad (\text{B.9})$$

$$\begin{aligned} \mathcal{C}(u) &= e^{pu}\left\{\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right)\{f_2^{(+)}\}^p + \left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right)\{d_1^{(+)}\}^p\right\} \\ &\quad + e^{-pu}\left\{\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right)\{f_2^{(-)}\}^p + \left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right)\{d_1^{(-)}\}^p\right\}, \end{aligned} \quad (\text{B.10})$$

$$\begin{aligned} \hat{\mathcal{A}}(u) &= e^{2pu}\{f_1^{(-)}f_2^{(-)}\}^p + e^{-2pu}\{f_1^{(+)}f_2^{(+)}\}^p + \{f_1^{(-)}f_2^{(+)}\}^p + \{f_1^{(+)}f_2^{(-)}\}^p \\ &\quad + \left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right), \end{aligned} \quad (\text{B.11})$$

$$\begin{aligned} \hat{\mathcal{D}}(u) &= e^{2pu}\{d_1^{(-)}d_2^{(-)}\}^p + e^{-2pu}\{d_1^{(+)}d_2^{(+)}\}^p + \{d_1^{(-)}d_2^{(+)}\}^p + \{d_1^{(+)}d_2^{(-)}\}^p \\ &\quad + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right), \end{aligned} \quad (\text{B.12})$$

$$\begin{aligned} \hat{\mathcal{B}}(u) &= -e^{pu}\left\{\left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\{d_2^{(-)}\}^p + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\{f_1^{(-)}\}^p\right\} \\ &\quad - e^{-pu}\left\{\left(\{g_1^{(+)}\}^p + \{g_1^{(-)}\}^p\right)\{d_2^{(+)}\}^p + \left(\{g_2^{(+)}\}^p + \{g_2^{(-)}\}^p\right)\{f_1^{(+)}\}^p\right\}, \end{aligned} \quad (\text{B.13})$$

$$\begin{aligned} \hat{\mathcal{C}}(u) &= -e^{pu}\left\{\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right)\{f_2^{(-)}\}^p + \left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right)\{d_1^{(-)}\}^p\right\} \\ &\quad - e^{-pu}\left\{\left(\{h_1^{(+)}\}^p + \{h_1^{(-)}\}^p\right)\{f_2^{(+)}\}^p + \left(\{h_2^{(+)}\}^p + \{h_2^{(-)}\}^p\right)\{d_1^{(+)}\}^p\right\}. \end{aligned} \quad (\text{B.14})$$

The results show that the average value functions become tedious when the number of the lattice sites  $N$  goes large. However, they can be worked out for an arbitrary  $N$  when the inhomogeneous parameters  $\{g_n^{(+)}\}$ ,  $\{g_n^{(-)}\}$ ,  $\{h_n^{(+)}\}$ ,  $\{h_n^{(-)}\}$  associated with each site  $n$  satisfy some constraints. One case is the parameters  $\{g_n^{(+)}\}$ ,  $\{g_n^{(-)}\}$ ,  $\{h_n^{(+)}\}$ ,  $\{h_n^{(-)}\}|_{n=1,\dots,N}$  obey the chiral Potts constraints [38]

$$\frac{\{g_n^{(+)}\}^p + \{g_n^{(-)}\}^p}{\{h_n^{(+)}\}^p + \{h_n^{(-)}\}^p} = \lambda, \quad (\text{B.15})$$

where  $\lambda$  is arbitrary constant. The other case is that the parameters  $d_n^{(+)}\}$ ,  $d_n^{(-)}\}$ ,  $g_n^{(+)}\}$ ,  $g_n^{(-)}\}$ ,  $h_n^{(+)}\}$ ,  $h_n^{(-)}\}$ ,  $f_n^{(+)}\}$  and  $f_n^{(-)}\}$  are independent of the site (i.e.,  $n$ ), which corresponds to the

homogeneous model. In the both cases, the average values of  $L$ -operators  $\{\mathcal{L}_n(u)|n=1, 2, \dots, N\}$  given in (3.12) (or  $\{\hat{\mathcal{L}}_n(u)|n=1, 2, \dots, N\}$  given in (3.13)) with different sites commute with each other. Thus one can diagonalize them simultaneously. Then the average values of the elements of monodromy matrices (3.10) and (3.11) can be easily obtained.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] R.J. Baxter, *Transfer matrix functional relations for the generalized  $\tau_2(t_q)$  model*, *J. Statist. Phys.* **117** (2004) 1 [[cond-mat/0409493](#)] [[INSPIRE](#)].
- [2] R.J. Baxter, *Superintegrable chiral Potts model: Thermodynamic properties, an inverse model and a simple associated Hamiltonian*, *J. Statist. Phys.* **57** (1989) 1 [[INSPIRE](#)].
- [3] V.V. Bazhanov and Yu. G. Stroganov, *Chiral Potts model as a descendant of the six vertex model*, *J. Statist. Phys.* **59** (1990) 799 [[INSPIRE](#)].
- [4] R.J. Baxter, V.V. Bazhanov and J.H.H. Perk, *Functional relations for transfer matrices of the chiral Potts model*, *Int. J. Mod. Phys. B* **4** (1990) 803 [[INSPIRE](#)].
- [5] G. von Gehlen and V. Rittenberg,  *$Z(n)$  Symmetric Quantum Chains With an Infinite Set of Conserved Charges and  $Z(n)$  Zero Modes*, *Nucl. Phys. B* **257** (1985) 351 [[INSPIRE](#)].
- [6] F.C. Alcaraz and A. Lima Santos, *Conservation Laws for  $Z(N)$  Symmetric Quantum Spin Models and Their Exact Ground State Energies*, *Nucl. Phys. B* **275** (1986) 436 [[INSPIRE](#)].
- [7] Yu. A. Bashilov and S.V. Pokrovsky, *Conservation Laws In The Quantum Version Of  $N$  Positional Potts Model*, *Commun. Math. Phys.* **76** (1980) 129 [[INSPIRE](#)].
- [8] H. Au-Yang, B.M. McCoy, J.H.H. Perk, S. Tang and M.-L. Yan, *Commuting transfer matrices in the chiral Potts models: Solutions of Star triangle equations with genus  $> 1$* , *Phys. Lett. A* **123** (1987) 219 [[INSPIRE](#)].
- [9] B.M. McCoy, J.H.H. Perk, S. Tang and C.-H. Sah, *Commuting Transfer Matrices for the Four State Selfdual Chiral Potts Model With a Genus Three Uniformizing Fermat Curve*, *Phys. Lett. A* **125** (1987) 9 [[INSPIRE](#)].
- [10] R.J. Baxter, J.H.H. Perk and H. Au-Yang, *New solutions of the star triangle relations for the chiral Potts model*, *Phys. Lett. A* **128** (1988) 138 [[INSPIRE](#)].
- [11] S.N. Ruijsenaars, *Relativistic Toda system*, *Commun. Math. Phys.* **133** (1990) 217.
- [12] G. Albertini, B.M. McCoy and J.H.H. Perk, *Eigenvalue Spectrum Of The Superintegrable Chiral Potts Model*, *Adv. Stud. Pure Math.* **19** (1989) 1 [[INSPIRE](#)].
- [13] R.J. Baxter, *The order parameter of the chiral Potts model*, *J. Statist. Phys.* **120** (2005) 1 [[cond-mat/0501226](#)] [[INSPIRE](#)].
- [14] R.J. Baxter, *Free energy of the solvable chiral Potts model*, *J. Statist. Phys.* **52** (1988) 639 [[INSPIRE](#)].
- [15] P. Fendley, *Free parafermions*, *J. Phys. A* **47** (2014) 075001 [[arXiv:1310.6049](#)] [[INSPIRE](#)].

- [16] R.J. Baxter, *The  $\tau_2$  model and parafermions*, *J. Phys. A* **47** (2014) 315001 [[arXiv:1310.7074](#)] [[INSPIRE](#)].
- [17] H. Au-Yang and J.H.H. Perk, *Parafermions in the tau(2) model*, *J. Phys. A* **47** (2014) 315002 [[INSPIRE](#)].
- [18] H. Au-Yang and J.H.H. Perk, *Parafermions in the tau-2 model II*, [arXiv:1606.06319](#).
- [19] J. Cao, W. Yang, K. Shi and Y. Wang, *Off-diagonal Bethe ansatz and exact solution of a topological spin ring*, *Phys. Rev. Lett.* **111** (2013) 137201 [[arXiv:1305.7328](#)] [[INSPIRE](#)].
- [20] J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Off-diagonal Bethe ansatz solution of the XXX spin-chain with arbitrary boundary conditions*, *Nucl. Phys. B* **875** (2013) 152 [[arXiv:1306.1742](#)] [[INSPIRE](#)].
- [21] J. Cao, W.-L. Yang, S. Cui, K.-J. Shi and Y. Wang, *Spin- $\frac{1}{2}$  XYZ model revisit: General solutions via off-diagonal Bethe ansatz*, *Nucl. Phys. B* **886** (2014) 185 [[arXiv:1307.0280](#)] [[INSPIRE](#)].
- [22] J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Off-diagonal Bethe ansatz solutions of the anisotropic spin-1/2 chains with arbitrary boundary fields*, *Nucl. Phys. B* **877** (2013) 152 [[arXiv:1307.2023](#)] [[INSPIRE](#)].
- [23] Y. Wang, W.-L. Yang, J. Cao and K. Shi, *Off-Diagonal Bethe Ansatz for Exactly Solvable Models*, Springer, (2015).
- [24] X. Xu, J. Cao, S. Cui, W.-L. Yang, K. Shi and Y. Wang, *Off-diagonal Bethe Ansatz solution of the  $\tau_2$ -model*, *JHEP* **09** (2015) 212 [[arXiv:1507.03367](#)] [[INSPIRE](#)].
- [25] X. Zhang, J. Cao, W.-L. Yang and K.-J. Shi, *Bethe states of the  $\tau_2$ -model with the periodic boundary condition*.
- [26] P.P. Kulish, N.Yu. Reshetikhin and E.K. Sklyanin, *Yang-Baxter Equation and Representation Theory. 1.*, *Lett. Math. Phys.* **5** (1981) 393 [[INSPIRE](#)].
- [27] P.P. Kulish and E.K. Sklyanin, *Quantum spectral transform method. Recent developments*, *Lect. Notes Phys.* **151** (1982) 61 [[INSPIRE](#)].
- [28] A.N. Kirillov and N.Y. Reshetikhin, *Exact solution of the integrable XXZ Heisenberg model with arbitrary spin. I. The ground state and the excitation spectrum*, *J. Phys. A* **20** (1987) 1565 [[INSPIRE](#)].
- [29] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, (1982).
- [30] V.E. Korepin, N.M. Bogoliubov and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Function*, Cambridge University Press, (1993).
- [31] E.K. Sklyanin, *Boundary Conditions for Integrable Quantum Systems*, *J. Phys. A* **21** (1988) 2375 [[INSPIRE](#)].
- [32] I.V. Cherednik, *Factorizing Particles on a Half Line and Root Systems*, *Theor. Math. Phys.* **61** (1984) 977 [[INSPIRE](#)].
- [33] H.J. de Vega and A. González-Ruiz, *Boundary K matrices for the XYZ, XXZ and XXX spin chains*, *J. Phys. A* **27** (1994) 6129 [[hep-th/9306089](#)] [[INSPIRE](#)].
- [34] S. Ghoshal and A.B. Zamolodchikov, *Boundary S matrix and boundary state in two-dimensional integrable quantum field theory*, *Int. J. Mod. Phys. A* **9** (1994) 3841 [*Erratum ibid. A* **9** (1994) 4353] [[hep-th/9306002](#)] [[INSPIRE](#)].

- [35] L. Mezincescu and R.I. Nepomechie, *Analytical Bethe Ansatz for quantum algebra invariant spin chains*, *Nucl. Phys. B* **372** (1992) 597 [[hep-th/9110050](#)] [[INSPIRE](#)].
- [36] R.E. Behrend, P.A. Pearce and D.L. O'Brien, *Interaction-round-a-face models with fixed boundary conditions: The ABF fusion hierarchy*, *J. Statist. Phys.* **84** (1996) 1 [[hep-th/9507118](#)] [[INSPIRE](#)].
- [37] W.-L. Yang and Y.-Z. Zhang, *T-Q relation and exact solution for the XYZ chain with general nondiagonal boundary terms*, *Nucl. Phys. B* **744** (2006) 312 [[hep-th/0512154](#)] [[INSPIRE](#)].
- [38] V.O. Tarasov, *Cyclic Monodromy Matrices For The R-Matrix Of The Six-Vertex Model And The Chiral Potts Model With Fixed Spin Boundary Conditions*, *Int. J. Mod. Phys. A* **07** (1992) 963.
- [39] L. Mezincescu, R.I. Nepomechie and V. Rittenberg, *Bethe Ansatz Solution of the Fateev-zamolodchikov Quantum Spin Chain With Boundary Terms*, *Phys. Lett. A* **147** (1990) 70 [[INSPIRE](#)].
- [40] L. Mezincescu and R.I. Nepomechie, *Fusion procedure for open chains*, *J. Phys. A* **25** (1992) 2533 [[INSPIRE](#)].
- [41] Y.-k. Zhou, *Row transfer matrix functional relations for Baxter's eight vertex and six vertex models with open boundaries via more general reflection matrices*, *Nucl. Phys. B* **458** (1996) 504 [[hep-th/9510095](#)] [[INSPIRE](#)].
- [42] T. Inami, S. Odake and Y.-Z. Zhang, *Reflection K matrices of the 19 vertex model and XXZ spin 1 chain with general boundary terms*, *Nucl. Phys. B* **470** (1996) 419 [[hep-th/9601049](#)] [[INSPIRE](#)].
- [43] J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Nested off-diagonal Bethe ansatz and exact solutions of the SU( $N$ ) spin chain with generic integrable boundaries*, *JHEP* **04** (2014) 143 [[arXiv:1312.4770](#)] [[INSPIRE](#)].
- [44] K. Hao, J. Cao, T. Yang and W.-L. Yang, *Exact solution of the XXX Gaudin model with generic open boundaries*, *Annals Phys.* **354** (2015) 401 [[arXiv:1408.3012](#)] [[INSPIRE](#)].
- [45] L. Frappat, R. Nepomechie and É. Ragoucy, *A Complete Bethe Ansatz solution of the open spin-s XXZ chain with general integrable boundary terms*, *J. Stat. Mech.* **0709** (2007) P09009 [[arXiv:0707.0653](#)] [[INSPIRE](#)].
- [46] A.G. Izergin and V.E. Korepin, *A Lattice model related to the nonlinear Schrödinger equation*, *Dokl. Akad. Nauk Ser. Fiz.* **259** (1981) 76 [[INSPIRE](#)].
- [47] P.P. Kulish and E.K. Sklyanin, *Quantum spectral transform method. Recent developments*, *Lect. Notes Phys.* **151** (1982) 61 [[INSPIRE](#)].
- [48] A.G. Izergin and V.E. Korepin, *Lattice versions of quantum field theory models in two-dimensions*, *Nucl. Phys. B* **205** (1982) 401 [[INSPIRE](#)].
- [49] V.V. Bazhanov and N.Yu. Reshetikhin, *Critical Rsos Models and Conformal Field Theory*, *Int. J. Mod. Phys. A* **4** (1989) 115 [[INSPIRE](#)].
- [50] G. von Gehlen, N. Iorgov, S. Pakuliak and V. Shadura, *The Baxter-Bazhanov-Stroganov model: separation of variables and the Baxter equation*, *J. Phys. A* **39** (2006) 7257 [[nlin/0603028](#)].
- [51] R.I. Nepomechie, *Bethe Ansatz solution of the open XX spin chain with nondiagonal boundary terms*, *J. Phys. A* **34** (2001) 9993 [[hep-th/0110081](#)] [[INSPIRE](#)].

- [52] R.I. Nepomechie, *Solving the open XXZ spin chain with nondiagonal boundary terms at roots of unity*, *Nucl. Phys. B* **622** (2002) 615 [[hep-th/0110116](#)] [[INSPIRE](#)].
- [53] R.I. Nepomechie, *Functional relations and Bethe Ansatz for the XXZ chain*, *J. Statist. Phys.* **111** (2003) 1363 [[hep-th/0211001](#)] [[INSPIRE](#)].
- [54] R.I. Nepomechie, *Bethe ansatz solution of the open XXZ chain with nondiagonal boundary terms*, *J. Phys. A* **37** (2004) 433 [[hep-th/0304092](#)] [[INSPIRE](#)].
- [55] J. Cao, H.-Q. Lin, K.-J. Shi and Y. Wang, *Exact solution of XXZ spin chain with unparallel boundary fields*, *Nucl. Phys. B* **663** (2003) 487 [[INSPIRE](#)].
- [56] W.-L. Yang, R.I. Nepomechie and Y.-Z. Zhang, *Q-operator and T-Q relation from the fusion hierarchy*, *Phys. Lett. B* **633** (2006) 664 [[hep-th/0511134](#)] [[INSPIRE](#)].
- [57] J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Exact solution of the alternating XXZ spin chain with generic non-diagonal boundaries*, *Annals Phys.* **361** (2015) 91 [[arXiv:1409.3646](#)] [[INSPIRE](#)].
- [58] J.M. Maillet, G. Niccoli and B. Pezelier, *Transfer matrix spectrum for cyclic representations of the 6-vertex reflection algebra I*, [arXiv:1607.02983](#) [[INSPIRE](#)].
- [59] E.K. Sklyanin, *The Quantum Toda Chain*, *Lect. Notes Phys.* **226** (1985) 196 [[INSPIRE](#)].
- [60] E.K. Sklyanin, *Goryachev-Chaplygin top and the inverse scattering method*, *J. Sov. Math.* **31** (1985) 3417 [[INSPIRE](#)].
- [61] E.K. Sklyanin, *Separation of variables - new trends*, *Prog. Theor. Phys. Suppl.* **118** (1995) 35 [[solv-int/9504001](#)] [[INSPIRE](#)].