Some generalization of Cauchy's and the quadratic functional equations

RADOSŁAW ŁUKASIK

Abstract. We find the solutions $f, g, h: S \rightarrow H$ of each of the functional equations

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = |\Lambda|g(x) + h(y), \quad x, y \in S,$$

where $(S, +)$ is an abelian semigroup, Λ is a finite subgroup of the automorphism group of S , $(H, +)$ is an abelian group.

Mathematics Subject Classification (2010). 39B52.

Keywords. Cauchy functional equation, quadratic functional equation.

1. Introduction

The generalization of the quadratic functional equation

$$f(x + y) + f(x + \sigma y) = 2f(x) + 2f(y), \quad x, y \in G,$$

where σ is an automorphism of the abelian group G such that $\sigma^2 = id_G$, $f, g: G \rightarrow \mathbb{C}$ was investigated by Stetkær [3].

In Stetkær [4] solved the functional equation

$$\frac{1}{N} \sum_{n=0}^{N-1} f(z + \omega^n \zeta) = g(z) + h(\zeta), \quad z, \zeta \in \mathbb{C},$$

where $N \in \{2, 3, \dots\}$ and ω is a primitive N th root of unity, $f, g, h: \mathbb{C} \rightarrow \mathbb{C}$ are continuous.

In the present paper we give the complete solution of the following functional equation

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = Lg(x) + h(y), \quad x, y \in S,$$

where $(S, +)$ is an abelian semigroup, Λ is a finite subgroup of the automorphism group of S , $L = \text{card}\Lambda$, $(H, +)$ is an abelian group uniquely divisible by $(L + 1)!$, $f, g, h: S \rightarrow H$.

2. Main result

In this work we use a theorem and a corollary proved by Mazur and Orlicz [2], and generalized by Djoković [1]:

Theorem 1. *Let $(S, +)$ be an abelian semigroup, $n \in \mathbb{N}$, $(H, +)$ be an abelian group uniquely divisible by $(n + 1)!$, $f: S \rightarrow H$ satisfies the equation*

$$\Delta_v^{n+1} f(u) = 0, \quad u, v \in S.$$

Then there exist $A_0 \in H$ and k -additive, symmetric mappings $A_k: S \rightarrow H$, $k \in \{1, \dots, n\}$ such that

$$f(x) = A_0 + A_1(x) + \dots + A_n(x, \dots, x), \quad x \in S.$$

Corollary 1. *Let $(S, +)$ be an abelian semigroup, $n \in \mathbb{N}$, $(H, +)$ be an abelian group uniquely divisible by $(n + 1)!$, $f, g: S \rightarrow H$ satisfy the equation*

$$\Delta_v^n f(u) = g(v), \quad u, v \in S.$$

Then there exist $A_0 \in H$ and k -additive, symmetric mappings $A_k: S \rightarrow H$, $k \in \{1, \dots, n\}$ such that

$$\begin{aligned} f(x) &= A_0 + A_1(x) + \dots + A_n(x, \dots, x), \quad x \in S, \\ g(x) &= n! A_n(x, \dots, x), \quad x \in S. \end{aligned}$$

We start with some lemmas.

Lemma 1. *For all $n \in \mathbb{N}$ we have*

$$\sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^i = 0, \quad i \in \{1, \dots, n-1\}, \quad n \neq 1, \quad (1)$$

$$\sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^n = n!. \quad (2)$$

Proof. First we prove (1) by induction on n .

For $n = 2$ we have

$$\sum_{k=1}^2 (-1)^{2-k} \binom{2}{k} k^i = -2 \cdot 1 + 1 \cdot 2 = 0.$$

Assume that (1) holds for $2, \dots, n$. Since

$$k \binom{n+1}{k} = (n+1) \binom{n}{k-1}, \quad k \in \{1, \dots, n\},$$

we have

$$\begin{aligned}\sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n+1}{k} k &= (n+1) \sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n}{k-1} \\ &= (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} = 0.\end{aligned}$$

Hence, for $i \in \{1, \dots, n-1\}$,

$$\begin{aligned}\sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n+1}{k} k^{i+1} &= (n+1) \sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n}{k-1} k^i \\ &= (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} (k+1)^i = (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \sum_{m=0}^i \binom{i}{m} k^m \\ &= (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} + (n+1) \sum_{m=1}^i \binom{i}{m} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^m \\ &= (n+1) \sum_{m=1}^i \binom{i}{m} \sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^m = 0.\end{aligned}$$

Now we prove (2) by induction on n .

For $n = 1$ we have

$$\sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^n = 1.$$

Assume that (2) holds for some $n \in \mathbb{N}$. Using (1) we get

$$\begin{aligned}\sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n+1}{k} k^{n+1} &= (n+1) \sum_{k=1}^{n+1} (-1)^{n+1-k} \binom{n}{k-1} k^n \\ &= (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} (k+1)^n = (n+1) \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \sum_{m=0}^n \binom{n}{m} k^m \\ &= (n+1) \sum_{m=0}^n \binom{n}{m} \sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^m = (n+1) \sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^n \\ &= (n+1)!.\end{aligned}$$

□

Lemma 2. *Let $(H, +)$ be an abelian group uniquely divisible by $n!$, $x_1, \dots, x_n \in H$ be such that*

$$\sum_{i=1}^n k^i x_i = 0, \quad k \in \{1, \dots, n\}.$$

Then $x_1 = \dots = x_n = 0$.

Proof. First, we notice that for the group $(H, +)$ from the unique divisibility by $n!$ we obtain the unique divisibility by $k!$ for each $k \in \{1, \dots, n\}$.

We prove this theorem by induction on n .

For $n = 1$ we have $x_1 = 0$. Assume that the theorem holds for $n - 1, n > 1$. In view of Lemma 1 we have

$$0 = \sum_{k=1}^n (-1)^{n-k} \binom{n}{k} \sum_{i=1}^n k^i x_i = \sum_{i=1}^n \left[\sum_{k=1}^n (-1)^{n-k} \binom{n}{k} k^i \right] x_i = n! x_n.$$

Hence $x_n = 0$ and

$$\sum_{k=1}^{n-1} k^i x_i = 0.$$

By the induction hypothesis we obtain

$$x_1 = \dots = x_{n-1} = 0 = x_n,$$

which finishes the proof. $\square$

Lemma 3. *Let $(S, +)$ be an abelian semigroup, Λ be a finite subgroup of the automorphism group of $S, L := \text{card} \Lambda, (H, +)$ be an abelian group uniquely divisible by $L!$. Further, let for each $k \in \{1, \dots, L\} A_k: S \rightarrow H$ be k -additive and symmetric mappings, $f: S \rightarrow H$ be a function given by the formula*

$$f(x) = A_1(x) + \dots + A_L(x, \dots, x), \quad x \in S.$$

Then a function f satisfies the equation

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = Lf(x) + \sum_{\lambda \in \Lambda} f(\lambda y), \quad x, y \in S \quad (3)$$

if and only if

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L.$$

Proof. First we show that for $x, y \in S$

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) - \sum_{\lambda \in \Lambda} f(\lambda y) - Lf(x) = \sum_{\lambda \in \Lambda} \sum_{i=1}^{L-1} \sum_{k=i+1}^L \binom{k}{i} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i).$$

Indeed, we have the following sequence of identities

$$\begin{aligned} & \sum_{\lambda \in \Lambda} f(x + \lambda y) - \sum_{\lambda \in \Lambda} f(\lambda y) - Lf(x) \\ &= \sum_{\lambda \in \Lambda} \sum_{k=1}^L [A_k(x + \lambda y, \dots, x + \lambda y) - A_k(\lambda y, \dots, \lambda y) - A_k(x, \dots, x)] \\ &= \sum_{\lambda \in \Lambda} \sum_{k=2}^L \sum_{i=1}^{k-1} \binom{k}{i} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) \\ &= \sum_{\lambda \in \Lambda} \sum_{i=1}^{L-1} \sum_{k=i+1}^L \binom{k}{i} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i), \quad x, y \in S. \end{aligned}$$

By the above equality we obtain that if

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 1 \leq k \leq L,$$

then f satisfies (3).

Now we assume that f satisfies (3). Then

$$\sum_{\lambda \in \Lambda} \sum_{i=1}^{L-1} \sum_{k=i+1}^L \binom{k}{i} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0.$$

For each $i \in \{1, \dots, L-1\}$ we define $g_i: S \times S \rightarrow H$ by the formula

$$g_i(x, y) := \sum_{\lambda \in \Lambda} \sum_{k=i+1}^L \binom{k}{i} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i), \quad x, y \in S.$$

Since

$$g_i(x, my) = m^i g_i(x, y), \quad m \in \mathbb{N}, \quad x, y \in S, \quad 1 \leq i \leq L-1,$$

then

$$\sum_{i=1}^{L-1} m^i g_i(x, y) = \sum_{i=1}^{L-1} g_i(x, my) = 0, \quad m \in \mathbb{N}, \quad x, y \in S.$$

In view of Lemma 2 we obtain

$$g_i(x, y) = 0, \quad x, y \in S, \quad i \in \{1, \dots, L-1\}.$$

Now, for each $i \in \{1, \dots, L-1\}, j \in \{1, \dots, L-i\}$, we define $h_{i,j}: S \times S \rightarrow H$ by the formula

$$h_{i,j} := \binom{i+j}{i} \sum_{\lambda \in \Lambda} A_{i+j}(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i), \quad x, y \in S.$$

Since

$$h_{i,j}(mx, y) = m^j h_{i,j}(x, y), \quad m \in \mathbb{N}, \quad x, y \in S, \quad 1 \leq j \leq L - i, \quad 1 \leq i \leq L - 1,$$

we have

$$\sum_{j=1}^{L-i} m^j h_{i,j}(x, y) = \sum_{j=1}^{L-i} h_{i,j}(mx, y) = g_i(mx, y) = 0,$$

for $m \in \mathbb{N}$, $x, y \in S$, $1 \leq i \leq L - 1$. In view of lemma 2 we obtain

$$h_{i,j}(x, y) = 0, \quad x, y \in S, \quad 1 \leq j \leq L - i, \quad 1 \leq i \leq L - 1,$$

so we get

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k - 1, \quad 2 \leq k \leq L.$$

□

Theorem 2. Let $(S, +)$ be an abelian semigroup, Λ be a finite subgroup of the automorphism group of S , $L := \text{card} \Lambda$, $(H, +)$ be an abelian group. Further, let $f: S \rightarrow H$ satisfy Eq. (3), $g: S \rightarrow H$ be a function given by

$$g(x) := - \sum_{i=0}^{L-1} (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} f \left(\sum_{\mu \in \Lambda_{i,j}} \mu x \right), \quad x \in S,$$

where $\Lambda_{i,j} \subset \Lambda$ are pairwise different sets such that $\text{card} \Lambda_{i,j} = L - i$ for $j \in \{1, \dots, \binom{L}{i}\}$, $i \in \{0, \dots, L\}$. Then

$$L \Delta_v^L f(u) = Lg(v), \quad u, v \in S.$$

Proof. First we observe that any solution of Eq. (3) in a semigroup without zero can be uniquely extended to a solution in a semigroup with a neutral element 0 by putting $f(0) = 0$. So we can assume that $(S, +)$ is a monoid.

Since

$$\lambda \Lambda_{i,j} = \Lambda_{i,k} \Rightarrow \lambda^{-1} \Lambda_{i,k} = \Lambda_{i,j}, \quad \lambda \in \Lambda, i \in \{0, \dots, L\}, j, k \in \left\{1, \dots, \binom{L}{i}\right\},$$

we have

$$\sum_{j=1}^{\binom{L}{i}} \sum_{\lambda \in \Lambda} f \left(\sum_{\mu \in \Lambda_{i,j}} \lambda \mu x \right) = L \sum_{j=1}^{\binom{L}{i}} f \left(\sum_{\mu \in \Lambda_{i,j}} \mu x \right), \quad x \in S. \quad (4)$$

Fix $u, v \in S$. Let

$$x_i := u + iv, \quad y_{i,j} := \sum_{\mu \in \Lambda_{i,j}} \mu v, \quad j \in \left\{1, \dots, \binom{L}{i}\right\}, \quad i \in \{0, \dots, L\}.$$

For $\lambda \in \Lambda, i \in \{0, \dots, L\}, j \in \{1, \dots, \binom{L}{i}\}$ we consider two cases:

- (i) $\lambda^{-1} \in \Lambda_{i,j}$. Hence $i \neq L$. Let $k \in \{1, \dots, \binom{L}{i+1}\}$ be such that $\Lambda_{i,j} = \Lambda_{i+1,k} \cup \{\lambda^{-1}\}$. Then

$$\begin{aligned} x_i + \lambda y_{i,j} &= u + iv + \sum_{\mu \in \Lambda_{i,j}} \lambda \mu v = u + (i+1)v + \sum_{\mu \in \Lambda_{i+1,k}} \lambda \mu v \\ &= x_{i+1} + \lambda y_{i+1,k}. \end{aligned}$$

- (ii) $\lambda^{-1} \notin \Lambda_{i,j}$. Hence $i \neq 0$. Let $k \in \{1, \dots, \binom{L}{i-1}\}$ be such that $\Lambda_{i-1,k} = \Lambda_{i,j} \cup \{\lambda^{-1}\}$. Then

$$\begin{aligned} x_i + \lambda y_{i,j} &= u + iv + \sum_{\mu \in \Lambda_{i,j}} \lambda \mu v = u + (i-1)v + \sum_{\mu \in \Lambda_{i-1,k}} \lambda \mu v \\ &= x_{i-1} + \lambda y_{i-1,k}. \end{aligned}$$

From the above consideration and equality (4) we obtain

$$\begin{aligned} 0 &= \sum_{i=0}^L (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} \sum_{\lambda \in \Lambda} f(x_i + \lambda y_{i,j}) \\ &= \sum_{i=0}^L (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} \left[Lf(x_i) + \sum_{\lambda \in \Lambda} f(\lambda y_{i,j}) \right] \\ &= L \sum_{i=0}^L (-1)^{L-i} \binom{L}{i} f(u + iv) + \sum_{i=0}^{L-1} (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} \sum_{\lambda \in \Lambda} f \left(\sum_{\mu \in \Lambda_{i,j}} \lambda \mu v \right) \\ &= L \sum_{i=0}^L (-1)^{L-i} \binom{L}{i} f(u + iv) + L \sum_{i=0}^{L-1} (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} f \left(\sum_{\mu \in \Lambda_{i,j}} \lambda \mu v \right) \\ &= L \Delta_v^L f(u) - Lg(v). \end{aligned}$$

□

Theorem 3. Let $(S, +)$ be an abelian semigroup, Λ be a finite subgroup of the automorphism group of $S, L := \text{card} \Lambda, (H, +)$ be an abelian group uniquely divisible by $(L+1)!$. Then a function $f: S \rightarrow H$ satisfies Eq. (3) if and only if there exist k -additive and symmetric mappings $A_k: S^k \rightarrow H, k \in \{1, \dots, L\}$ such that

$$f(x) = A_1(x) + \cdots + A_L(x, \dots, x), \quad x \in S,$$

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L.$$

Proof. The proof of this theorem follows straight from Theorem 2, Corollary 1 and Lemma 3. $\square$

Now we show some applications of the above theorems.

Theorem 4. *Let $(S, +)$ be an abelian semigroup, Λ be a finite subgroup of the automorphism group of S , $L := \text{card} \Lambda$, $(H, +)$ be an abelian group uniquely divisible by $(L+1)!$. Then a function $f: S \rightarrow H$ satisfies the equation*

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = Lf(x) + Lf(y), \quad x, y \in S \quad (5)$$

if and only if there exist k -additive, symmetric mappings $A_k: S^k \rightarrow H$, $k \in \{1, \dots, L\}$ such that

$$f(x) = A_1(x) + \cdots + A_L(x, \dots, x), \quad x \in S,$$

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L,$$

$$A_k(\mu x, \dots, \mu x) = A_k(x, \dots, x), \quad x \in S, \quad \mu \in \Lambda, \quad 1 \leq k \leq L.$$

Proof. It is easy to check that if f satisfies the three above conditions then f satisfies Eq. (5). We observe that if f satisfies Eq. (5), then f satisfies Eq. 3 and $f(\mu x) = f(x)$, $\mu \in \Lambda$, $x \in S$.

Indeed, for $x, y \in S$, $\mu \in \Lambda$ we have

$$Lf(y) = -Lf(x) + \sum_{\lambda \in \Lambda} f(x + \lambda y) = -Lf(x) + \sum_{\lambda \in \Lambda} f(x + \lambda \mu y) = Lf(\mu y),$$

which gives $f(\mu x) = f(x)$. Hence $Lf(y) = \sum_{\lambda \in \Lambda} f(\lambda y)$ for $y \in S$, thus f satisfies Eq. (3). In view of Theorem 3 we obtain the first and the second condition of the thesis. Finally, we show that

$$A_k(\mu x, \dots, \mu x) = A_k(x, \dots, x), \quad x \in S, \quad \mu \in \Lambda, \quad 1 \leq k \leq L.$$

Fix $\mu \in \Lambda$. We observe that

$$0 = f(\mu x) - f(x) = \sum_{k=1}^L [A_k(\mu x, \dots, \mu x) - A_k(x, \dots, x)], \quad x \in S.$$

For each $i \in \{1, \dots, L\}$ we define $g_i: S \rightarrow H$ by the formula

$$g_i(x) := A_i(\mu x, \dots, \mu x) - A_i(x, \dots, x), \quad x \in S.$$

Since

$$g_i(mx) = m^i g_i(x), \quad m \in \mathbb{N}, \quad x \in S,$$

we have

$$0 = \sum_{i=1}^L g_i(mx) = \sum_{i=1}^L m^i g_i(x), \quad m \in \mathbb{N}, \quad x \in S.$$

Hence, in view of Lemma 2, we obtain

$$g_i(x) = 0, \quad 1 \leq i \leq L, \quad x \in S,$$

which ends the proof. $\square$

Theorem 5. Let $(S, +)$ be an abelian semigroup, Λ be a finite subgroup of the automorphism group of S , $L := \text{card} \Lambda$, $(H, +)$ be an abelian group uniquely divisible by $L!$. Then a function $f: S \rightarrow H$ satisfies the equation

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = Lf(x), \quad x, y \in S \quad (6)$$

if and only if there exist k -additive, symmetric mappings $A_k: S^k \rightarrow H$, $k \in \{1, \dots, L-1\}$ and $A_0 \in H$ such that

$$f(x) = A_0 + A_1(x) + \dots + A_{L-1}(x, \dots, x), \quad x \in S,$$

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k, \quad 2 \leq k \leq L-1.$$

Proof. Let us note that if a function f satisfies the above conditions then f satisfies Eq. (6).

First we observe that if $(S, +)$ is a semigroup without a neutral element 0, then we can uniquely extend each solution of (6) to a solution on the semigroup $(S \cup \{0\}, +)$. Indeed, we have

$$\begin{aligned} L \sum_{\lambda \in \Lambda} f(\lambda y) &= \sum_{\lambda \in \Lambda} \sum_{\mu \in \Lambda} f(\lambda y + \mu x) \\ &= \sum_{\mu \in \Lambda} \sum_{\lambda \in \Lambda} f(\mu x + \lambda y) = L \sum_{\mu \in \Lambda} f(\mu x), \quad x, y \in S, \end{aligned}$$

and

$$Lf \left(\sum_{\lambda \in \Lambda} \lambda y \right) = \sum_{\mu \in \Lambda} f \left(\mu \sum_{\lambda \in \Lambda} \lambda y \right) = \sum_{\mu \in \Lambda} f(\mu x), \quad x, y \in S.$$

Putting $f(0) := f(\sum_{\lambda \in \Lambda} \lambda y)$ for $y \in S$. It is easy to check that the solution extended in this way satisfies Eq. 3 for the semigroup S with zero.

So we can assume that $(S, +)$ is a monoid. Since

$$Lf(0) = \sum_{\lambda \in \Lambda} f(\lambda y), \quad y \in S,$$

then $f_0 := f - f(0)$ satisfies Eq. 3. Let $\Lambda_{i,j} \subset \Lambda, j \in \{1, \dots, \binom{L}{i}\}, i \in \{0, \dots, L\}$, be sets from Theorem 2. We observe that for $x \in S$ we have

$$\sum_{\lambda \in \Lambda} f_0 \left(\sum_{\mu \in \Lambda_{i,j}} \lambda \mu x \right) = \sum_{\lambda \in \Lambda} f \left(\lambda \sum_{\mu \in \Lambda_{i,j}} \mu x \right) - Lf(0) = 0.$$

By equality (4) we obtain

$$L \sum_{i=0}^{L-1} (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} f_0 \left(\sum_{\mu \in \Lambda_{i,j}} \mu x \right) = \sum_{i=0}^{L-1} (-1)^{L-i} \sum_{j=1}^{\binom{L}{i}} \sum_{\lambda \in \Lambda} f_0 \left(\sum_{\mu \in \Lambda_{i,j}} \lambda \mu x \right) = 0.$$

Then Theorem 2 holds with $g = 0$. In view of Theorem 1, Lemma 3 and putting $A_0 := f(0)$ we obtain that there exist k -additive and symmetric mappings $A_k: S^k \rightarrow H, k \in \{1, \dots, L-1\}$ such that

$$f(x) = A_0 + A_1(x) + \dots + A_{L-1}(x, \dots, x), \quad x \in S,$$

$$\binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) = 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L-1,$$

Finally, we show that

$$\sum_{\lambda \in \Lambda} A_k(\lambda x, \dots, \lambda x) = 0, \quad x \in S, \quad 2 \leq k \leq L-1.$$

For each $i \in \{1, \dots, L-1\}$ we define $g_i: S \rightarrow H$ by the formula

$$g_i(x) := \sum_{\lambda \in \Lambda} A_i(\lambda x, \dots, \lambda x), \quad x \in S.$$

Since

$$g_i(mx) = m^i g_i(x), \quad m \in \mathbb{N}, \quad x \in S, \quad 1 \leq i \leq L-1,$$

we have

$$0 = \sum_{\lambda \in \Lambda} f(\lambda x) - Lf(0) = \sum_{i=1}^L g_i(mx) = \sum_{i=1}^L m^i g_i(x), \quad m \in \mathbb{N}, \quad x \in S.$$

In view of Lemma 2

$$g_i(x) = 0, \quad 1 \leq i \leq L-1, \quad x \in S,$$

which ends the proof. $\square$

Our main result reads as follows

Theorem 6. *Let $(S, +)$ be an abelian monoid, Λ be a finite subgroup of the automorphism group of $S, L := \text{card} \Lambda, (H, +)$ be an abelian group uniquely divisible by $(L+1)!$. Then functions $f, g, h: S \rightarrow H$ satisfy the equation*

$$\sum_{\lambda \in \Lambda} f(x + \lambda y) = Lg(x) + h(y), \quad x, y \in S \quad (7)$$

if and only if there exist k -additive, symmetric mappings $A_k: S^k \rightarrow H$, $k \in \{1, \dots, L\}$ and $A_0, B_0 \in H$ such that

$$\begin{aligned} f(x) &= A_0 + A_1(x) + \dots + A_L(x, \dots, x), \quad x \in S, \\ g(x) &= B_0 + A_1(x) + \dots + A_L(x, \dots, x), \quad x \in S, \\ h(x) &= LA_0 - LB_0 + \sum_{\lambda \in \Lambda} A_1(\lambda x) + \dots + \sum_{\lambda \in \Lambda} A_L(\lambda x, \dots, \lambda x), \quad x \in S, \\ \binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) &= 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L. \end{aligned}$$

Proof. It is easy to check that if the four equalities above hold, then functions f, g, h satisfy Eq. (7).

Assume that functions f, g, h satisfy Eq. (7). First we observe that

$$\begin{aligned} Lf(0) &= Lg(0) + h(0), \\ Lf(x) &= Lg(x) + h(0) = Lg(x) - Lg(0) + Lf(0), \quad x \in S. \end{aligned}$$

Therefore, we get

$$f(x) = g(x) - g(0) + f(0), \quad x \in S, \quad (8)$$

and

$$Lg(0) + h(y) = \sum_{\lambda \in \Lambda} f(\lambda y) = \sum_{\lambda \in \Lambda} [g(\lambda y) - g(0)] + Lf(0), \quad y \in S,$$

hence

$$h(y) = \sum_{\lambda \in \Lambda} [g(\lambda y) - g(0)] + Lf(0) - Lg(0), \quad y \in S. \quad (9)$$

Let $g_0 := g - g(0)$. From the above equalities we have

$$\begin{aligned} \sum_{\lambda \in \Lambda} g_0(x + \lambda y) &= \sum_{\lambda \in \Lambda} [g(x + \lambda y) - g(0) + f(0)] - Lf(0) \\ &= \sum_{\lambda \in \Lambda} f(x + \lambda y) - Lf(0) = Lg(x) + h(y) - Lf(0) \\ &= Lg_0(x) + \sum_{\lambda \in \Lambda} g_0(\lambda y), \quad x, y \in S. \end{aligned}$$

In view of Theorem 3, there exist k -additive and symmetric mappings $A_k: S^k \rightarrow H, k \in \{1, \dots, L\}$ such that

$$\begin{aligned} g_0(x) &= A_1(x) + \dots + A_L(x, \dots, x), \quad x \in S, \\ \binom{k}{i} \sum_{\lambda \in \Lambda} A_k(x, \dots, x, \underbrace{\lambda y, \dots, \lambda y}_i) &= 0, \quad x, y \in S, \quad 1 \leq i \leq k-1, \quad 2 \leq k \leq L. \end{aligned}$$

Let $B_0 := g(0), A_0 := f(0)$. Then from equalities 8 and 9 we obtain the thesis. $\square$

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Djoković, D.Ž.: A representation theorem for $(X_{1-1})(X_{2-1})\cdots(X_{n-1})$ and its applications. *Ann. Polon. Math.* **22**, 189–198 (1969)
- [2] Mazur, S., Orlicz, W.: Grundlegende Eigenschaften der Polynomischen Operationen I, II, *Studia Math.* **5**, 50–68, 179–189 (1934)
- [3] Stetkær, H.: Functional equation on abelian groups with involution, II. *Aequat. Math.* **55**, 227–240 (1998)
- [4] Stetkær, H.: Functional equations involving means of functions on the complex plane. *Aequat. Math.* **56**, 47–62 (1998)

Radosław Łukasik
Institute of Mathematics
University of Silesia
Bankowa 14
40-007 Katowice
Poland
e-mail: rlukasik@math.us.edu.pl

Received: November 12, 2010

Revised: November 9, 2011