Around a problem of Nicole Brillouët-Belluot, II

JANUSZ MORAWIEC

Abstract. For every $\alpha \in \mathbb{R}$ we determine all increasing bijections $f: (0, +\infty) \rightarrow (0, +\infty)$ such that $f(1) \neq 1$ and $f(x)f^{-1}(x) = x^\alpha$ for every $x \in (0, +\infty)$.

Mathematics Subject Classification (2010). Primary 39B22; Secondary 39B12, 26A18.

Keywords. Iterative functional equation, bijection, iterate, increasing solution.

In this note we answer Problem 3.8 posed in my article [4]. More precisely, given $\alpha \in \mathbb{R}$ we determine all increasing bijections $f: (0, +\infty) \rightarrow (0, +\infty)$ such that $f(1) \neq 1$ and

$$f(x)f^{-1}(x) = x^\alpha \quad \text{for every } x \in (0, +\infty). \quad (1.1)$$

Theorem 1. (i) If (1.1) admits a solution f which is an increasing bijection from $(0, +\infty)$ onto $(0, +\infty)$, then we have $\alpha > 2$. Let $\beta \in (1, +\infty)$ be the unique number satisfying $\beta + \frac{1}{\beta} = \alpha$.

If $f(1) = c \neq 1$ and $f_0 = f|_{[\min\{1, c\}, \max\{1, c\}]}$, then f_0 is continuous, $f_0(c) = c^\alpha$ and

$$\left(\frac{y}{x}\right)^{\frac{1}{\beta}} \leq \frac{f_0(y)}{f_0(x)} \leq \left(\frac{y}{x}\right)^\beta \quad (1.2)$$

for all $x, y \in [\min\{1, c\}, \max\{1, c\}]$ with $x < y$.

(ii) Conversely, let $c \in (0, 1) \cup (1, +\infty)$ and let $f_0: [\min\{1, c\}, \max\{1, c\}] \rightarrow \mathbb{R}$ be a continuous function such that $f_0(1) = c$, $f_0(c) = c^\alpha$ and (1.2) holds for all $x, y \in [\min\{1, c\}, \max\{1, c\}]$ with $x < y$. Then f_0 can be uniquely extended to an increasing bijection $f: (0, +\infty) \rightarrow (0, +\infty)$ satisfying (1.1).

Moreover:

- (a) if $c \in (0, 1)$, then $f(x) < \min\{x^\beta, x^{\frac{1}{\beta}}\}$ for every $x \in (0, +\infty)$;
- (b) if $c \in (1, +\infty)$, then $f(x) > \max\{x^\beta, x^{\frac{1}{\beta}}\}$ for every $x \in (0, +\infty)$.

Proof. (i) The case $f(1) = 1$ was treated in [4]. Then we have $\alpha > 0$ by Lemma 3.2 of [4].

Fix an increasing bijective solution $f: (0, +\infty) \rightarrow (0, +\infty)$ of (1.1) such that $f(1) \neq 1$.

Put $c = f(1)$ and $f_0 = f|_{[\min\{1, c\}, \max\{1, c\}]}$. Then f_0 is continuous, $f_0(1) = c$ and $f_0(c) = f^2(1) = [f(1)]^\alpha = c^\alpha$.

Fix $x, y \in [\min\{1, c\}, \max\{1, c\}]$ with $x < y$. Since all the iterates of f are increasing, by Lemma 2.3 of [4] we have

$$\frac{[f_0(x)]^{a_n}}{x^{a_{n-1}}} < \frac{[f_0(y)]^{a_n}}{y^{a_{n-1}}} \quad \text{and} \quad \frac{x^{a_{n+1}}}{[f_0(x)]^{a_n}} < \frac{y^{a_{n+1}}}{[f_0(y)]^{a_n}} \quad \text{for every } n \in \mathbb{N}.$$

Hence

$$\left(\frac{y}{x}\right)^{\frac{a_{n-1}}{a_n}} \leq \frac{f_0(y)}{f_0(x)} \leq \left(\frac{y}{x}\right)^{\frac{a_{n+1}}{a_n}} \quad \text{for every } n \in \mathbb{N}. \quad (1.3)$$

In particular, $a_{n-1} < a_{n+1}$ for every $n \in \mathbb{N}$. Finally, using Lemma 3.3 of [4] we get $\alpha > 2$ and passing with n to $+\infty$ in (1.3) we obtain (1.2).

(ii) Define a function $g_0: [\min\{0, \log c\}, \max\{0, \log c\}] \rightarrow \mathbb{R}$ by setting $g_0(x) = \log f_0(e^x)$. Then g_0 is continuous, $g_0(0) = \log c$, $g_0(\log c) = \alpha \log c$ and $\frac{1}{\beta} \leq \frac{g_0(y) - g_0(x)}{y - x} \leq \beta$ for all $x, y \in [\min\{0, \log c\}, \max\{0, \log c\}]$ with $x < y$.

Since $0 < \frac{1}{\beta} < 1 < \beta$, we conclude from Theorem 2 of [5] that g_0 can be uniquely extended to a continuous function $g: \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$g^2(x) = \alpha g(x) - x \quad \text{for every } x \in \mathbb{R}, \quad (1.4)$$

and this unique extension g satisfies

$$\begin{aligned} g(x) &< \min \left\{ \beta x, \frac{1}{\beta} x \right\} && \text{for every } x \in \mathbb{R}, \text{ if } c \in (0, 1), \\ g(x) &> \max \left\{ \beta x, \frac{1}{\beta} x \right\} && \text{for every } x \in \mathbb{R}, \text{ if } c \in (1, +\infty), \\ \frac{1}{\beta}(y - x) &\leq g(y) - g(x) \leq \beta(y - x) && \text{for all } x, y \in \mathbb{R} \text{ with } x < y. \end{aligned}$$

Consequently, g is strictly increasing and maps $\mathbb{R}$ onto $\mathbb{R}$. This jointly with (1.4) implies $g(x) + g^{-1}(x) = \alpha x$ for every $x \in \mathbb{R}$.

Now define a function $f: (0, +\infty) \rightarrow (0, +\infty)$ by setting $f(x) = e^{g(\log x)}$. Then f is an increasing bijection satisfying (1.1) such that (a) and (b) hold. The proof of the theorem is complete. $\square$

Remark 2. An answer to the problem posed during The Forty-ninth International Symposium on Functional Equations by Nicole Brillouët-Belluot (see [2]) as well as the result presented during The Fiftieth International Symposium on Functional Equations by Zoltán Boros (see [1]) can be derived from Theorem 10 of [5] in the same way as above (involving Lemmas 2.1–2.5 of [3] in the case of the problem of Nicole Brillouët-Belluot).

Acknowledgments

I thank Professor Karol Baron for calling my attention to the paper by Seiji Nabeya. This research was supported by Silesian University Mathematics Department (Iterative Functional Equations and Real Analysis program).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- [1] Boros, Z.: Talk given during the fiftieth international symposium on functional equations. *Aequat. Math.* **86**, 293 (2013)
- [2] Brillouët-Belluot, N.: Problem posed during The forty-ninth international symposium on functional equations. *Aequat. Math.* **84**, 312 (2012)
- [3] Morawiec, J.: On a problem of Nicole Brillouët-Belluot. *Aequat. Math.* **84**, 219–225 (2012)
- [4] Morawiec, J.: Around a problem of Nicole Brillouët-Belluot. *Aequat. Math.* doi:[10.1007/s00010-013-0216-8](https://doi.org/10.1007/s00010-013-0216-8)
- [5] Nabeya, S.: On the functional equation $f(p + qx + rf(x)) = a + bx + cf(x)$. *Aequat. Math.* **11**, 199–211 (1974)

Janusz Morawiec
Institute of Mathematics
University of Silesia
Bankowa 14
40-007 Katowice
Poland
e-mail: morawiec@math.us.edu.pl

Received: October 22, 2013

Revised: December 12, 2013