Correction

Correction to: Oscillations of difference equations with several deviated arguments

G. E. CHATZARAKIS, S. PINELAS AND I. P. STAVROULAKIS

Correction to: Aequat. Math. 88 (2014), 105–123

<https://doi.org/10.1007/s00010-013-0238-2>

Theorem 2.2. Assume that the sequences $(\tau_i(n))$, $1 \leq i \leq m$ are increasing, (1.1) holds,

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^m p_i(n) > 0 \quad (2.2)$$

and for every $i = 1, 2, \dots, m$

$$\liminf_{n \rightarrow \infty} \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j) > \frac{1}{e}. \quad (2.3)$$

Then all solutions of Eq. (E_D) oscillate.

Proof. Assume, for the sake of contradiction, that $(x(n))_{n \geq -w}$ is a nonoscillatory solution of (E_D) . Then it is either eventually positive or eventually negative. As $(-x(n))_{n \geq -w}$ is also a solution of (E_D) , we may restrict ourselves only to the case where $x(n) > 0$ for all large n . Let $n_1 \geq -w$ be an integer such that $x(n) > 0$ for all $n \geq n_1$. Then, there exists $n_2 \geq n_1$ such that

$$x(\tau_i(n)) > 0 \quad \forall n \geq n_2, \quad 1 \leq i \leq m.$$

As it was shown in Theorem 2.1, the sequence $(x(n))$ is eventually decreasing.

The original article can be found online at <https://doi.org/10.1007/s00010-013-0238-2>.

Set

$$b_i(n) = \left(\frac{n - \tau_i(n)}{n - \tau_i(n) + 1} \right)^{n - \tau_i(n) + 1}, \quad n \geq 1, \quad 1 \leq i \leq m. \quad (2.4)$$

Clearly

$$\frac{1}{4} \leq b_i(n) \leq \frac{1}{e}, \quad n \geq 1, \quad 1 \leq i \leq m. \quad (2.5)$$

By (2.3), there exist an integer $n_0 \geq n_2$ and a small positive number ε_0 such that

$$\sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j) > \frac{1}{e} + \varepsilon_0 \quad \text{for } n \geq n_0. \quad (2.6)$$

Let

$$d = e \left(\frac{1}{e} + \varepsilon_0 \right). \quad (2.7)$$

Combining (2.5), (2.6) and (2.7), we have that

$$\begin{aligned} \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} \frac{p_k(j)}{b_i(n)} &\geq \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} \frac{p_k(j)}{1/e} \\ &= e \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j) > e \left(\frac{1}{e} + \varepsilon_0 \right) = d > 1, \end{aligned}$$

or

$$\sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} \frac{p_k(j)}{b_i(n)} > d > 1. \quad (2.8)$$

Since $(x(n))$ is decreasing, clearly

$$\frac{x(\tau_i(n))}{x(n)} \geq 1 \quad \text{for all } n \geq n_0, \quad 1 \leq i \leq m. \quad (2.9)$$

By (E_D) , we have

$$\frac{x(n+1)}{x(n)} = 1 - \sum_{i=1}^m p_i(n) \frac{x(\tau_i(n))}{x(n)} \quad \text{for all } n \geq n_0. \quad (2.10)$$

On the other hand

$$\begin{aligned} \frac{x(n)}{x(\tau_i(n))} &= \prod_{j=\tau_i(n)}^{n-1} \frac{x(j+1)}{x(j)} \\ &= \prod_{j=\tau_i(n)}^{n-1} \left(1 - \sum_{k=1}^m p_k(j) \frac{x(\tau_k(j))}{x(j)} \right) \\ &\leq \prod_{j=\tau_i(n)}^{n-1} \left(1 - \sum_{k=1}^m p_k(j) \right) \end{aligned}$$

or

$$\frac{x(n)}{x(\tau_i(n))} \leq \prod_{j=\tau_i(n)}^{n-1} \left(1 - \sum_{k=1}^m p_k(j) \right). \quad (2.11)$$

By using (2.11) and the well-known inequality between the arithmetic and geometric means, we find that

$$\begin{aligned} \frac{x(n)}{x(\tau_i(n))} &\leq \prod_{j=\tau_i(n)}^{n-1} \left(1 - \sum_{k=1}^m p_k(j)\right) \\ &\leq \left[1 - \frac{1}{n - \tau_i(n)} \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j)\right]^{n-\tau_i(n)}, \end{aligned}$$

or

$$\frac{x(\tau_i(n))}{x(n)} \geq \left[1 - \frac{1}{n - \tau_i(n)} \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j)\right]^{-(n-\tau_i(n))}. \quad (2.12)$$

In view of

$$y(1-y)^\rho \leq \frac{\rho^\rho}{(1+\rho)^{1+\rho}}, \quad \forall y \in (0, 1), \quad \rho \in \mathbb{N}$$

inequality (2.12) gives

$$\frac{x(\tau_i(n))}{x(n)} \geq \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j) \left(\frac{n - \tau_i(n) + 1}{n - \tau_i(n)}\right)^{n-\tau_i(n)+1}. \quad (2.13)$$

Combining (2.13), (2.4) and (2.8), we obtain

$$\frac{x(\tau_i(n))}{x(n)} \geq \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} \frac{p_k(j)}{b_i(n)} \geq d \quad \text{for all } n \geq n_0. \quad (2.14)$$

Similarly,

$$\begin{aligned} \frac{x(n)}{x(\tau_i(n))} &= \prod_{j=\tau_i(n)}^{n-1} \frac{x(j+1)}{x(j)} \\ &= \prod_{j=\tau_i(n)}^{n-1} \left(1 - \sum_{k=1}^m p_k(j) \frac{x(\tau_k(j))}{x(j)}\right) \\ &\leq \prod_{j=\tau_i(n)}^{n-1} \left(1 - d \sum_{k=1}^m p_k(j)\right) \\ &\leq \left[1 - \frac{d}{n - \tau_i(n)} \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} p_k(j)\right]^{n-\tau_i(n)}. \end{aligned}$$

Therefore

$$\frac{x(\tau_i(n))}{x(n)} \geq d \sum_{k=1}^m \sum_{j=\tau_i(n)}^{n-1} \frac{p_k(j)}{b_i(n)} > d^2 \quad \text{for all } n \geq n_3 \geq n_0.$$

Applying this procedure ℓ -times, we obtain

$$\frac{x(\tau_i(n))}{x(n)} > d^\ell \quad \text{for all large } n. \quad (2.15)$$

On the other hand, since (2.2) holds, there exists a subsequence of integers $\theta(n)$ such that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^m p_i(\theta(n)) \geq c > 0.$$

By (2.10), we have

$$0 < \frac{x(n+1)}{x(n)} = 1 - \sum_{i=1}^m p_i(n) \frac{x(\tau_i(n))}{x(n)}$$

or

$$\frac{x(\tau(n))}{x(n)} \sum_{i=1}^m p_i(n) < 1,$$

where $(\tau(n))$ is defined by (1.20). Thus

$$\frac{x(\tau(\theta(n)))}{x(\theta(n))} \sum_{i=1}^m p_i(\theta(n)) < 1$$

or

$$\frac{x(\tau(\theta(n)))}{x(\theta(n))} < \frac{1}{\sum_{i=1}^m p_i(\theta(n))} \leq \frac{1}{c} < +\infty \quad \text{for large } n,$$

i.e., $\liminf_{n \rightarrow \infty} \frac{x(\tau(n))}{x(n)}$ exists. This contradicts (2.15).

The proof of the theorem is complete. $\square$

An immediate consequence of Theorem 2.2, is the following corollary:

Corollary 2.2. Assume that (2.2) holds, and for every $i = 1, 2, \dots, m$

$$\liminf_{n \rightarrow \infty} \sum_{k=1}^m \sum_{j=n-k_i}^{n-1} p_k(j) > \frac{1}{e}. \quad (2.3')$$

Then all solutions of Eq. (E'_D) oscillate.

Theorem 3.2. Assume that the sequences $(\sigma_i(n))$, $1 \leq i \leq m$ are increasing, (1.1') holds,

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^m p_i(n) > 0 \quad (3.2)$$

and for every $i = 1, 2, \dots, m$

$$\liminf_{n \rightarrow \infty} \sum_{k=1}^m \sum_{j=n+1}^{\sigma_i(n)} p_k(j) > \frac{1}{e}. \quad (3.3)$$

Then all solutions of Eq. (E_A) oscillate.

Example 4.2. Consider the delay difference equation

$$\Delta x(n) + \frac{1}{3e}x(n-1) + \frac{5}{6e}x(n-2) = 0, \quad n \geq 0. \quad (4.2)$$

Observe that

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^2 p_i(n) = \frac{1}{3e} + \frac{5}{6e} = \frac{7}{6e} > 0$$

and

$$\liminf_{n \rightarrow \infty} \sum_{k=1}^2 \sum_{j=n-1}^{n-1} p_k(j) = p_1(n-1) + p_2(n-1) = \frac{7}{6e} > \frac{1}{e},$$

$$\begin{aligned} \liminf_{n \rightarrow \infty} \sum_{k=1}^2 \sum_{j=n-2}^{n-1} p_k(j) &= p_1(n-2) + p_1(n-1) + p_2(n-2) + p_2(n-1) \\ &= 2 \cdot \frac{7}{6e} > \frac{1}{e}. \end{aligned}$$

Thus, all the conditions of Theorem 2.2 are satisfied and therefore all solutions of Eq. (4.2) oscillate.

Observe, however, that

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^2 \sum_{j=\tau(n)}^n p_i(j) = \limsup_{n \rightarrow \infty} \sum_{i=1}^2 \sum_{j=n-1}^n p_i(j) = \frac{7}{3e} < 1,$$

that is, condition (2.1) of Theorem 2.1 is not satisfied.

Example 4.4. Consider the advanced difference equation

$$\nabla x(n) - \frac{1}{2e}x(n+1) - \frac{3}{5e}x(n+2) = 0, \quad n \geq 1. \quad (4.4)$$

Observe that

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^2 p_i(n) = \frac{1}{2e} + \frac{3}{5e} = \frac{11}{10e} > 0$$

and

$$\liminf_{n \rightarrow \infty} \sum_{k=1}^2 \sum_{j=n+1}^{n+1} p_k(j) = p_1(n+1) + p_2(n+1) = \frac{11}{10e} > \frac{1}{e},$$

$$\begin{aligned} \liminf_{n \rightarrow \infty} \sum_{k=1}^2 \sum_{j=n+1}^{n+2} p_k(j) &= p_1(n+1) \\ &\quad + p_1(n+2) + p_2(n+1) + p_2(n+2) \\ &= 2 \cdot \frac{11}{10e} > \frac{1}{e}. \end{aligned}$$

Thus, all the conditions of Theorem 3.2 are satisfied and therefore all solutions of Eq. (4.4) oscillate.

Observe, however, that

$$\limsup_{n \rightarrow \infty} \sum_{i=1}^2 \sum_{j=n}^{\sigma(n)} p_i(j) = \limsup_{n \rightarrow \infty} \sum_{i=1}^2 \sum_{j=n}^{n+1} p_i(j) = \frac{11}{5e} < 1,$$

which means that condition (3.1) of Theorem 3.1 is not satisfied.

G. E. Chatzarakis

Department of Electrical and Electronic Engineering Educators
School of Pedagogical and Technological Education (ASPETE)

14121 N. Heraklio, Athens

Greece

e-mail: geaxatz@otenet.gr;

gea.xatz@aspete.gr

S. Pinelas

Departamento de Ciências Exactas e Naturais

Academia Militar

Av. Conde Castro Guimarães

2720-113 Amadora

Portugal

e-mail: sandra.pinelas@gmail.com

I. P. Stavroulakis

Department of Mathematics

University of Ioannina

451 10 Ioannina

Greece

e-mail: ipstav@cc.uoi.gr