

# Symmetric Minkowski planes ordered by separation

Helmut Karzel, Jarosław Kosiorek and Andrzej Matraś

Dedicated to Mario Marchi

**Abstract.** In Karzel et al. (J. Geom. 99:116–127, 2009) we introduced for a symmetric Minkowski plane  $\mathfrak{M} := (P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  an order concept by the notion of an orthogonal valuation for the circles of  $\Lambda$  and showed that there is a one to one correspondence between the valuations and the halforderings of the accompanying commutative field. Here we consider an order concept which is based on the notion of *separation* for quadruples of concyclic points and establish the connections between these two notions. Our main result (cf. Theorem 3.3) states that these concepts are equivalent.

**Mathematics Subject Classification (2010).** 51B20(2000).

**Keywords.** Minkowski planes, order, valuation, separation.

## Introduction

Order concepts in geometry are usually based on both, *points* and *blocks*. Minkowski planes belong to the chain structures and so there appear two types of blocks, *generators* and *chains* latter also called *circles* (in the case of Minkowski planes). In [1] we showed that in a symmetric Minkowski plane  $\mathfrak{M} := (P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  an order structure can be introduced where only the set  $\Lambda$  of circles is involved, namely by a *valuation* which associates to each circle  $C \in \Lambda$  a value  $[C]$  which is 1 or  $-1$ . Then, since  $\Lambda$  can be endowed with a multiplication “ $\cdot$ ” such that  $(\Lambda, \cdot)$  becomes a group, it turns out that the valuation is a homomorphism of  $(\Lambda, \cdot)$  in the cyclic group  $(\{1, -1\}, \cdot)$  of order two hence  $[A \cdot B] = [A] \cdot [B]$  for  $A, B \in \Lambda$ . Furthermore to a symmetric Minkowski plane  $\mathfrak{M}$  there corresponds a commutative field  $(\mathbf{F}, +, \cdot)$  such that  $(\Lambda, \cdot)$  is isomorphic to the projective linear group  $PGL(2, \mathbf{F})$  and if  $[ ]$  is an orthogonal valuation of

$\mathfrak{M}$  then there is a halfordering  $\eta$  of the commutative field  $(\mathbf{F}, +, \cdot)$  such that if  $A_f \in GL(2, \mathbf{F})$  is a matrix representing a chain  $A \in \Lambda$  then  $[A] = \eta(\det(A_f))$ .

In [2] Kroll (cf. Subsection 1.6) defined order structures for Benz planes—and so also for Minkowski planes—starting inter alia from the notion of *separation*, i.e. a function which maps each quadruple  $(a, b, c, d)$  consisting of concyclic points with  $a, b \neq c, d$  on a value  $[a, b|c, d]$  in  $\{1, -1\}$  such that certain conditions are satisfied (cf. Sect. 1.5).

In Sect. 1.5 we recall for symmetric Minkowski planes  $\mathfrak{M}$  the notions *orthogonal valuation*, *order valuation* and then *halfordered* and *ordered* symmetric Minkowski planes  $(\mathfrak{M}, [\ ])$  (based on valuations). Theorem 1.13. describes properties of these structures and Theorem 1.14. recalls relations to the halforderings and orderings of the corresponding commutative field  $(\mathbf{F}, +, \cdot)$ .

Following Kroll we introduce the concepts *separation* and *order separation* by conditions **(T1)**, **(T2)** and **(T1)**, **(T2)**, **(T3)**, respectively.

Theorem 2.1 shows how one can derive from an orthogonal valuation  $[\ ]$  of a symmetric Minkowski plane  $\mathfrak{M}$  a separation  $\tau$  of  $\mathfrak{M}$ . The separation  $\tau$  is harmonic iff the valuation  $[\ ]$  is harmonic, and  $\tau$  is an order separation iff  $[\ ]$  is an order valuation.

Finally in Sect. 3 we start from a pair  $(\mathfrak{M}, \tau)$  where  $\mathfrak{M}$  is a symmetric Minkowski plane and  $\tau$  a separation of  $\mathfrak{M}$  and prove in Theorem 3.1 some properties of  $(\mathfrak{M}, \tau)$ . Then we show in Theorem 3.2 how one can derive from  $\tau$  a halfordering  $\eta_\tau$  for the corresponding field  $(\mathbf{F}, +, \cdot)$ . Combining these results starting from a halfordered symmetric Minkowski  $(\mathfrak{M}, [\ ])$  we can derive firstly (as in [1]) a halfordering  $\eta$  of the corresponding field  $(\mathbf{F}, +, \cdot)$  and secondly a separation  $\tau$  of  $\mathfrak{M}$  and then a halfordering  $\eta_\tau$  of  $(\mathbf{F}, +, \cdot)$ . We show  $\eta = \eta_\tau$  and this proves our main result Theorem 3.3.

## 1. Preliminaries and supplements

### 1.1. Hyperbola structures and Minkowski planes

In this paper we will use mostly the same notations as in [1]. Let  $(P, \mathfrak{G}_1, \mathfrak{G}_2)$  be a net. As in [1] let  $P^{(3)} := \{\{a, b, c\} \in \binom{P}{3} \mid \{a, b, c\} \text{ is joinable}\}$  and let  $\mathfrak{C}$  be the set of all chains of  $(P, \mathfrak{G}_1, \mathfrak{G}_2)$ . The quadruple  $(P, \mathfrak{C}, \mathfrak{G}_1, \mathfrak{G}_2)$  is then called *maximal chain structure*.

For  $a, b, c, d \in P$  we add the notions  $(a, b, c, d)^\square := \{ab, ba, cd, dc\}$  and  $\{a, b, c, d\}^\square := \{\{a, b, c, d\}, (a, b, c, d)^\square, (a, c, d, b)^\square, (a, d, b, c)^\square\}$ .

If  $\{x, y, z, u\} \in \{a, b, c, d\}^\square$  then  $\{x, y, z, u\}^\square = \{a, b, c, d\}^\square$ .

A quadruple  $(P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  is called *hyperbola structure* if  $\Lambda \subseteq \mathfrak{C}$  such that:

- (I)  $\forall \{a, b, c\} \in P^{(3)} \exists_1 L \in \Lambda : \{a, b, c\} \subseteq L -$  we set  $(a, b, c)^\circ := L -$ .

If  $(P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  is a hyperbola structure then a subset  $S$  of  $P$  is called *concylic* if there is a *circle*  $L \in \Lambda$  such that  $S \subseteq L$  – if  $|S| \geq 3$  then  $L$  is uniquely determined and we set then  $S^\circ := L$ . A hyperbola structure  $(P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  is called *symmetric Minkowski plane* if the so called “*Symmetry Axiom*” is satisfied which we can express in the following form (cf. [1, (2.7)]) :

**(S)** If a quadruple  $(a, b, c, d)$  consists of concyclic points then also the set  $(a, b, c, d)^\square$  is concyclic.

In a symmetric Minkowski plane, if  $F := \{a, b, c, d\}$  is a set consisting of four distinct concyclic points then by Axiom **(S)**, the set  $F^\square$  consists of four subsets of points and each subset consisting of four distinct concyclic points so that (by Axiom **(I)**) we can form the set

$$\Lambda_F := \{\{a, b, c, d\}^\circ, (a, b, c, d)^\square, (a, c, d, b)^\square, (a, d, b, c)^\square\}$$

consisting of four distinct circles which are orthogonal in pairs. A quadruple  $(a, b, c, d)$  consisting of four distinct concyclic points is called *harmonic* if  $\{c, d\} = (a, b, c)^\circ \cap (ab, ba, c)^\circ$ . Let  $P^{(4h)}$  denote the *set of all harmonic quadruples*.

From now on let  $\mathfrak{M} = (P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  be a symmetric Minkowski plane, let a circle  $E \in \Lambda$  be fixed and for  $A, B \in \mathfrak{C}$  let  $A \cdot B := t(A, E, B) = \widehat{AB}(E)$  then (cf. [1, p. 118 f]):

**Theorem 1.1.**  $(\mathfrak{C}, \cdot)$  is a group with the neutral element  $E$ ,  $t(A, B, C) = A \cdot B^{-1} \cdot C$  and the orthogonality of chains is described by:

$$A \perp B \Leftrightarrow A \cdot B^{-1} = B \cdot A^{-1} \quad \text{and} \quad A \neq B.$$

Moreover  $\Lambda$  is a subgroup of  $(\mathfrak{C}, \cdot)$ .

We recall that a quadruple  $(A, B, C, D)$  of circles is called a *quadrangle* if  $D = t(A, B, C)$ , a quadrangle is called a *square* if  $A \perp B \perp C \perp D \perp A$  and a *total square* if moreover  $A \perp C$  and  $B \perp D$ . If  $F$  is a set of four distinct concyclic points then  $\Lambda_F$  is a total square,

## 1.2. Properties of symmetric Minkowski planes

In a symmetric Minkowski planes  $\mathfrak{M} = (P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$  we define  $\Gamma^- := \{\widehat{AB} \mid A, B \in \Lambda\}$ , denote by  $\Gamma := \langle \Gamma^- \rangle$  the group generated by  $\Gamma^-$  and set

$$(\Lambda \times \Lambda)_\perp := \{(A, B) \in (\Lambda \times \Lambda) \mid A \perp B\} \text{ and}$$

$$(\Lambda \times \Lambda)_{\perp,s} := \{(A, B) \in (\Lambda \times \Lambda)_\perp \mid A \cap B \neq \emptyset\}.$$

For a set  $S$  of concyclic points with  $|S| \geq 3$  there is exactly one circle in  $\Lambda$  containing  $S$  which shall be denoted also by  $S^\circ$ .

Finally the notions of 1- and 2-perspectivities can be defined:

Let  $A, B \in \Lambda$  then the map

$$[A \xrightarrow{1} B] : A \rightarrow B; x \mapsto xB \text{ and } [A \xrightarrow{2} B] : A \rightarrow B; x \mapsto Bx$$

is called 1- and 2-perspectivity, respectively.

From [1, Section 2.1, (2.5),(2.6),(2.7)] follows:

**Theorem 1.2.** *Let  $A, B, C, X \in \Lambda$  and let  $(a, b, c, d) \in X$  with  $a, b \neq c, d$ , then:*

- (1)  $\Gamma \leqslant \text{Aut}(P, \Lambda, \mathfrak{G}_1 \cup \mathfrak{G}_2)$ .
- (2) *If  $a \neq b$  then  $\{ab, ba, cd\} \in P^{(3)}$ ,  $dc \in K := (ab, ba, cd)^\circ$  and  $K \perp X$ .*
- (3) *If  $|\{a, b, c, d\}| \geq 3$  then  $X = (a, b, c, d)^\circ$  and  $K := ((a, b, c, d)^\square)^\circ$  are uniquely determined circles of  $\Lambda$  and  $X \perp K$ ; if  $a = b$  or  $c = d$  then  $K = ((c, d, a)^\beta)^\circ$  and  $a \in X \cap K$  or  $K = ((a, b, c)^\beta)^\circ$  and  $c \in X \cap K$ , respectively.*
- (4) *If  $|\{a, b, c, d\}| = 4$  and  $F := \{a, b, c, d\} \subseteq E$  then  $\Lambda_F$  is a subgroup of  $(\Lambda, \cdot)$  and consists of four circles which are orthogonal in pairs (cf. Theorem 1.1), i.e.  $\Lambda_F$  is a total square.*
- (5) *If  $A, B \perp C$ ,  $A \cap C = B \cap C = \{p, q\}$  and  $p \neq q$  then  $A = B$ .*

*Proof.* (5) Assume  $A \neq B$  hence  $A \cap B = \{p, q\}$  then by [1, (3.5.1)],  $pq, qp \in C$ . But since  $p, q \in C$  and  $p \neq q$  we have  $pq, qp \notin C$ . Consequently  $A = B$ .  $\square$

**Theorem 1.3.** *Let  $K \in \Lambda$ ,  $a, b, c, d, e \in K$  with  $a, b \neq c, d, e$  and  $a \neq b$  and let  $L_c := \{ab, ba, de\}^\circ$ ,  $L_d := \{ab, ba, ec\}^\circ$ ,  $L_e := \{ab, ba, cd\}^\circ$  and  $M := \widetilde{L_e L_c}(L_d) = L_e \cdot L_d^{-1} \cdot L_c$  then  $M = \{ab, ba, d\}^\circ$ ,  $M \perp K$  and  $d \in M \cap K$ , i.e.  $(M, K) \in (\Lambda \times \Lambda)_{\perp s}$ .*

*Proof.* Since  $\{ab, ba, cd, dc\} \subseteq L_e$ ,  $\{ab, ba, ce, ec\} \subseteq L_d$  and  $\{ab, ba, de, ed\} \subseteq L_c$  we have  $ab, ba \in \text{Fix} \widetilde{L_e L_c}$  and  $\widetilde{L_e L_c}(ce) = (de)(cd) = d$  hence  $M = \widetilde{L_e L_c}(L_d) = \{ab, ba, d\}^\circ$  with  $M \perp K$  and  $d \in M \cap K$  and so  $(M, K) \in (\Lambda \times \Lambda)_{\perp s}$ .  $\square$

By [1, (2.5.2) and (3.9)] we have:

**Theorem 1.4.** *The set  $\Gamma^- := \{\widetilde{AB} \mid A, B \in \Lambda\}$  consists of automorphisms of  $(P, \Lambda)$  and acts transitively on the set of circles  $\Lambda$  preserving orthogonality and if  $\gamma$  is an element of the group  $\Gamma$  generated by  $\Gamma^-$  then there exist exactly two elements  $A, B \in \Lambda$  such that either  $\gamma = \widetilde{AB}$  or  $\gamma = \widetilde{AB} \circ \widetilde{E}$ . Moreover  $\widetilde{AB} \circ \widetilde{CD} = (A \cdot D^{-1})(C^{-1} \cdot B) \circ \widetilde{E}$  and  $\Gamma^+ := \widetilde{E} \circ \Gamma^-$  acts regularly on the set  $P^{(3)}$ .*

Therefore, if  $A, B \in \Lambda$  and  $C := \widetilde{AE}(B) = A \cdot B^{-1} \cdot E = A \cdot B^{-1}$  we have:

$$(A, B) \in (\Lambda \times \Lambda)_{\perp s} \Leftrightarrow (E, C) \in (\Lambda \times \Lambda)_{\perp s}.$$

Consequently,  $\forall (A, B), (C, D) \in (\Lambda \times \Lambda)_{\perp s} : |A \cap B| = |C \cap D|$ .

### 1.3. The corresponding field and the characteristic of a symmetric Minkowski plane

As in [1] we denote by  $(\mathbf{F}, +, \cdot)$  the commutative field corresponding to the symmetric Minkowski plane  $\mathfrak{M}$  (cf. [1, sec.1]) and we define  $\text{char } \mathfrak{M} := \text{char } \mathbf{F}$ .

If  $\begin{pmatrix} c_1, c_2 \\ c_3, c_4 \end{pmatrix} \in GL(2, \mathbf{F})$  represents the circle  $C$  then  $C \perp E$  implies  $c_1 = -c_4$  and if  $c_1 = -c_4$  then:

$C \cap E \neq \emptyset \Leftrightarrow$  the characteristic equation  $\Xi_C(t) = t^2 - (c_1^2 + c_2 \cdot c_3) = 0$  is solvable  $\Leftrightarrow (c_1^2 + c_2 \cdot c_3) \in \mathbf{F}^{(2)} := \{\lambda^2 \mid \lambda \in \mathbf{F} \setminus \{0\}\}$ .

Hence if  $(c_1^2 + c_2 \cdot c_3) \in \mathbf{F}^{(2)}$  then:  $|C \cap E| = 2$  if  $\text{char}\mathbf{F} \neq 2$  and  $|C \cap E| = 1$  if  $\text{char}\mathbf{F} = 2$ . This supplements the statements of [1, (3.7)]:

**Theorem 1.5.** *Let  $(A, B) \in (\Lambda \times \Lambda)_{\perp,s}$  then:*

- (1)  $|A \cap B| = 2 \Leftrightarrow \text{char}\mathfrak{M} \neq 2$ .
- (2)  $|A \cap B| = 1 \Leftrightarrow \text{char}\mathfrak{M} = 2$ .
- (3) *If  $\text{char}\mathfrak{M} \neq 2$  then  $\forall (a, b, c) \in P^{(3)} \exists d \in P$  such that  $(a, b, c, d)$  is a harmonic quadruple and  $d$  is given by  $\{a, b, c\}^\circ \cap \{ab, ba, c\}^\circ = \{c, d\}$ . If  $\text{char}\mathfrak{M} = 2$ , there are no harmonic quadruples.*

Now let  $C, D \in \Lambda$  be circles with  $C, D \perp E$  and  $C \cap E \neq \emptyset, D \cap E \neq \emptyset$  and let  $\begin{pmatrix} c_1, c_2 \\ c_3, -c_1 \end{pmatrix}, \begin{pmatrix} d_1, d_2 \\ d_3, -d_1 \end{pmatrix} \in GL(2, \mathbf{F})$  be a representing matrices. Then we may assume  $d_1^2 + d_2 \cdot d_3 = c_1^2 + c_2 \cdot c_3 = 1$  and moreover  $c_1 = 1$  if  $c_2 \cdot c_3 = 0$  and  $d_1 = 1$  if  $d_2 \cdot d_3 = 0$ . Let  $M_C := \begin{pmatrix} 0, c_2 \\ 1, -c_1 \end{pmatrix}$  if  $c_2 \neq 0$ ,  $M_C := \begin{pmatrix} c_1, 1 \\ c_3, 0 \end{pmatrix}$  if  $c_2 = 0 \neq c_3$  and  $M_C := \begin{pmatrix} 1, 1 \\ 1, -1 \end{pmatrix}$  if  $c_2 = c_3 = 0$  and  $\text{char}\mathbf{F} \neq 2$  and let  $M_D$  be defined accordingly. Then if  $L \in \Lambda$  is the circle represented by the matrix  $M_D \cdot M_C^{-1}$ . Then the map  $\widetilde{LL^{-1}}$  fixes  $E$  and maps  $C$  onto  $D$ . This gives us:

**Theorem 1.6.**  *$\Gamma$  acts transitively on the set  $(\Lambda \times \Lambda)_{\perp,s}$ .*

In the same way using representing matrices one confirms the *Three Reflection Theorems*:

**Theorem 1.7.** *Let  $A, B, C \in \Lambda$  with  $A \cap B \cap C \neq \emptyset$  and  $p \in A \cap B \cap C$  then:*

- (1) *If  $|A \cap B \cap C| \geq 2$  then:  $\exists D \in \Lambda : \widetilde{A} \circ \widetilde{B} \circ \widetilde{C} = \widetilde{D}$ .*
- (2) *If there is an  $L \in \Lambda$  with  $p \in L$  and  $A, B, C \perp L$  then:  $\exists D \in \Lambda : \widetilde{A} \circ \widetilde{B} \circ \widetilde{C} = \widetilde{D}$  (and then  $p \in D$  and  $D \perp L$  or  $D = L$ ).*

By the definition of the maps  $\widetilde{AB}$  one obtains the following representations of 1- and 2-perspectivities:

**Proposition 1.8.** *Let  $A, B, C \in \Lambda$  and  $D := \widetilde{AB}(C)$  then:*

- (1)  $[A \xrightarrow{1} B] = \widetilde{BA}|_A$  and  $[A \xrightarrow{2} B] = \widetilde{AB}|_A$ .
- (2)  $\widetilde{AB}|_C = [A \xrightarrow{2} D] \circ [C \xrightarrow{1} A] = [B \xrightarrow{1} D] \circ [C \xrightarrow{2} B]$ .
- (3)  $\bar{A} : E \rightarrow E; x \mapsto E(xA) = [A \xrightarrow{2} E] \circ [E \xrightarrow{1} A](x) = \widetilde{AE} \circ \widetilde{AE}(x)$ .
- (4) *If  $A \perp B$  then  $\psi := [B \xrightarrow{2} A] \circ [A \xrightarrow{1} B] = \widetilde{BA} \circ \widetilde{BA}|_A$  and if for  $x \in A, x' := \widetilde{B}(x)$  then:*

$$x' = \psi(x) \in A, (x')' = x, xx' \in B \quad \text{and} \quad \widetilde{A}(xx') = x'x \in B.$$

**Theorem 1.9.** *Let  $(a, b, c, d)$  and  $(a', b', c', d')$  be harmonic quadruples and let  $\gamma \in \Gamma$  then:*

- (1)  $(b, a, c, d), (c, d, a, b), (ab, ba, cd, dc), (ab, ba, c, d)$  and  $(\gamma(a), \gamma(b), \gamma(c), \gamma(d))$  are harmonic quadruples.
- (2)  $\exists \sigma \in \Gamma : \sigma(a) = a', \sigma(b) = b', \sigma(c) = c', \sigma(d) = d'$ .
- (3) The circles  $K := \{a, b, c\}^\circ$ ,  $L := \{ab, ba, cd\}^\circ$ ,  $M := \{ab, ba, c\}^\circ$  and  $N := \{a, b, cd\}^\circ$  are orthogonal in pairs hence  $\mathfrak{H} := (K, L, M, N)$  is a total square in  $\Lambda$  and  $(K, M), (K, N), (L, M), (L, N) \in (\Lambda \times \Lambda)_{\perp, s}$  and  $K \cap M = \{c, d\}$ ,  $K \cap N = \{a, b\}$ ,  $L \cap M = \{ab, ba\}$  and  $L \cap N = \{cd, dc\}$ .  $K^{-1} \cdot \{K, L, M, N\}$  is a subgroup of  $(\Lambda, \cdot)$  isomorphic to the Klein four group.

*Proof.* (1) The definition *harmonic* is symmetric in the first two and in the last two arguments and from  $\{c, d\} = \{a, b, c\}^\circ \cap \{ab, ba, c\}^\circ$  we obtain  $\{a, b, c\}^\circ = (\{c, d, a\}^\circ \cap ab, ba, c, d)$  are four distinct concyclic points hence by (S),  $(ab, ba, c, d)^\square = \{a, b, cd, dc\}$  is concyclic, i.e.  $\{cd, dc, a\}^\circ \ni a, b$  and so  $\{a, b\} = \{c, d, a\}^\circ \cap \{cd, dc, a\}^\circ$ , i.e.  $(c, d, a, b) \in P^{(4h)}$ .

(2) By Theorem 1.6 there is exactly one  $\sigma \in \Gamma^+$  with  $\sigma(a) = a'$ ,  $\sigma(b) = b'$ ,  $\sigma(c) = c'$  and so  $\{\sigma(c), \sigma(d)\} = \{\sigma(a), \sigma(b), \sigma(c)\}^\circ \cap \{\sigma(a)\sigma(b), \sigma(b)\sigma(a), \sigma(c)\}^\circ = \{a', b', c'\}^\circ \cap \{a'b', b'a', c'\}^\circ = \{c', d'\}$  implying  $\sigma(d) = d'$ .  $\square$

**Theorem 1.10.** Let  $a, b, c, d$  be four distinct concyclic points,  $K := \{a, b, c\}^\circ$ ,  $C := \{bc, cb, a\}^\circ$ ,  $D := \{bd, db, a\}^\circ$ ,  $H := \{cd, dc, a\}^\circ$  and let  $B \in \Lambda$  [according to Theorem 1.7(2)] be such that  $\tilde{B} = \tilde{D} \circ \tilde{H} \circ \tilde{C}$ . Then  $\tilde{B}(\tilde{C}(d)) = \tilde{D}(c)$  and  $(a, b, \tilde{C}(d), \tilde{D}(c))$  is harmonic.

*Proof.* By the definitions we have:  $a \in C, H, D, K$ ;  $C, H, D \perp K$ ,  $\tilde{C}(b) = c$ ,  $\tilde{H}(c) = d$  and  $\tilde{D}(d) = b$  hence, by Theorem 1.9(2),  $B \perp K$ ,  $\tilde{B}(a) = \tilde{D} \circ \tilde{H} \circ \tilde{C}(a) = a$ ,  $\tilde{B}(b) = \tilde{D} \circ \tilde{H} \circ \tilde{C}(b) = b$  and  $\tilde{B}(\tilde{C}(d)) = \tilde{D} \circ \tilde{H} \circ \tilde{C}(\tilde{C}(d)) = \tilde{D} \circ \tilde{H}(d) = \tilde{D}(c)$ . Since moreover  $\tilde{C}(d), \tilde{D}(c) \in K$  we have  $\{\tilde{C}(d), \tilde{D}(c)\} = \{a, b, \tilde{C}(d)\}^\circ \cap \{ab, ba, \tilde{C}(d)\}^\circ$  i.e.  $(a, b, \tilde{C}(d), \tilde{D}(c))$  is harmonic.  $\square$

#### 1.4. Orthogonal valuations and separations in symmetric Minkowski planes

In [1, p. 120 and p. 126] a map  $[\ ] : \Lambda \rightarrow \{1, -1\}; X \mapsto [X]$  was called *orthogonal valuation* if

- (O 1) For each square  $(A, B, C, D) : [A] \cdot [B] \cdot [C] \cdot [D] = 1$ . and *homomorphic valuation* if this equation holds true even for each quadrangle. An orthogonal valuation is called *order valuation* if
- (O 2) For each total square  $(A, B, C, D) :$  exactly two of the values  $[A], [B], [C], [D]$  equal 1.

By Theorems 1.4, 1.5 and [1, (4.5) and (4.8)] we have:

**Theorem 1.11.** Let  $[\ ] : \Lambda \rightarrow \{1, -1\}; X \mapsto [X]$  be an orthogonal valuation of  $\mathfrak{M}$ , let  $E \in \Lambda$  with  $[E] = 1$  be fixed, for  $A, B \in \Lambda$  let  $A \cdot B := t(A, E, B)$  and for  $\gamma = \widetilde{AB} \in \Gamma^-$  or  $\gamma = \widetilde{AB} \circ \widetilde{E} \in \Gamma^+$  let  $[\gamma]' := [A] \cdot [B]$ . Then:

- (1)  $[ ]$  is a homomorphic valuation hence  $\forall A, B, C \in \Lambda : [t(A, B, C)] = [A \cdot B^{-1} \cdot C] = [A] \cdot [B] \cdot [C]$  and  $[A \cdot B] = [A] \cdot [B]$ .
- (2)  $['] : (\Gamma, \circ) \rightarrow (\{1, -1\}, \cdot); \gamma \mapsto [\gamma]'$  is a homomorphism.
- (3)  $\forall X \in \Lambda, \forall \gamma \in \Gamma : [\gamma(X)] = [\gamma]' \cdot [X]$ .
- (4) There is an  $\iota \in \{1, -1\}$  such that:  $\forall (A, B) \in (\Lambda \times \Lambda)_{\perp s} : [A] \cdot [B] = \iota$ .
- (5) If  $[ ]$  is an order valuation then  $\iota = -1$ .

An orthogonal valuation  $[ ]$  is called *harmonic* if  $\iota = -1$  and *anharmonic* if  $\iota = 1$ .

Let  $P^{(4)} := \{(a, b, c, d) \in P^4 \mid \{a, b, c, d\} \text{ concyclic} \wedge a, b \neq c, d\}$ .

A map  $\tau : P^{(4)} \rightarrow (\{1, -1\}, \cdot) ; (a, b, c, d) \mapsto [a, b|c, d]$  is called a *separation* if the following conditions are satisfied:

- (T1) For all concyclic  $a, b, c, d, e \in P$  with  $a, b \neq c, d, e$  holds:  
 $[a, b|c, d] \cdot [a, b|d, e] = [a, b|c, e]$ .
- (T2) For all  $(a, b, c, d) \in P^{(4)}$ , for all  $L \in \Lambda$ :  
 $[a, b|c, d] = [La, Lb|Lc, Ld] = [aL, bL|cL, dL]$ , i.e. the separation function is invariant by 1- and 2-perspectivities.

If moreover the axiom

- (T3)  $\forall (a, b, c, d) \in P^{(4)}$  with  $a \neq b, c \neq d$ : exactly one of the values  $[a, b|c, d]$ ,  $[a, c|d, b]$ ,  $[a, d|b, c]$  is  $-1$

is valid then  $\tau$  is called an *order separation*.

A separation is called *harmonic* if for a harmonic quadruple  $(a, b, c, d)$  (and then for all) the value  $[a, b|c, d]$  is  $-1$ . This definition corresponds E. Sperner's definition of a harmonic separation.

### 1.5. Kroll's order concepts for Minkowski planes

Kroll [2] developed a thoroughly theory of order questions for Benz planes. Benz planes contain as a subclass the Minkowski planes. We present some of Kroll's concepts specialized for Minkowski planes  $\mathfrak{M} := (P, \Lambda, \mathfrak{G}_1, \mathfrak{G}_2)$ : Let  $(\Lambda \times P \times P)^* := \{(A, b, c) \in \Lambda \times P \times P \mid b, c \notin A\}$ . Then a map  $\alpha : (\Lambda \times P \times P)^* \rightarrow \{1, -1\}; (A, b, c) \mapsto (A|b, c)$  is called *order function* of  $\mathfrak{M}$  if :

- O 1**  $\forall A \in \Lambda$  and  $\forall x, y, z \in P \setminus A : (A|x, y) \cdot (A|y, z) \cdot (A|z, x) = 1$ .
- O 2** (circle relation)  $\forall A, B, C \in \Lambda$  such that  $A \cap C = B \cap C$  and  $|A \cap C| \geq 1$  and  $\forall x, y \in C \setminus A : (A|x, y) = (B|x, y)$ .
- OV**  $\forall G \in \mathfrak{G}_1 \cup \mathfrak{G}_2, \forall A, B \in \Lambda$  with  $A \cap G = B \cap G$  and  $\forall c, d \in G \setminus A : (A|c, d) = (B|c, d)$  (cf. [2], p. 224 and 231]).

If  $(\mathfrak{M}, \alpha)$  is a Minkowski plane with an order function, Kroll derives a separation  $\tau_\alpha$  by:

- (V) For  $K \in \Lambda$ , for  $a, b, c, d \in K$  with  $a, b \neq c, d$  and  $A, L \in \Lambda$  with  $A \cap K = \{a\}, L \cap K = \{a, b\}$  let  $[a, b|c, d] := (A|c, d) \cdot (L|c, d)$ . ([2], p. 234])

He shows that  $\tau_\alpha$  satisfies the conditions **(T1)** and **(T2)** if  $\mathfrak{M}$  is a symmetric Minkowski plane (cf. [2, p. 234 and p. 237], T4\*i). Conversely every separation of a symmetric Minkowski plane is a separation in the sense of Kroll.

## 2. From an orthogonal valuation to a separation

**Theorem 2.1.** Let  $(\mathfrak{M}, [\cdot])$  be a halfordered symmetric Minkowski plane, let  $\iota$  be the value corresponding to  $(\mathfrak{M}, [\cdot])$  according to Theorem 1.11(4) and let  $\tau : P^{(4)} \rightarrow \{1, -1\}; (a, b, c, d) \mapsto [a, b|c, d]$  be the map defined by  $[a, b|c, d] := 1$  if  $a = b$  and  $[a, b|c, d] := \iota \cdot [(a, b, c, d)^\square] \cdot [(a, b, c, d)^\circ]$  if  $a \neq b$  then:

- (1)  $\tau$  is a separation and if  $(a, b, c, d)$  is harmonic then  $[a, b|c, d] = \iota$ . Hence the derived separation  $\tau$  is harmonic if and only if the valuation  $[\cdot]$  is harmonic.
- (2) If  $[\cdot]$  is an order valuation then  $\tau$  is an order separation.

*Proof.* **(T1)** Let  $(a, b, c, d), (a, b, d, e) \in P^{(4)}$ . If  $a = b$  then  $[a, b|c, d] = [a, b|d, e] = [a, b|c, e] = 1$  and the equation **(T1)** is valid. Let  $a \neq b$  then  $(a, b, c), (a, b, d) \in P^{(3)}$ , so we can form the circles  $K := \{a, b, c\}^\circ$  and  $K' := \{a, b, d\}^\circ$  and by the assumption,  $(a, b, c, d), (a, b, d, e) \in P^{(4)}$ , we have  $a, b, c, d, e \in K = K'$ . If we define the circles  $L_c, L_d, L_e$  and  $M$  according to Theorem 1.3 then  $(M, K) \in (\Lambda \times \Lambda)_{\perp s}$  hence by Theorem 1.11(4),  $[M] \cdot [K] = \iota$  and by definition of  $\tau$  we have  $[a, b|c, d] := \iota \cdot [K] \cdot [L_e], [a, b|d, e] := \iota \cdot [K] \cdot [L_c]$  and  $[a, b|e, c] := \iota \cdot [K] \cdot [L_d]$ . By Theorem 1.11(1)), this implies:

$$[a, b|c, d] \cdot [a, b|d, e] \cdot [a, b|e, c] = \iota \cdot [K] \cdot [L_e] \cdot \iota \cdot [K] \cdot [L_c] \cdot \iota \cdot [K] \cdot [L_d] = \iota \cdot [K] \cdot [L_e \cdot L_d^{-1} \cdot L_c] = \iota \cdot [K] \cdot [M] = \iota \cdot \iota = 1.$$

**(T2)** Let  $K, L \in \Lambda, \{a, b, c, d\} \in \binom{\Lambda}{4}$  and  $H := \{ab, ba, cd\}^\circ$  then by Theorem 1.2(2),  $H \perp K$  hence  $[a, b|c, d] = [K] \cdot [H] \cdot \iota$  and by Proposition 1.8,

$$\pi_1 := [K \xrightarrow{1} L] = \widetilde{LK}|_K, \quad \pi_2 := [K \xrightarrow{2} L] = \widetilde{KL}|_K.$$

Then  $\widetilde{KL}(K) = L = \widetilde{LK}(K)$  and if  $H_1 := \widetilde{LK}(H), H_2 := \widetilde{KL}(H)$  then  $H_1, H_2 \perp L$  and so  $H_i = ((\pi_i(a), \pi_i(b), \pi_i(c), \pi_i(d))^\square)^\circ$ . Thus

$[\pi_i(a), \pi_i(b)|\pi_i(c), \pi_i(d)] = [L] \cdot [H_i] \cdot \iota$  with  $H_1 = L \cdot H^{-1} \cdot K, H_2 = K \cdot H^{-1} \cdot L$  implying by Theorem 1.11(1),  $[H_i] = [K] \cdot [H] \cdot [L]$ . Therefore  $\iota \cdot [\pi_i(a), \pi_i(b)|\pi_i(c), \pi_i(d)] = [K] \cdot [H] \cdot [L] = [K] \cdot [H] = \iota \cdot [a, b|c, d]$ .

Now let  $(a, b, c, d)$  be harmonic and let  $K, L, M, N$  be defined according to Theorem 1.9(3). Then by Theorem 1.9(3) and Theorem 1.11(4),  $[a, b|c, d] = [K] \cdot [L] \cdot \iota = [K] \cdot [M] \cdot [M] \cdot [L] \cdot \iota = \iota \cdot \iota \cdot \iota = \iota$ .

(2) By Theorem 1.11(5),  $\iota = -1$ . Therefore using the notations of Theorem 1.2(3) we have:

$$[a, b|c, d] = -[X] \cdot [Y], \quad [a, c|d, b] = -[X] \cdot [Z], \quad [a, d|b, c] = -[X] \cdot [U],$$

by Theorem 1.2(4) the quadruple  $(X, Y, Z, U)$  is a total square and so by **(O2)**, exactly two of the values  $[X], [Y], [Z], [U]$  are 1. But from this observations follows **(T3)**.  $\square$

### 3. From a separation to a halfordering of the corresponding field

We start from the assumption that the Minkowski plane  $\mathfrak{M}$  is provided with a separation  $\tau$ .

**Theorem 3.1.** *Let  $(a, b, c, d) \in P^{(4)}$  with  $a \neq b; c \neq d$  and let  $\gamma \in \Gamma$  then:*

1.  $[\gamma(a), \gamma(b)|\gamma(c), \gamma(d)] = [a, b|c, d].$
2.  $[a, b|c, d] \cdot [a, c|d, b] \cdot [a, d|b, c] = [a', b'|c', d']$  where  $(a', b', c', d')$  is any harmonic point quadruple.

*Proof.* (1) Follows from Proposition 1.8 and **(T2)**.

(2) Let  $C := \{bc, cb, a\}^\circ$  and  $D := \{bd, db, a\}^\circ$ . Then  $\tilde{C}(a) = a$ ,  $\tilde{C}(b) = c$ ,  $\tilde{C}(c) = b$  and  $\tilde{D}(a) = a$ ,  $\tilde{D}(b) = d$  and so by (1),  $[a, c|d, b] = [a, b|\tilde{C}(d), c]$  and  $[a, d|b, c] = [a, b|d, \tilde{D}(c)].$  By (T1) we have  $[a, b|c, d] \cdot [a, c|d, b] \cdot [a, d|b, c] = [a, b|c, d] \cdot [a, b|\tilde{C}(d), c] \cdot [a, b|d, \tilde{D}(c)] = [a, b|d, \tilde{C}(d)] \cdot [a, b|d, \tilde{D}(c)] = [a, b|\tilde{C}(d), \tilde{D}(c)].$  By Theorem 1.10,  $[a, b|\tilde{C}(d), \tilde{D}(c)]$  is a harmonic quadruple. By Theorem 1.9 and **(T2)** the  $\tau$ -value for all harmonic quadruples is the same.  $\square$

We recall that the commutative field  $(\mathbf{F}, +, \cdot)$  corresponding to the symmetric Minkowski plane  $\mathfrak{M}$  can be obtained as follows:

**Theorem 3.2.** *Let  $0, 1, \infty$  be three distinct points of a fixed circle  $E \in \Lambda$  and let  $\mathbf{F} := E \setminus \{\infty\}$ . For  $a, b \in \mathbf{F}$  and  $c \in \mathbf{F}^* := \mathbf{F} \setminus \{0\}$  let  $A^+$ ,  $C^- \in \Lambda$  be circles determined by:  $A^+ := E$  if  $a = 0$ ,  $0a \in A^+$  and  $A^+ \cap E = \{\infty\}$  if  $a \neq 0$ ,  $C^- := \{0, \infty, 1c\}^\circ$  and let  $\overline{A^+} := \widetilde{A^+E} \circ \widetilde{A^+E}|_E$ ,  $\overline{C^-} := \widetilde{C^-E} \circ \widetilde{C^-E}|_E$  [cf. Proposition 1.8(3)],  $a + b := \overline{A^+}(b)$ ,  $c \cdot b := \overline{C^-}(b)$ . Then:*

- (1)  $(\mathbf{F}, +, \cdot)$  is a commutative field.
- (2) The function  $\eta : \mathbf{F}^* \rightarrow \{1, -1\}$ ;  $x \mapsto [0, \infty|1, x]$  induced by the separation  $\tau$  is a halfordering of  $(\mathbf{F}, +, \cdot)$ .
- (3) If  $\tau$  is an order separation then  $\eta$  is an ordering of  $(\mathbf{F}, +, \cdot)$ .

*Proof.* (2) Let  $a, b \in \mathbf{F}^*$ . By the definition of  $\overline{A^+}$ ,  $\overline{A^+}(0) = 0$ ,  $\overline{A^+}(\infty) = \infty$ ,  $\overline{A^+}(1) = a$  and  $\overline{A^+}(b) = a \cdot b$  and so by Theorem 3.1,  $[0, \infty|1, b] = [0, \infty|a, a \cdot b]$  hence together with **(T1)**,  $\eta(a \cdot b) = [0, \infty|1, a \cdot b] = [0, \infty|1, a] \cdot [0, \infty|a, a \cdot b] = [0, \infty|1, a] \cdot [0, \infty|1, b] = \eta(a) \cdot \eta(b)$ .

Since  $\widetilde{A^+E}(\infty) = \infty$  we have by Theorem 3.1,  $[\infty, x|y, z] = [\infty, a+x|a+y, a+z]$  for  $x, y, z \in \mathbf{F}$ . Now let  $a \neq 0, 1$ . Then by using this formula and **(T2)**,

$$\begin{aligned}
[\infty, 0|1, a] &= \eta(a), \\
[\infty, 1|a, 0] &= [\infty, 0|a - 1, -1] = [\infty, 0|1, -1] \cdot [\infty, 0|1, a - 1] \\
&= \eta(-1) \cdot \eta(a - 1) = \eta(1 - a) \text{ and} \\
[\infty, a|0, 1] &= [\infty, 0| - a, 1 - a] = [\infty, 0|1, -a] \cdot [\infty, 0|1, 1 - a] \\
&= \eta(-a) \cdot \eta(1 - a) = \eta(a^2 - a) = \eta(a) \cdot \eta(a - 1).
\end{aligned}$$

(3) If  $\tau$  is an order separation then exactly one of these three values is equal  $-1$ . Hence if  $\eta(a) = 1$  then  $-\eta(1 - a) = \eta(a - 1) = \eta((-1) \cdot ((1 - a))) = \eta(-1) \cdot \eta(1 - a)$  thus  $\eta(-1) = -1$  and so  $\eta(-a) = \eta((-1) \cdot a) = -\eta(a)$ . Now let  $x, y \in \mathbf{F}^*$  with  $\eta(x) = \eta(y) = 1$ . For  $a := -x$  we obtain  $\eta(-x) = -1$  and so  $\eta(1 - a) = \eta(1 + x) = 1$ . By  $\eta(x^{-1} \cdot y) = (\eta(x))^{-1} \cdot \eta(y) = 1$ , this implies  $\eta(x + y) = \eta(x \cdot (1 + x^{-1}y)) = \eta(x) \cdot \eta(1 + x^{-1}y) = 1$  and this is the monotony law for the addition.  $\square$

Starting from a symmetric Minkowski plane  $\mathfrak{M}$  endowed with an orthogonal valuation [ ], we derived a separation  $\tau$  for  $\mathfrak{M}$  via Theorem 2.1 and from the separation  $\tau$ , a halfordering  $\eta_1$  for the corresponding field  $(\mathbf{F}, +, \cdot)$  via Theorem 3.2. We show that  $\eta_1$  and the halfordering  $\eta_2$  derived from  $(\mathfrak{M}, [ ])$  via [1](4.7) are equal. Let  $x \in \mathbf{F}^*$  and  $X := \{0\infty, \infty 0, 1x\}^\circ$ . Then  $X \perp E$ , by Theorem 1.11(1),  $[E] = 1$  and so  $\eta_1(x) = [0, \infty|1, x] = [E] \cdot [X] \cdot \iota = [X] \cdot \iota$  and moreover,  $\begin{pmatrix} 0 & 1 \\ x & 0 \end{pmatrix}$  is a matrix representing  $X$ . Therefore by [1](4.7),  $[X] = \eta_2(\det(\begin{pmatrix} 0 & 1 \\ x & 0 \end{pmatrix})) = \eta_2(-x) = \eta_2(-1) \cdot \eta_2(x)$  hence  $\eta_1(x) = \iota \cdot \eta_2(-1) \cdot \eta_2(x)$ . The circles  $E$  and  $I$  represented by the matrices  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  (with the determinants 1 and  $-1$ ) are orthogonal and containing the point 1. Hence  $(E, I) \in (\Lambda \times \Lambda)_{\perp s}$  and so by Theorem 1.11(4),  $\iota = [E] \cdot [I] = \eta_2(1) \cdot \eta_2(-1) = \eta_2(-1)$  and so  $\eta_1(x) = \eta_2(x)$  for all  $x \in \mathbf{F}^*$ . This gives us our main result:

**Theorem 3.3.** *Let  $\mathfrak{M}$  be a symmetric Minkowski plane and let  $(\mathbf{F}, +, \cdot)$  be the corresponding commutative field. Then there are a one to one correspondences between the orthogonal valuations of  $\mathfrak{M}$ , the separations of  $\mathfrak{M}$  and the halforderings of  $(\mathbf{F}, +, \cdot)$  where the order valuations correspond with the order separations and with the orderings of the field.*

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] Karzel, H., Kosiorek, J., Matraś, A.: *Ordered symmetric Minkowski planes I*. J. Geom. **93**, 116–127 (2009)
- [2] Kroll, H.-J.: *Anordnungsfragen in Benz-Ebenen*. Abh. Math. Semin. Univ. Hambg. **46**, 217–255 (1977)

Helmut Karzel  
Zentrum Mathematik  
T.U. München  
80290 Munich  
Germany  
e-mail: karzel@ma.tum.de

Jarosław Kosiorek and Andrzej Matraś  
Faculty of Mathematics and Computer Science  
University of Warmia and Mazury in Olsztyn  
Żołnierska 14  
10-561 Olsztyn  
Poland  
e-mail: kosiorek@matman.uwm.edu.pl  
e-mail: matras@uwm.edu.pl

Received: March 13, 2006.

Revised: December 17, 2009.