The degenerate two well problem for piecewise affine maps

Bernard Dacorogna, Paolo Marcellini and Emanuele Paolini

Abstract. The *two well problem* consists in finding maps u which satisfy some boundary conditions and whose gradient Du assumes values in the two wells $\mathbb{S}_A, \mathbb{S}_B$. Here $\mathbb{S}_A$ (similarly $\mathbb{S}_B$) is the *well* generated by a 2×2 matrix A , i.e., $\mathbb{S}_A$ is the set of matrices of the form RA , where R is a rotation. We study specifically the case when at least one of the two matrices A, B is singular and we characterize piecewise affine maps u satisfying almost everywhere the differential inclusion $Du(x) \in \mathbb{S}_A \cup \mathbb{S}_B$. In particular we describe the *lamination* and *angle* properties, which turn out to be different from those of the nonsingular case described in detail in [15]. We also show that the two well problem can be solved in some cases involving singular matrices, in strict contrast to the nonsingular (and not *orthogonal*) case.

Mathematics Subject Classification. 35F50 Nonlinear first-order systems.

Keywords. Partial differential equations, Two well problem, Differential inclusions.

1. Introduction

Arrigo Cellina, to whom this paper is dedicated, gave a relevant contribution to the field of *differential inclusions*, with application to several different types of problems, starting from the celebrated book [1] on set-valued maps, published in 1984 in collaboration with Jean-Pierre Aubin. In the context of *vector-valued maps*, which we consider in the research presented here, we mainly refer to the article [5], in collaboration with Stefania Perrotta, where a Dirichlet problem for differential inclusions involving *orthogonal matrices* is studied.

The differential inclusion problem is the following: let A, B be given 2×2 matrices and let us denote respectively by $\mathbb{S}_A$ and $\mathbb{S}_B$ the *wells* generated by A and B . That is, the set of matrices $\mathbb{S}_A$ (similarly $\mathbb{S}_B$) is the *well*, generated by A , of the form

$$\mathbb{S}_A = SO(2)A = \{RA : \text{where } R \in SO(2), \text{ i.e. } R \text{ is a rotation}\}.$$

For a given open set $\Omega \subset \mathbb{R}^2$ and a given boundary datum φ , the 2-dimensional *Dirichlet problem* for two wells consists in finding a map $u : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which satisfies the conditions

$$\begin{cases} Du(x) \in \mathbb{S}_A \cup \mathbb{S}_B & \text{a.e. } x \in \Omega \\ u(x) = \varphi(x) & x \in \partial\Omega. \end{cases} \quad (1)$$

Above Du represents the 2×2 gradient matrix of the map u . The *two well problem* is relevant in nonlinear elasticity and is a model for vector-valued differential inclusions. The original mathematical formulation is due to Ball and James [2, 3] (see also [19] and [9, 15] for details and references).

The case when A, B are *orthogonal matrices* has been first considered in the quoted paper [5] by Cellina and Perrotta (for 3×3 matrices and zero boundary condition). Is this a special case? Why to consider only *orthogonal* matrices A, B ?

The reason relies on the fact that it turns out to be impossible to solve the same two well problem in the same context, i.e., by mean of *piecewise affine maps*, unless we start from orthogonal matrices A, B . Precisely, we recently proved in [15] that the two well problem, for *nonsingular* matrices A and B , can be solved by means of piecewise affine maps if and only if

$$BA^{-1} \in O(2) \quad \text{and} \quad \det(BA^{-1}) = -1. \quad (2)$$

To give a more complete picture we recall that a map u , solving (1) with φ affine and satisfying some natural compatibility conditions, exists in the Sobolev class $W^{1,\infty}(\Omega; \mathbb{R}^2)$ of Lipschitz continuous maps (see [4, 6–9, 16, 18]). However, for *orthogonal matrices* it was also proved (see [5, 10–14, 17]) that solutions exist in the class of piecewise *affine* maps. As we said, in the class of nonsingular matrices A, B (i.e., with nonzero determinant) the Dirichlet problem (1)—for instance with zero boundary datum—can be solved, in the class of piecewise affine maps, (see [15]) essentially only if A and B are orthogonal matrices (namely, they satisfy (2)).

What about singular matrices A and/or B ? In fact this is a case which does not enter in the previous analysis. The aim of this paper is to show that if at least one of the two matrices has zero determinant then different properties happen to the geometry of the singularities of maps u satisfying the differential inclusion

$$Du(x) \in \mathbb{S}_A \cup \mathbb{S}_B, \quad \text{a.e. } x \in \Omega; \quad (3)$$

precisely, if u is a piecewise affine map as in (3), then its gradient matrix Du at every internal vertex of gradient-discontinuity satisfy new *lamination* and *angle conditions*, different from that one valid in the nonsingular case and described in detail in [15]. The new geometrical situation is proposed in Sect. 3 for one singular matrix and in Sect. 4 when both matrices A and B are singular.

Finally in Sect. 5 we show that the Dirichlet problem can be solved in some cases involving singular matrices, in strict contrast with respect to what happens for the nonsingular case, where the Dirichlet problem for the two well

problem lacks a solution in the class of piecewise affine maps, unless A and B are orthogonal matrices.

2. General notations

The set of 2×2 *orthogonal* matrices is denoted by $O(2)$, this is the set of $R \in \mathbb{R}^{2 \times 2}$ such that $R^t R = I$. The set of *special orthogonal* matrices, denoted by $SO(2)$ is the set of $R \in O(2)$ such that $\det R = 1$. We write a generic matrix in $SO(2)$ as

$$R_\varphi = \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix}$$

with $\varphi \in (-\pi, \pi]$.

We recall that the *singular values* of a 2×2 matrix A , denoted by

$$0 \leq \lambda_1(A) \leq \lambda_2(A),$$

are defined to be the square root of the eigenvalues of the symmetric and positive semidefinite matrix $A^t A \in \mathbb{R}^{2 \times 2}$. As well known, the singular values decomposition theorem asserts that for any $A \in \mathbb{R}^{2 \times 2}$, there exist $R, Q \in O(2)$ such that

$$RAQ = \text{diag}(\lambda_1(A), \lambda_2(A)) = \begin{pmatrix} \lambda_1(A) & 0 \\ 0 & \lambda_2(A) \end{pmatrix}.$$

In the sequel we will adopt the following notation and definitions.

Notation 1. If A is a 2×2 matrix, we denote with $\mathbb{S}_A$ the set of matrices

$$\mathbb{S}_A = SO(2) \cdot A = \{RA : R \in SO(2)\}.$$

Definition 2 (*Piecewise affine maps*). Let $\Omega \subset \mathbb{R}^2$ be an open set and let $u : \Omega \rightarrow \mathbb{R}^2$.

(i) We define the *singular set* of u as the set

$$\Sigma_u = \{x \in \Omega : u \text{ is not differentiable in } x\}.$$

(ii) We say that a map $u : \Omega \rightarrow \mathbb{R}^2$ is *piecewise affine* in Ω if

- u is continuous on Ω ,
- Σ_u is relatively closed in Ω ,
- $\Omega \setminus \Sigma_u$ has a finite number of connected components
- Du is constant on each connected component of $\Omega \setminus \Sigma_u$.

(iii) We say that u is *locally piecewise affine* if for every open set Ω' such that $\overline{\Omega'}$ is a compact subset of Ω , we have that u is piecewise affine in Ω' .

Definition 3. Let Σ be a locally finite union of closed segments in an open set $\Omega \subset \mathbb{R}^2$. We say that a point of Ω is a *vertex* of Σ if either it is an end point of a segment or a point where at least two segments meet.

3. The semi-degenerate case

In the present section we consider the case where one of the matrix is invertible, say A , and the other is not. We first fix the notations.

Notation 4. Let $\mu = \lambda_2(BA^{-1})$ be the largest singular value of BA^{-1} (the smallest being, under our hypotheses, necessarily 0). When $\mu \geq 1$ we let

$$m = \sqrt{\mu^2 - 1} \quad \text{and} \quad \nu_{\pm} = (\pm 1, \sqrt{\mu^2 - 1}) = (\pm 1, m).$$

Note that if $\mu = 1$ then $\nu_{\pm} = \pm e_1$. We also define $\theta \in [0, \pi/2)$ through

$$\cos \theta = \frac{1}{\mu} \quad \text{and} \quad \sin \theta = \frac{\sqrt{\mu^2 - 1}}{\mu} = \frac{m}{\mu}. \quad (4)$$

As in [15], with a change of variables we can reduce all of our analysis to the case where

$$A = I \quad \text{and} \quad B = \Lambda = \text{diag}(0, \mu),$$

with $\mu \geq 1$, to be consistent with (4). We will assume this special structure in all the following lemmas.

We now analyze when edges and vertices can occur and we start with edges.

Lemma 5 (Edge). Let $\nu = (\nu_1, \nu_2) \neq 0$ and $\varphi, \psi \in (-\pi, \pi]$. The map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu \rangle = x_1 \nu_1 + x_2 \nu_2 > 0 \\ R_{\psi} \Lambda x & \text{if } \langle x; \nu \rangle = x_1 \nu_1 + x_2 \nu_2 < 0 \end{cases}$$

is continuous across the line $\langle x; \nu \rangle = 0$ if and only if $\mu \geq 1$, ν is parallel to $\nu_{\pm}$ and $\psi = \pm \theta$. More precisely (see Fig. 1, middle case)

$$R_{-\varphi} u(x) = \begin{cases} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \text{if } x_1 + mx_2 > 0 \\ R_{\theta} \Lambda \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 & -m \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} -mx_2 \\ x_2 \end{pmatrix} & \text{if } x_1 + mx_2 < 0 \end{cases}$$

or (see Fig. 1, left hand case)

$$R_{-\varphi} u(x) = \begin{cases} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \text{if } x_1 - mx_2 < 0 \\ R_{-\theta} \Lambda \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 & m \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} mx_2 \\ x_2 \end{pmatrix} & \text{if } x_1 - mx_2 > 0. \end{cases}$$

Remark 6. (i) Note that within the well $\mathbb{S}_B$ there might be rank one connections, while this cannot happen within the well $\mathbb{S}_A$. More precisely if $\varphi, \psi \in (-\pi, \pi]$, then the map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$R_{-\varphi} u(x) = \begin{cases} \Lambda x & \text{if } \langle x; \nu \rangle = x_1 \nu_1 + x_2 \nu_2 > 0 \\ R_{\psi} \Lambda x & \text{if } \langle x; \nu \rangle = x_1 \nu_1 + x_2 \nu_2 < 0 \end{cases}$$

FIGURE 1. A scheme of 3 possible edges in the semi-degenerate case

is continuous across the line $\langle x; \nu \rangle = 0$ if and only if ν is parallel to $e_2 = (0, 1)$. In other words, for every $\varphi, \psi \in (-\pi, \pi]$, the map $R_{-\varphi} u$ does not depend on x_1 more precisely (see Fig. 1, right hand case)

$$R_{-\varphi} u(x) = \begin{cases} x = (0, \mu x_2) & \text{if } x_2 > 0 \\ ((-\mu \sin \psi) x_2, (\mu \cos \psi) x_2) & \text{if } x_2 < 0. \end{cases}$$

(ii) If $\mu = 1$, the lemma reads as

$$R_{-\varphi} u(x) = \begin{cases} x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \text{if } x_1 > 0 \\ \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ x_2 \end{pmatrix} & \text{if } x_1 < 0 \end{cases}$$

Proof. The map u is continuous only if

$$\det(I - R_\psi \Lambda) = 1 - \mu \cos \psi = 0.$$

This can happen if and only if $\mu \geq 1$ and $\psi = \pm\theta$ where

$$\cos \theta = \frac{1}{\mu} \quad \text{and} \quad \sin \theta = \frac{\sqrt{\mu^2 - 1}}{\mu} = \frac{m}{\mu}.$$

It remains to show that ν is parallel to $\nu_\pm = (\pm 1, m)$. We discuss only the case where $\psi = \theta$, the case $\psi = -\theta$ being handled similarly. Since the map is continuous across the line $\langle x; \nu \rangle = 0$, we should have

$$0 = (I - R_\psi \Lambda) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & \mu \sin \theta \\ 0 & 1 - \mu \cos \theta \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & m \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

This is equivalent to $x_1 + mx_2 = 0$ and thus ν is parallel to $\nu_+ = (1, m)$, as wished. $\square$

Recall that $\mu \geq 1$. If $\mu = 1$ then the directions $\nu_\pm = (\pm 1, 0)$ are parallel to e_1 while ν is parallel to e_2 , hence all the edges of the singular set are parallel to e_2 , respectively e_1 , and a vertex can occur either with a *T-shaped configuration* (i.e., 3 angles of measure $\pi/2, \pi/2$, and π) or with 4 orthogonal angles; a direct verification excludes both cases. Thus a vertex can exist only if $\mu > 1$.

We will show that these vertices can only be of order 3, 4, 5 and 6. We now consider separately all these cases. In the following lemmas we assume

that $\mu > 1$. The configurations considered in each lemma are schematized in Fig. 2 below.

Lemma 7 (Vertex of order 3). *Let $\varphi, \chi, \psi \in (-\pi, \pi]$. The map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by*

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle < 0 \\ R_\chi \Lambda x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \\ R_\psi \Lambda x & \text{otherwise} \end{cases}$$

is continuous, across the lines of discontinuities of the gradient, if and only if $\chi = -\psi = \theta$.

Remark 8. (i) Therefore the only possibility of having, under the hypotheses of the lemma, a continuous u is that (see Fig. 2, the four cases in (3a))

$$R_{-\varphi} u(x_1, x_2) = \begin{cases} (x_1, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 > 0 \\ R_\theta \Lambda x = (-mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_2 < 0 \\ R_{-\theta} \Lambda x = (mx_2, x_2) & \text{otherwise.} \end{cases}$$

(ii) There can also be, for example, (but it is essentially the same, the only difference is the opening of the angles) a vertex of order 3 of the form

FIGURE 2. Vertices of order 3, 4, 5 and 6

(see Fig. 2, the four cases in (3b))

$$R_{-\varphi} u(x_1, x_2) = \begin{cases} (mx_2, x_2) & \text{if } x_1 - mx_2 > 0 \text{ and } x_2 > 0 \\ (-mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_2 < 0 \\ (x_1, x_2) & \text{otherwise.} \end{cases}$$

Proof. We can apply Lemma 5 to the two lines of discontinuity ($\langle x; \nu_+ \rangle = 0$ and $\langle x; \nu_- \rangle = 0$) to find that

$$\chi = -\psi = \theta.$$

Remark 6 ensures the continuity across $x_2 = 0$. $\square$

Lemma 9 (Vertex of order 4). *Let $\varphi, \chi_1, \chi_2, \psi \in (-\pi, \pi]$. The map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by*

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle < 0 \\ R_{\chi_1} \Lambda x & \text{if } \langle x; \nu_- \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \\ R_{\chi_2} \Lambda x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \\ R_{\psi} \Lambda x & \text{if } \langle x; e_2 \rangle > 0 \end{cases}$$

is continuous, across the lines of discontinuities of the gradient, if and only if $\chi_1 = -\chi_2 = \theta$.

Remark 10. (i) Therefore the only possibility of having, under the hypotheses of the lemma, a continuous u is that (see Fig. 2, the first case in (4))

$$R_{-\varphi} u(x) = \begin{cases} (x_1, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 > 0 \\ R_{-\theta} \Lambda x = (mx_2, x_2) & \text{if } x_1 - mx_2 < 0 \text{ and } x_2 < 0 \\ R_{\theta} \Lambda x = (-mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_2 < 0 \\ R_{\psi} \Lambda x = ((-\mu \sin \psi) x_2, (\mu \cos \psi) x_2) & \text{if } x_2 > 0. \end{cases}$$

(ii) There can also be (but it is essentially the same, the only difference is the aperture of the angles) one other vertex of order 4 (see Fig. 2, the second case in (4)).

(iii) However there cannot be a vertex of order 4 of the form

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; \nu_- \rangle < 0 \\ R_{\chi_1} \Lambda x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; \nu_- \rangle > 0 \\ R_{\psi} x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle > 0 \\ R_{\chi_2} \Lambda x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle < 0 \end{cases}$$

i.e. necessarily $\chi_1 = -\chi_2 = \theta$ but no ψ can verify the third part

$$R_{-\varphi} u(x) = \begin{cases} (x_1, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_1 - mx_2 > 0 \\ (mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_1 - mx_2 < 0 \\ \begin{pmatrix} x_1 \cos \psi - x_2 \sin \psi \\ x_1 \sin \psi + x_2 \cos \psi \end{pmatrix} & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 < 0 \\ (-mx_2, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 > 0. \end{cases}$$

Proof. We can apply Lemma 5 to the two lines of discontinuity ($\langle x; \nu_+ \rangle = 0$ and $\langle x; \nu_- \rangle = 0$) to find that

$$\chi = -\psi = \theta.$$

Remark 6 ensures the continuity across $x_2 = 0$. $\square$

The proofs of the next two lemmas are similar to the ones above and we do not discuss the details.

Lemma 11 (Vertex of order 5). *Let $\varphi, \chi_1, \chi_2, \chi_3, \psi \in (-\pi, \pi]$. The map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by*

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle < 0 \\ R_{\chi_1} \Lambda x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle > 0 \\ R_{\psi} x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; \nu_- \rangle > 0 \\ R_{\chi_2} \Lambda x & \text{if } \langle x; \nu_- \rangle < 0 \text{ and } \langle x; e_2 \rangle > 0 \\ R_{\chi_3} \Lambda x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \end{cases}$$

is continuous, across the lines of discontinuities of the gradient, if and only if

$$\chi_1 = -\theta, \quad \chi_2 = -3\theta, \quad \chi_3 = \theta, \quad \psi = -2\theta.$$

Remark 12. (i) Therefore the only possibility, under the hypotheses of the lemma, to have a continuous u is that (see Fig. 2, the first case in (5))

$$R_{-\varphi} u(x_1, x_2) = \begin{cases} (x_1, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 > 0 \\ R_{-\theta} \Lambda x = (mx_2, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 < 0 \\ R_{-2\theta} x & \text{if } x_1 + mx_2 > 0 \text{ and } x_1 - mx_2 < 0 \\ R_{-3\theta} \Lambda x = R_{-2\theta}(mx_2, x_2) & \text{if } x_1 - mx_2 > 0 \text{ and } x_2 > 0 \\ R_{\theta} \Lambda x = (-mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_2 < 0. \end{cases}$$

(ii) There can also be another vertex of order 5 (essentially the symmetric one), see Fig. 2, the second case in (5).

Lemma 13 (Vertex of order 6). *Let $\varphi, \chi, \chi', \psi, \psi', \omega \in (-\pi, \pi)$. The map $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by*

$$R_{-\varphi} u(x) = \begin{cases} x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; \nu_- \rangle < 0 \\ R_{\chi} \Lambda x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \\ R_{\chi'} \Lambda x & \text{if } \langle x; \nu_- \rangle > 0 \text{ and } \langle x; e_2 \rangle < 0 \\ R_{\omega} x & \text{if } \langle x; \nu_+ \rangle > 0 \text{ and } \langle x; \nu_- \rangle > 0 \\ R_{\omega+\psi} \Lambda x & \text{if } \langle x; \nu_- \rangle < 0 \text{ and } \langle x; e_2 \rangle > 0 \\ R_{\omega+\psi'} \Lambda x & \text{if } \langle x; \nu_+ \rangle < 0 \text{ and } \langle x; e_2 \rangle > 0 \end{cases}$$

is continuous, across the lines of discontinuities of the gradient, if and only if

$$\chi = \psi' = -\chi' = -\psi = \theta.$$

Remark 14. Therefore the only possibility, under the hypotheses of the lemma, to have a continuous u is that (see Fig. 2, (6))

$$R_{-\varphi} u(x_1, x_2) = \begin{cases} (x_1, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_1 - mx_2 > 0 \\ R_{\theta} \Lambda x = (-mx_2, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_2 < 0 \\ R_{-\theta} \Lambda x = (mx_2, x_2) & \text{if } x_1 - mx_2 < 0 \text{ and } x_2 < 0 \\ R_{\omega} x = R_{\omega}(x_1, x_2) & \text{if } x_1 + mx_2 > 0 \text{ and } x_1 - mx_2 < 0 \\ R_{\omega-\theta} \Lambda x = R_{\omega}(mx_2, x_2) & \text{if } x_1 - mx_2 > 0 \text{ and } x_2 > 0 \\ R_{\omega+\theta} \Lambda x = R_{\omega}(-mx_2, x_2) & \text{if } x_1 + mx_2 < 0 \text{ and } x_2 > 0 \end{cases}$$

Finally we have the main result of the present section, which follows from the previous lemmas in a straightforward way. In the theorem we will assume that $A, B \in \mathbb{R}^{2 \times 2}$ with A invertible and $B \neq 0$ non-invertible (the case where

$B = 0$ is elementary, see Remark 16). We recall that $\mu = \lambda_2(BA^{-1})$ (while, of course, $\lambda_1(BA^{-1}) = 0$) and, if $\mu \geq 1$,

$$\nu_{\pm} = \left(\pm 1, \sqrt{\mu^2 - 1} \right) = (\pm 1, m).$$

We let $\nu \neq 0$ be a (uniquely defined up to rescaling) vector such that $B\nu = 0$.

Theorem 15 (The semi-degenerate case). *Let Ω be an open set of $\mathbb{R}^2$. Let $A, B \in \mathbb{R}^{2 \times 2}$ with A invertible and $B \neq 0$ non-invertible. Let $u: \Omega \rightarrow \mathbb{R}^2$ be a piecewise affine map satisfying*

$$Du \in \mathbb{S}_A \cup \mathbb{S}_B \quad \text{a.e. in } \Omega.$$

Case 1 $\mu > 1$. Then every edge of Σ_u is perpendicular to one of the three vectors: ν_+, ν_-, ν . Moreover if an edge perpendicular to $\nu_{\pm}$ is crossed, then the matrix Du changes from a rotation of A to a rotation of B or vice-versa. While if an edge perpendicular to ν is crossed, then the matrix Du changes between two different rotations of B . The segments can possibly meet in vertices of order 3, 4, 5 or 6. The possible local configurations around a vertex are the 13 cases depicted in Fig. 2.

Case 2 $\mu = 1$. In this case ν_+, ν_- are parallel each other and every edge of Σ_u is perpendicular either to ν_+ (and ν_-) or ν . Moreover if an edge perpendicular to $\nu_{\pm}$ is crossed, then the matrix Du changes from a rotation of A to a rotation of B or vice-versa. While if an edge perpendicular to ν is crossed, then the matrix Du changes between two different rotations of B . In this case no vertex is possible.

Case 3 $\mu < 1$. No interface between A and B can exist. Hence either $\Sigma_u = \emptyset$ or Σ_u is composed by segments perpendicular to ν and $Du(x) \in \mathbb{S}_B$ for a.e. $x \in \Omega$.

Remark 16. When A is invertible and $B = 0$, we are in the case of incompatible wells with no rank one connection inside each of the wells. Therefore if u is a piecewise C^1 solution of

$$Du \in \mathbb{S}_A \cup \mathbb{S}_B \quad \text{a.e. in } \Omega,$$

then u is affine and thus $\Sigma_u = \emptyset$.

Proof. If $\mu < 1$ then there is not a θ satisfying (4); hence no edge between A and B can exist, as stated in Lemma 5 and Remark 6. If $\mu = 1$ the directions $\nu_{\pm} = (\pm 1, 0)$ are parallel each other, hence all the edges of the singular set are parallel either to $\nu_{\pm}$ or to ν . As we already said a vertex cannot occur. We therefore discuss only the last possibility $\mu > 1$.

As in [15], by mean of a change of variables, we can reduce ourselves to the case when $A = I$ and $B = \text{diag}(0, \mu)$. By Lemma 5 and Remark 6 we know that every segment of Σ_u is perpendicular to one of the vectors: $\nu, \nu_{\pm}$. Then around a vertex of Σ_u we can have edges with 6 possible directions.

By deciding if a direction is or is not used, we obtain a total of $2^6 = 64$ possible configurations. However only 13 (cf. Fig. 2) of them are possible. To exclude the others we notice that the map u must satisfy the following conditions:

- (i) the configuration must have at least a vertex where at least two edges with different directions meet; i.e., we exclude the empty configuration, configurations with only one edge and configurations with two aligned edges;
- (ii) on both sides of the edges which are perpendicular to ν (the horizontal edges) the matrix Du must be a rotation of B ;
- (iii) crossing a side which is perpendicular to $\nu_{\pm}$ the matrix Du changes from a rotation of A to a rotation of B (or vice-versa);
- (iv) the configuration with order 4 where the two directions perpendicular to ν are not used is not possible in view of Remark 10 (iii). In fact this configuration would be possible only if $\lambda\mu = -1$ (see [15], where the two matrices are invertible), but we have here $\lambda = 0$.

Conditions (ii)–(iv) allow us to exclude the vertices represented in Fig. 3. Notice that in the configurations with 4 and 6 edges, the map is not uniquely determined even if we fix the value in one region. $\square$

We end this section with some other considerations. An example of a complete singular set Σ_u is depicted in Fig. 4.

Interestingly enough we can show, with the example depicted in Fig. 5, that a set Σ satisfying the local properties considered in the previous proposition, is not always the singular set of some Lipschitz continuous map u .

Thus in this case we cannot have a *recovery* theorem (as in Theorem 4.9 of [11] or in the *sufficient condition* of Theorem 25 in [15]). Moreover there are also cases (when we use the configuration with vertices of order 4 or 6) where a map with the prescribed singular set can be constructed but is not uniquely determined even if we fix the value in one region and the set Ω is simply connected.

FIGURE 3. The represented vertices cannot be in the singular set of any map u

FIGURE 4. An example of the singular set: in the dark regions the map has gradient in the non-singular well $\mathbb{S}_A$, while in the complementary regions its gradient belong to the singular well $\mathbb{S}_B$

FIGURE 5. A set Σ which locally satisfies the conditions stated in Lemmas 5, 7 at every vertex and every edge, but which does not correspond to the singular set of any map

4. The fully-degenerate case

We will now discuss the case where both matrices A and B are not invertible. We start with a very elementary lemma whose proof is straightforward.

Lemma 17. *Let $A \in \mathbb{R}^{2 \times 2}$ with $\det A = 0$. Then there exists $\alpha \in \mathbb{R}^2$ such that*

$$\mathbb{S}_A = (SO(2)e_1) \otimes \alpha.$$

Thus if $A, B \in \mathbb{R}^{2 \times 2}$ with $\det A = \det B = 0$. Then there exist $\alpha, \beta \in \mathbb{R}^2$ such that

$$\mathbb{S}_A \cup \mathbb{S}_B = (SO(2)e_1) \otimes \{\alpha, \beta\}.$$

We now have a lemma concerning edges between two different wells.

Lemma 18. *Let $\alpha, \beta \in \mathbb{R}^2$ and $\Omega \subset \mathbb{R}^2$ be an open set. Let u be piecewise affine and such that*

$$Du \in (SO(2)e_1) \otimes \{\alpha, \beta\} \quad \text{a.e. in } \Omega.$$

Then three cases can happen for the lamination between the two different wells (we do not consider here discontinuities in the same well; see the remark below).

Case 1 (*double laminations*). If the vectors α and β are linearly independent, then only two lines of discontinuity are possible; one with normal $\alpha + \beta$ and the other one with normal $\alpha - \beta$.

Case 2 (*single lamination*). If the vectors α and β are linearly dependent and $\alpha \neq 0$ (or $\beta \neq 0$), then only one line of discontinuity is possible, with normal α (respectively β).

Case 3 (*no lamination*). If $\alpha = \beta = 0$, then u is constant on every connected component.

Remark 19. As in the semi-degenerate case, lamination can occur within the same well. The line of discontinuities are then orthogonal to α and β respectively. For example in the first well we can have, for every $\varphi, \psi \in (-\pi, \pi]$,

$$u(x) = \begin{cases} (R_\varphi e_1 \otimes \alpha) x = \begin{pmatrix} (\alpha_1 x_1 + \alpha_2 x_2) \cos \varphi \\ (\alpha_1 x_1 + \alpha_2 x_2) \sin \varphi \end{pmatrix} & \text{if } \langle \alpha; x \rangle = \alpha_1 x_1 + \alpha_2 x_2 > 0 \\ (R_\psi e_1 \otimes \alpha) x = \begin{pmatrix} (\alpha_1 x_1 + \alpha_2 x_2) \cos \psi \\ (\alpha_1 x_1 + \alpha_2 x_2) \sin \psi \end{pmatrix} & \text{if } \langle \alpha; x \rangle = \alpha_1 x_1 + \alpha_2 x_2 < 0 \end{cases}$$

Proof. In Case 3 nothing is to be proved, so we discuss the two other cases. We start with the following observation. Along a line of discontinuity of the gradient we have that

$$Du = \begin{cases} R_\theta e_1 \otimes \alpha = \begin{pmatrix} \alpha_1 \cos \theta & \alpha_2 \cos \theta \\ \alpha_1 \sin \theta & \alpha_2 \sin \theta \end{pmatrix} \\ R_\varphi e_1 \otimes \beta = \begin{pmatrix} \beta_1 \cos \varphi & \beta_2 \cos \varphi \\ \beta_1 \sin \varphi & \beta_2 \sin \varphi \end{pmatrix}. \end{cases}$$

Since the map u is continuous across the line of discontinuity of the gradient we should have

$$\det(R_\theta e_1 \otimes \alpha - R_\varphi e_1 \otimes \beta) = \det(R_{(\theta-\varphi)} e_1 \otimes \alpha - e_1 \otimes \beta) = 0$$

which leads to

$$(\alpha_2 \beta_1 - \alpha_1 \beta_2) \sin(\theta - \varphi) = 0.$$

Therefore only three possibilities can happen

$$\alpha_2 \beta_1 - \alpha_1 \beta_2 = 0, \quad \varphi = \theta \quad \text{or} \quad \varphi = \theta + \pi.$$

Case 1 We now consider the case where α and β are linearly independent. According to the previous computations we must have $\varphi = \theta$ or $\varphi = \theta + \pi$. Therefore only two possibilities happen for the gradient

$$Du = \begin{cases} R_\theta e_1 \otimes \alpha = \begin{pmatrix} \alpha_1 \cos \theta & \alpha_2 \cos \theta \\ \alpha_1 \sin \theta & \alpha_2 \sin \theta \end{pmatrix} \\ R_\theta e_1 \otimes \beta = \begin{pmatrix} \beta_1 \cos \theta & \beta_2 \cos \theta \\ \beta_1 \sin \theta & \beta_2 \sin \theta \end{pmatrix} \end{cases}$$

or

$$Du = \begin{cases} R_\theta e_1 \otimes \alpha = \begin{pmatrix} \alpha_1 \cos \theta & \alpha_2 \cos \theta \\ \alpha_1 \sin \theta & \alpha_2 \sin \theta \end{pmatrix} \\ R_{(\theta+\pi)} e_1 \otimes \beta = R_\theta e_1 \otimes (-\beta) = -\begin{pmatrix} \beta_1 \cos \theta & \beta_2 \cos \theta \\ \beta_1 \sin \theta & \beta_2 \sin \theta \end{pmatrix}. \end{cases}$$

The two lines of discontinuities of the gradient are therefore orthogonal in the first case to $\alpha - \beta$ and in the second case to $\alpha + \beta$. More precisely in the first case the line of discontinuity is parallel to

$$(\alpha_1 - \beta_1)x_1 + (\alpha_2 - \beta_2)x_2 = 0$$

and in the second case the line is parallel to

$$(\alpha_1 + \beta_1)x_1 + (\alpha_2 + \beta_2)x_2 = 0$$

The case 1 is therefore settled.

Case 2 We now consider the case where α and β are linearly dependent. Assume, without loss of generality, that $\alpha \neq 0$, we therefore have $\beta = t\alpha$ and hence along a line of discontinuity of the gradient we have that

$$Du = \begin{cases} R_{\theta}e_1 \otimes \alpha \\ R_{\varphi}(te_1) \otimes \alpha. \end{cases}$$

Thus the only possible line of discontinuity has normal α . More precisely the line of discontinuity is parallel to the line

$$\alpha_1x_1 + \alpha_2x_2 = 0.$$

Case 2 is established and hence the proof of the theorem is complete. $\square$

5. The Dirichlet problem in the degenerate case

We finally discuss an example of the Dirichlet problem in the degenerate case. This turns out to be elementary. It is in fact essentially a reduction to a scalar problem. We emphasize that this example strictly contrasts with the nonsingular case; in fact, if A and B are invertible but not orthogonal matrices (in the sense of (2)), then we proved in [15] that the corresponding Dirichlet problem lacks a solution in the class of piecewise affine maps.

Example 20. Let a, b, c be positive real numbers. Let Ω be the rectangle in $\mathbb{R}^2$ with vertices in $(a, b), (-a, b), (a, -b), (-a, -b)$. Let us consider the degenerate matrices

$$A = \begin{pmatrix} \frac{c}{a} & 0 \\ \frac{c}{a} & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & \frac{c}{b} \\ 0 & \frac{c}{b} \end{pmatrix}.$$

Then there exists a piecewise (in fact with only four pieces) affine solution of

$$\begin{cases} Du(x) \in \mathbb{S}_A \cup \mathbb{S}_B & \text{a.e. } x \in \Omega \\ u(x) = 0 & x \in \partial\Omega. \end{cases}$$

Proof. We will solve a scalar differential problem for each component u^1, u^2 of the map $u : \Omega \rightarrow \mathbb{R}^2$, as in the more general “pyramid” construction in Theorem 2.10 in [9]. In this specific case we can exhibit an explicit solution given by $u = \begin{pmatrix} u^1 \\ u^2 \end{pmatrix}$ with

$$u^1(x_1, x_2) = u^2(x_1, x_2) = \begin{cases} c - \frac{c}{a}|x_1|, & \text{when } a|x_2| \leq b|x_1| \\ c - \frac{c}{b}|x_2|, & \text{when } a|x_2| > b|x_1| \end{cases}$$

whose gradient is given by

$$Du = \begin{pmatrix} u_{x_1}^1 & u_{x_2}^1 \\ u_{x_1}^2 & u_{x_2}^2 \end{pmatrix} = \pm \begin{pmatrix} \frac{c}{a} & 0 \\ \frac{c}{a} & 0 \end{pmatrix} = \pm A \in \mathbb{S}_A$$

when $a|x_2| < b|x_1|$, and similarly $Du = \pm B \in \mathbb{S}_B$ in the other case. $\square$

Acknowledgements

As we said above, this paper is dedicated to Arrigo Cellina. We know Arrigo since long time and we appreciate his mathematical rigor associated to fantasy and originality, which made the mathematics that he touched in his studies interesting to us, often attracting our interest and mathematical curiosity.

References

- [1] Aubin J.P., Cellina A. (1984) Differential inclusions. Grundlehren der Mathematischen Wissenschaften, vol. 264. Springer, Berlin
- [2] Ball, J.M., James, R.D.: Fine phase mixtures as minimizers of energy. Arch. Rational Mech. Anal **100**, 15–52 (1987)
- [3] Ball, J.M., James, R.D.: Proposed experimental tests of a theory of fine microstructure and the two well problem. Phil. Trans. Royal Soc. London A **338**, 389–450 (1991)
- [4] Celada, P., Perrotta, S.: Functions with prescribed singular values of the gradient. Nonlinear Differ. Equ. Appl. **5**, 383–396 (1998)
- [5] Cellina, A., Perrotta, S.: On a problem of potential wells. J. Convex Anal. **2**, 103–115 (1995)
- [6] Dacorogna, B., Marcellini, P.: Théorème d’existence dans le cas scalaire et vectoriel pour les équations de Hamilton-Jacobi. C. R. Acad. Sci. Paris Sér. I Math **322**, 237–240 (1996)
- [7] Dacorogna, B., Marcellini, P.: Sur le problème de Cauchy-Dirichlet pour les systèmes d’équations non linéaires du premier ordre. C. R. Acad. Sci. Paris Sér. I Math **323**, 599–602 (1996)
- [8] Dacorogna, B., Marcellini, P.: General existence theorems for Hamilton-Jacobi equations in the scalar and vectorial case. Acta Math. **178**, 1–37 (1997)
- [9] Dacorogna, B., Marcellini, P.: Implicit partial differential equations. Progress in Nonlinear Differential Equations and Their Applications, vol. 37. Birkhäuser, Boston (1999)
- [10] Dacorogna, B., Marcellini, P., Paolini, E.: An explicit solution to a system of implicit differential equations. Annales de l’I.H.P. Analyse Non linéaire **25**, 163–171 (2008)

- [11] Dacorogna, B., Marcellini, P., Paolini, E.: Lipschitz-continuous local isometric immersions: rigid maps and origami. *J. Math. Pures Appl* **90**, 66–81 (2008)
- [12] Dacorogna, B., Marcellini, P., Paolini, E.: On the n -dimensional Dirichlet problem for isometric maps. *J. Funct. Anal.* **255**, 3274–3280 (2008)
- [13] Dacorogna, B., Marcellini, P., Paolini, E.: Functions with orthogonal Hessian. *Diff. Integral Equ.* **23**, 51–60 (2010)
- [14] Dacorogna, B., Marcellini, P., Paolini, E.: Origami and partial differential equations. *Notices AMS* **57**, 598–606 (2010)
- [15] Dacorogna, B., Marcellini, P., Paolini, E.: The two well problem for piecewise affine maps. *Adv. Differ. Equ.* (2012, to appear)
- [16] De Blasi, F.S., Pianigiani, G.: On the Dirichlet problem for Hamilton-Jacobi equations: a Baire category approach. *NoDEA Nonlinear Differential Equations Appl.* **6**, 13–34 (1999)
- [17] Iwaniec, T., Verchota, G., Vogel, A.: The failure of rank one connections. *Arch. Ration. Mech. Anal.* **163**, 125–169 (2002)
- [18] Müller, S., Sverak, V.: Attainment results for the two-well problem by convex integration. In: Jost, J. (ed.) *Geometric analysis and the calculus of variations*, pp. 239–251. International Press, Somerville (1996)
- [19] Sverak, V.: On the problem of two wells. In: Ericksen J. et al. (eds.) *Microstructure and phase transitions*, IMA Vol. Appl. Math. vol. 54, pp. 183–189 (1993). Springer-Verlag, Berlin

Bernard Dacorogna
Section de Mathématiques
EPFL
1015 Lausanne
Switzerland

Paolo Marcellini and Emanuele Paolini
Dipartimento di Matematica “U. Dini”
Università di Firenze
Viale Morgagni 67/A
50134 Firenze
Italy
e-mail: marcellini@math.unifi.it

Received: 13 December 2011.

Accepted: 27 March 2012.