Editorial

H. Urbach

Published online: 10 November 2012
© Springer-Verlag Berlin Heidelberg 2012

Hardly a week goes by without some German TV show reporting on the misconduct of German physicians. Headlines like “nosocomial infections responsible for more deaths than traffic accidents” or “manipulated donor lists for organ transplantations” always focus on the fates of individual patients who received insufficient treatment or were even harmed at the hospital. The fact that these patients were also protected from serious diseases tends to be downplayed and some physicians are blamed even though they acted correctly.

It was only a matter of time before interventional neuroradiologists also found themselves in the crosshairs. We are accused of treating patients without using randomized controlled trials to ensure the treatment’s efficacy, despite the fact that, given the rapid evolution of the field of interventional neuroradiology, at least some randomized controlled trials become worthless before they are even finished. We are criticized for using different approval processes for medical products in the EU and the US. Stricter approval criteria than those established by the FDA might protect some patients from unexpected complications, which—with respect to flow diverter—were not foreseeable, are not fully understood, and therefore are hardly testable in animal models. However, stricter approval criteria may also rob patients of the chance to benefit from a new therapy.

The consequences become obvious if we consider the “history” of mechanical thrombectomy. With dissenting

opinions, the FDA approved the Merci retriever for thrombectomies in 2004. Eight years later, a randomized controlled trial, ISAT III, has failed to show the superiority of mechanical thrombectomy to i.v. lysis, and most devices used in the trial were Merci retrievers. What would be the role of thrombectomy today if skilled and enthusiastic German neuroradiologists had not started using the Solitaire stent for mechanical thrombectomy? The superiority of stentriever, namely, the Solitaire and Trevo stent, as compared with the Merci retriever has since been proven, and the Solitaire stent was approved by the FDA in April 2012. We all know many patients who were treated before FDA approval and who would raise their arms—assuming they can—if they were asked whether this was the right decision.

I neither want to criticize the FDA’s approval processes nor want to ignore the need for randomized controlled trials. However, offering no or ineffective treatment in cases where randomized controlled trials are not practicable, have not even started, or are still not finished cannot be the solution. Industrial registries are of great help, but it is our obligation to collect these data independently of these registries, to disseminate information, and to share not only all available data but also our personal experiences with the individual patient. Each physician–patient relationship ultimately comes down to a relationship between individuals, and neither party can do away with their responsibilities by hiding behind registries or even randomized controlled trials.

H. Urbach (✉)
Radiologische Klinik Neuroradiologie Universitätsklinikum
Bonn, Sigmund-Freud-Str. 25, 53105 Bonn, Germany
e-mail: Horst.Urbach@ukb.uni-bonn.de