13. Lin SM, Huang CD, Lin HC et al (2006) A modified goal-directed protocol improves clinical outcomes in intensive care unit patients with septic shock: a randomized controlled trial. *Shock* 26:551–557
14. Bouchard J, Soroko SB, Chertow GM et al (2009) Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury. *Kidney Int* 76:422–427
15. Grams ME, Estrella MM, Coresh J et al (2011) Fluid balance, diuretic use, and mortality in acute kidney injury. *Clin J Am Soc Nephrol* 6:966–973
16. Honore PM, Jacobs R, Joannes-Boyau O et al (2013) Con: Dialy- and continuous renal replacement (CRRT) trauma during renal replacement therapy: still under-recognized but on the way to better diagnostic understanding and prevention. *Nephrol Dial Transplant* 28:2723–2727
17. Carl DE, Grossman C, Behnke M et al (2010) Effect of timing of dialysis on mortality in critically ill, septic patients with acute renal failure. *Hemodial Int* 14:11–17
18. Bagshaw SM, Uchino S, Bellomo R et al (2009) Timing of renal replacement therapy and clinical outcomes in critically ill patients with severe acute kidney injury. *J Crit Care* 24:129–140
19. Rabindranath K, Adams J, Macleod AM, Muirhead N (2007) Intermittent versus continuous renal replacement therapy for acute renal failure in adults. *Cochrane Database Syst Rev* 3:CD003773
20. Vinsonneau C, Camus C, Combes A et al (2006) Continuous venovenous haemodiafiltration versus intermittent haemodialysis for acute renal failure in patients with multiple-organ dysfunction syndrome: a multicentre randomised trial. *Lancet* 368:379–385
21. Schneider AG, Bellomo R, Bagshaw SM et al (2013) Choice of renal replacement therapy modality and dialysis dependence after acute kidney injury: a systematic review and meta-analysis. *Intensive Care Med* 39(6):987–997
22. Palevsky PM, Zhang JH, O'Connor TZ et al (2008) Intensity of renal support in critically ill patients with acute kidney injury. *N Engl J Med* 359:7–20
23. Bellomo R, Cass A, Cole L et al (2009) Intensity of continuous renal-replacement therapy in critically ill patients. *N Engl J Med* 361:1627–1638
24. Joannes-Boyau O, Honore PM, Perez P et al (2013) High-volume versus standard-volume haemofiltration for septic shock patients with acute kidney injury (IVOIRE study): a multicentre randomized controlled trial. *Intensive Care Med* 39:1535–1546
25. Vesconi S, Cruz DN, Fumagalli R et al (2009) Delivered dose of renal replacement therapy and mortality in critically ill patients with acute kidney injury. *Crit Care* 13:R57

Med Klin Intensivmed Notfmed 2014 · 109:331–335
DOI 10.1007/s00063-014-0392-x
Online publiziert: 7.Juni 2014
© Springer-Verlag Berlin Heidelberg 2014

A.C. Hoppner¹ · W. Klingler²

¹ Klinik für Neurologie, Rehabilitationskrankenhaus Ulm

² Neuroanästhesie, Neurochirurgische Universitätsklinik im Bezirkskrankenhaus Günzburg

Erratum zu: Notfall- und intensivmedizinische Versorgung des Status epilepticus

In Abbildung 2 ist die Dosisangabe des Medikaments Clonazepam fehlerhaft. Die korrekte Dosierung lautet 0,014–0,028 mg/kgKG.

Die Autoren bitten, die korrekte Angabe zu beachten und den Fehler zu entschuldigen.

Die Autoren

Korrespondenzadressen

Dr. A.C. Hoppner
Klinik für Neurologie,
Rehabilitationskrankenhaus Ulm
Oberer Eselsberg 45, 89081 Ulm
anselm.hoppner@uni-ulm.de

PD Dr. W. Klingler
Neuroanästhesie, Neurochirurgische Universitätsklinik im Bezirkskrankenhaus Günzburg
Ludwig-Heilmeyer-Str. 2, 89312 Günzburg
werner.klingler@uni-ulm.de

Die Online-Version des Originalartikels können Sie unter <http://dx.doi.org/10.1007/s00063-014-0372-1> finden.