Erratum to: Strahlentherapie und Onkologie, Volume 186 (No. 10)

Erratum to:

Guckenberger M, Ok S, Polat B, Sweeney RA,
Flentje M. Toxicity after Intensity-Modulated,
Image-Guided Radiotherapy for Prostate Cancer.
Strahlenther Onkol 2010;186:535–43
(DOI 10.1007/s00066-010-2144-z)

Due to a technical error in the publishing process, several figures were inaccurately presented in Table 1. Please note the corrected data of “Tumor stage” and “Risk group.”

Table 1. Patient and treatment characteristics. PO: prostate only; PSA: prostate-specific antigen; PTV: planning target volume; TURP: transurethral resection of the prostate; WP: whole pelvis.

Tabelle 1. Patienten- und Behandlungscharakteristika. PO: nur Prostata; PSA: prostataspezifisches Antigen; PTV: Planungszielvolumen; TURP: transurethrale Prostataresektion; WP: gesamtes Becken.

	PO (n = 75)		WP (n = 25)	
Age (years), median (range)	73 (54–80)		73 (58–80)	
PSA level (ng/ml), median (range)	7.7 (1.5–32)		24.8 (3.2–118)	
	Patients		Patients	
	n	%	n	%
Gleason Score				
• 2–6	38	50.7	5	20.0
• 7	33	44.0	9	36.0
• 8–9	3	4.0	10	40.0
• Unknown	1	1.3	1	4.0
Tumor stage				
• T1	46	61.3	6	24.0
• T2	27	36.0	11	44.0
• T3	2	2.7	7	28.0
• T4	0	0.0	1	4.0
Androgen suppression	23	30.7	16	64.0
TURP	19	25.3	6	24.0
Risk group				
• Low-risk	29	38.7	0	0.0
• Intermediate-risk	29	38.7	3	12.0
• High-risk	17	22.7	22	88.0
Total dose (mean dose PTV-2)				
• 73.91 Gy	24	32.0	2	8.0
• 76.23 Gy	51	68.0	23	92.0

Strahlenther Onkol 2010;186:705
DOI 10.1007/s00066-010-7144-5
Published Online: November 29, 2010