Enhanced functionality of T cell receptor-redirected T cells is defined by the transgene cassette

Matthias Leisegang · Boris Engels · Peter Meyerhuber ·
Elisa Kieback · Daniel Sommermeyer · Shao-An Xue ·
Simone Reuß · Hans Stauss · Wolfgang Uckert

Published online: 29 May 2008
© Springer-Verlag 2008

Erratum to: J Mol Med DOI 10.1007/s00109-008-0317-3

It has been brought to our attention by the authors that the following corrections are required for the above referenced manuscript.

The sentence on page 5, right column, first sentence after subheading should read: “*To confirm the previous findings for a human TCR, the same cassettes were used to compare the expression of TCR WT-1.*”

On page 6, in the chapter “Cell culture”, the sentence concerning the composition of murine T cell medium should read as follows: “*The murine cell lines 58 [25] and B3Z [26] and murine spleen cells were grown in RPMI 1640 medium (Invitrogen, Karlsruhe, Germany) supplemented with 10% FCS (PAN Biotech, Aidenbach, Germany), 50 μ M 2-mercaptoethanol (Sigma-Aldrich, Seelze, Germany), 1 mM sodium pyruvate, 100 IU/ml penicillin, streptomycin and 1 mM HEPES (all: Invitrogen).*”

The authors like to thank Chiron (Marburg, Germany) for providing Interleukin-2.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00109-008-0317-3>.

M. Leisegang · E. Kieback · D. Sommermeyer · S. Reuß ·
W. Uckert (✉)
Max-Delbrück-Center for Molecular Medicine,
Robert-Rössle-Straße 10,
13092 Berlin, Germany
e-mail: wuckert@mdc-berlin.de

B. Engels
Department of Pathology
and Committee on Immunology,
University of Chicago,
Chicago, Illinois, USA

P. Meyerhuber · W. Uckert
Institute of Biology,
Humboldt-University of Berlin,
Berlin, Germany

S.-A. Xue · H. Stauss
Royal Free Hospital,
Department of Immunology
and Molecular Pathology,
University College London,
London, UK