Microspore culture preferentially selects unreduced (2n) gametes from an interspecific hybrid of *Brassica napus* L. $\times$ *Brassica carinata* Braun

Matthew N. Nelson · Annaliese S. Mason ·
Marie-Claire Castello · Linda Thomson ·
Guijun Yan · Wallace A. Cowling

Published online: 7 August 2009
© Springer-Verlag 2009

Erratum to: Theor Appl Genet (2009) 119:497–505
DOI 10.1007/s00122-009-1056-8

Unfortunately, the author's name Nagaharu U, was wrongly typeset both in the Introduction and in the References. The correct citation in the Introduction should be (U 1935) and not (Nagaharu 1935) and the correct reference is as given below:

U N (1935) Genome analysis in *Brassica* with special reference to the experimental formation of *B. napus* and peculiar mode of fertilization. Jpn J Bot 7:389–452.

The online version of the original article can be found under doi: [10.1007/s00122-009-1056-8](https://doi.org/10.1007/s00122-009-1056-8).

M. N. Nelson (✉) · A. S. Mason · M.-C. Castello · L. Thomson ·
G. Yan · W. A. Cowling
Faculty of Natural and Agricultural Sciences,
School of Plant Biology, The University of Western Australia,
35 Stirling Highway, Crawley, WA 6009, Australia
e-mail: mnelson@plants.uwa.edu.au