

## Coenzyme Q<sub>10</sub> improves endothelial dysfunction of the brachial artery in Type II diabetes mellitus

G. F. Watts, D. A. Playford, K. D. Croft, N. C. Ward, T. A. Mori, V. Burke

Department of Medicine, University of Western Australia, Royal Perth Hospital, Perth, Australia

### Abstract

*Aim/hypothesis.* We assessed whether dietary supplementation with coenzyme Q<sub>10</sub> improves endothelial function of the brachial artery in patients with Type II (non-insulin-dependent) diabetes mellitus and dyslipidaemia.

*Methods.* A total of 40 patients with Type II diabetes and dyslipidaemia were randomized to receive 200 mg of coenzyme Q<sub>10</sub> or placebo orally for 12 weeks. Endothelium-dependent and independent function of the brachial artery was measured as flow-mediated dilatation and glyceryl-trinitrate-mediated dilatation, respectively. A computerized system was used to quantitate vessel diameter changes before and after intervention. Arterial function was compared with 18 non-diabetic subjects. Oxidative stress was assessed by measuring plasma F<sub>2</sub>-isoprostane concentrations, and plasma antioxidant status by oxygen radical absorbance capacity.

*Results.* The diabetic patients had impaired flow-mediated dilatation [3.8% (SEM 0.5) vs 6.4% (SEM 1.0),  $p = 0.016$ ], but preserved glyceryl-trinitrate-mediated dilatation, of the brachial artery compared with non-di-

abetic subjects. Flow-mediated dilatation of the brachial artery increased by 1.6% (SEM 0.3) with coenzyme Q<sub>10</sub> and decreased by -0.4% (SEM 0.5) with placebo ( $p = 0.005$ ); there were no group differences in the changes in pre-stimulatory arterial diameter, post-ischaemic hyperaemia or glyceryl-trinitrate-mediated dilatation response. Coenzyme Q<sub>10</sub> treatment resulted in a threefold increase in plasma coenzyme Q<sub>10</sub> ( $p < 0.001$ ) but did not alter plasma F<sub>2</sub>-isoprostanes, oxygen radical absorbance capacity, lipid concentrations, glycaemic control or blood pressure.

*Conclusion/interpretation.* Coenzyme Q<sub>10</sub> supplementation improves endothelial function of conduit arteries of the peripheral circulation in dyslipidaemic patients with Type II diabetes. The mechanism could involve increased endothelial release and/or activity of nitric oxide due to improvement in vascular oxidative stress, an effect that might not be reflected by changes in plasma F<sub>2</sub>-isoprostane concentrations. [Diabetologia (2002) 45: 420–426]

**Keywords** Coenzyme Q<sub>10</sub>, endothelial function, nitric oxide, diabetes.

Received: 14 August 2001 and in revised form: 15 November 2001

*Corresponding author:* G. F. Watts, PhD, MD, Department of Medicine, University of Western Australia, Royal Perth Hospital, GPO Box X2213, Perth, WA 6847, Australia.

e-mail: gfwatts@cyllene.uwa.edu.au

*Abbreviations:* CoQ, Coenzyme Q<sub>10</sub>; FMD, flow-mediated dilatation; NMD, glyceryl-trinitrate-mediated dilatation; ORAC, oxygen radical absorbance capacity

In Type II (non-insulin-dependent) diabetes mellitus the pathogenesis of vascular disease, its most common complication, remains unclear [1]. Endothelial dysfunction reflects the disordered physiology of several endothelium-derived vasoactive factors, in particular nitric oxide [2]. Endothelial dysfunction occurs commonly in diabetes and is an early feature of vasculopathy [3, 4]. Increased oxidative stress due to the effects of hyperglycaemia and its sequelae is a recognized feature of diabetes [5]. It might cause endothelial dysfunction through the inactivation and

decreased synthesis of nitric oxide by reactive oxygen species [6] such as superoxide.

While epidemiology suggests that conventional antioxidant vitamins can benefit vascular disease, the evidence from controlled clinical trials is less secure [7], especially in diabetes [8]. Vitamin E supplementation has not consistently been shown to improve endothelial-dependent vasodilator tone in diabetic patients [9, 11]. Moreover, with vitamin C, the only study reported in Type II diabetes was not placebo-controlled and involved acute intra-arterial administration of this antioxidant [12].

Coenzyme Q<sub>10</sub> (CoQ) is a critical intermediate of the mitochondrial electron transport chain that regulates cytoplasmic redox potential and it can inhibit superoxide generation by endothelial cells [13, 14]. CoQ is a more powerful antioxidant than vitamin E, inhibiting its pro-oxidant activity [15, 16]. Its deficiency can occur in diabetes in relation to impaired mitochondrial substrate metabolism [17] and increased oxidative stress [6]. Mitochondrial CoQ deficiency could be involved in the pathogenesis of Type II diabetes by impairing beta-cell function [18]. Low serum CoQ concentrations have been negatively correlated with poor glycaemic control and diabetic complications [19, 20]. Accordingly, some clinical trials have shown that CoQ could improve glycaemic control and blood pressure in diabetes [14, 18, 21, 22]. Hence, CoQ could have a potential role in the treatment of diabetes and its complications [14, 18, 23]. However, to date no studies have reported the effects of CoQ on vascular dysfunction or cardiovascular disease in diabetes.

We report a placebo-controlled observation of the effect of CoQ supplementation on vascular function of the peripheral circulation in Type II diabetic patients. Endothelial function was quantitated as postischaemic flow-mediated dilatation of the brachial artery using a new edge-detection software system that increases the precision of measurements [24]. We also measured potential changes in oxidative stress by measuring plasma F<sub>2</sub>-isoprostanes and measuring the oxygen radical absorbance capacity of plasma.

## Subjects and methods

**Subjects.** A total of 40 patients with Type II diabetes diagnosed by standard criteria and with dyslipidaemia were recruited from the community. Dyslipidaemia was defined as a fasting serum triglyceride of greater than 1.8 mmol/l or HDL cholesterol of less than 1.0 mmol/l with total cholesterol of less than 6.5 mmol/l and a total cholesterol-to-HDL cholesterol ratio of more than 4. Patients were excluded based on the following criteria: age older than 75 years, BMI greater than 40 kg/m<sup>2</sup>, history of myocardial infarction or stroke, insulin therapy, smoking, macroalbuminuria, serum creatinine greater than 150 µmol/l, liver abnormalities, use of antioxidants or lipid-regulating therapy, uncontrolled hypertension (> 160/

90 mmHg), and treatment with angiotensin-converting enzyme inhibitors, calcium antagonists or aspirin. Volunteers underwent a clinical examination, urinalysis and a 12-lead ECG. The vascular function in the diabetic patients was compared with 18 healthy, non-diabetic normolipidaemic subjects of similar age [mean age 54 years (SD 12.0); cholesterol 5 mmol/l (SD 0.4), HDL cholesterol 1.6 (0.3), triglyceride 0.9 (95% CI 0.8, 1.1)].

**Study design.** This study is part of a larger study examining the effects of CoQ and lipid-regulating therapies on vascular function of peripheral arteries measured by several techniques. We report on the effect of CoQ monotherapy on postischaemic dilation of the brachial artery. Eligible patients entered a run-in period of 6 weeks during which they were instructed to consume an isocaloric fat-modified diet of constant antioxidant composition. They then underwent the brachial artery reactivity test described below, after which they were randomized double-blind to treatment with either CoQ (Blackmores, Sydney, NSW, Australia) or matching placebo in a trial of 12 weeks duration. Coenzyme Q<sub>10</sub> (200 mg) was taken as two 50 mg capsules orally twice a day. Volunteers were interviewed every 2 weeks to assess their compliance with therapeutic units and brachial artery reactivity was re-studied 12 weeks after randomization. The Ethics Committee of the Royal Perth Hospital approved the study and all volunteers gave their written consent.

**Laboratory methods.** Venous blood was collected after a 12 h fast at baseline and at 12 weeks. Serum total cholesterol, triglyceride and HDL cholesterol were measured using enzymatic, colorimetric methods (Boehringer Mannheim, Mannheim, Germany) on a Hitachi 917 biochemical analyser (Hitachi, Tokyo, Japan). High-density lipoprotein cholesterol was measured after precipitation of apolipoprotein B-100 (apoB) containing lipoproteins with dextran sulphate. Low-density lipoprotein cholesterol was estimated by the Friedewald formula and by a direct assay when triglycerides were more than 3.5 mmol/l. The particle size of LDL was estimated by non-denaturing gel electrophoresis. Glycated haemoglobin (HbA<sub>1c</sub>) was measured by high performance liquid-chromatography (HPLC, BioRad Laboratories, Sydney, Australia). Plasma glucose and insulin were assayed using an enzymatic method (Boehringer) and an automated immuno-enzymometric assay (Tosoh, Kyobashi, Tokyo, Japan), respectively. Serum and urinary creatinine were measured by the modified Jaffe reaction. Total serum CoQ concentration was assayed by reverse-phase high performance liquid-chromatography using electrochemical detection [25]. Plasma F<sub>2</sub>-isoprostanes were analysed using gas-chromatography mass-spectrometry with electron capture negative chemical ionization [26]. Plasma oxygen radical absorbance capacity (ORAC) was measured using a fluorescent assay using a Trolox standard [27].

**Brachial artery ultrasonography.** Brachial artery ultrasonography was carried out [28]. Briefly, a 12-megahertz transducer connected to an Acuson Aspen ultrasound (Acuson, Mountain View, Calif, USA) and fixed in position by a stereotactic clamp 5 to 10 cm proximal to the ante-cubital crease was used to image the brachial artery. Images were recorded before and after vasodilatory stimuli and recorded on s-VHS videotape (Sony MQSE 180). Continuous ECG monitoring was done in all studies. Reactive hyperaemia of the brachial artery was induced after release of a pneumatic tourniquet placed around the left forearm and inflated to 50 mmHg above systolic blood pressure for 5 min. Pulse wave Doppler flow velocities were used to derive flow rate (ml/min) pre-reactive and post-reactive.

**Table 1.** Clinical, biochemical and vascular characteristics of the patients in the placebo and CoQ groups at baseline

Characteristics	Placebo group	CoQ group
<i>n</i> (male/female)	13/2	18/2
Age (years)	54.1 (10.4)	52.7 (6.2)
BMI (kg/m <sup>2</sup> )	31.3 (5.4)	29.9 (3.3)
SBP (mmHg)	139.1 (15.1)	128.0 (18.4)
DBP (mmHg)	81.0 (5.8)	75.8 (9.3)
Glucose (mmol/l)	6.9 (2.1)	8.2 (2.7)
HbA <sub>1c</sub> (%)	6.2 (0.8)	6.9 (1.4)
Insulin (mU/l)	15.1 (7.4)	12.2 (5.4)
Cholesterol (mmol/l)	5.3 (0.6)	5.3 (0.9)
Triglyceride (mmol/l)	2.5 (2.1, 3.0)	2.0 (1.7, 2.5)
HDL-cholesterol (mmol/l)	1.00 (0.08)	0.95 (0.15)
LDL-cholesterol (mmol/l)	3.2 (0.7)	3.2 (0.9)
LDL size (nm)	25.3 (0.8)	25.0 (0.8)
Coenzyme Q <sub>10</sub> (mmol/l)	1.2 (0.3)	1.5 (0.3)*
ORAC activity (μmol/l)	3638 (3270, 4046)	3723 (3275, 4233)
Plasma F <sub>2</sub> -isoprostanes (pmol/l)	1297 (1023, 1643)	1102 (892, 1361)
Brachial artery:		
Baseline diameter (mm)	4.3 (0.6)	4.2 (0.5)
Resting blood flow (ml/min)	186.6 (72.8)	205.9 (113.7)
Reactive hyperaemia (%)	462.2 (239.8)	431.9 (254.7)
Flow-mediated dilatation (%)	4.5 (3.0)	2.8 (3.0)
Nitrate-mediated dilatation (%)	16.9 (6.4)	16.3 (5.2)

Values are means (SD) or geometric means and 95% CI  
\**p* = 0.03 vs placebo group

tive hyperaemia. After the brachial artery diameter returned to baseline 400 μg glyceryl trinitrate was administered sublingually to assess endothelium-independent vasodilatory response. All images obtained were assessed blindly by two independent observers and only pairs of scans that were of consistently acceptable quality were included in this analysis. Analysis of flow-mediated dilatation (FMD) and glyceryl-trinitrate-mediated dilatation (NMD) of the brachial artery was carried out using a semi-automated edge detection software system [24] and operated by an experienced observer, who was blinded to the treatment group assignment. Responses were calculated as percentage change in brachial artery diameter from baseline. The analytical (intra-observer) coefficient of variation of the computerized technique is 6.7% compared with 32.5% using ultrasonic callipers, a more conventional visual estimation. The resolving power of the method tested on 'phantom arteries' is 8.3 μm.

**Statistical methods.** Data were analysed parametrically after logarithmic transformation of variables where appropriate. Discrete variables were compared by Chi-square test. Treatment effects were analysed using general linear models with adjustments for baseline values and resting brachial artery diameter. Association between variables were examined by linear regression methods. Statistical significance was defined at a *p* value of less than 0.05.

## Results

The diabetic patients were mostly middle-aged men and were on average, overweight, normotensive, in good glycaemic control and had typical diabetic dyslipidaemia. Of the 40 patients randomized, 39 completed the study and one withdrew due to an incidental illness. High quality ultrasound images were obtained in all patients randomized to CoQ. However, satisfactory image quality was not obtained in ultrasound scans from four patients randomized to placebo. Thus, our analysis refers to the residual 35 diabetic patients. Their characteristics did not differ from those withdrawn from the study.

Plasma F<sub>2</sub>-isoprostanes were not different between the diabetic patients and the non-diabetic control subjects [1245 μmol/l (95% CI 1075, 1442) vs 1310 (1085, 1583), *p* = 0.670]. However, plasma ORAC was lower in the diabetics patients than in the control subjects [3766 μmol/l (95% CI 3476, 4081) vs 4729 (4551, 4914) *p* < 0.001].

Compared with the control subjects, the diabetic patients overall had lower post-ischaemic FMD of the brachial artery [3.8% (SEM 0.5) vs 6.4 (1.0), *p* = 0.016] but similar NMD responses [16.5% (SEM 0.9) vs 18.5 (1.7), *p* = 0.291].

No differences were shown in the characteristics (Table 1) between patients randomized to placebo and CoQ treatment (*p* > 0.05), except for higher baseline plasma CoQ concentrations (*p* = 0.03) in the CoQ group. Only four patients in the study took metformin, one in the placebo and three in the CoQ group. The plasma glucose immediately before ultrasonography did not differ (*p* = 0.20) between the groups (Table 1).

The changes in baseline diameter, reactive hyperaemia, FMD and NMD of the brachial artery in the CoQ were compared with the placebo group (Table 2). There was an improvement in FMD of the brachial artery in the CoQ compared with the placebo group, without changes or differences in baseline arterial diameter, reactive hyperaemia or NMD. In an analysis with absolute FMD (mm) at 12 weeks as the dependent variable, there was still a favourable treatment effect of CoQ (*p* = 0.002) after adjusting for baseline brachial artery diameter (mm) at 12 weeks and pre-randomization percentage FMD response. This analysis together with the lack of significant change in resting brachial artery diameter (Table 2) demonstrates that the favourable effect of CoQ on percentage FMD response was independent of changes in basal brachial artery tone. There were no differences between the groups in the plasma glucose concentration immediately before ultrasonography post-intervention [8.2 mmol/l (SEM 0.8) for CoQ vs 7.3 mmol/l (0.7) for placebo, *p* = 0.43].

Treatment with CoQ was associated with an increase in plasma CoQ concentrations, from

**Table 2.** Changes in baseline diameter, reactive hyperaemia, flow-mediated dilatation (FMD) and nitrate-mediated dilatation (NMD) of the brachial artery in the diabetic patients treated with placebo or CoQ supplementation for 12 weeks

Variable	Placebo group	CoQ group	<i>p</i> value
Change in baseline artery diameter (mm)	0.05 (0.09)	-0.02 (0.08)	0.611
Change in resting blood flow (ml/min)	-43.3 (19.7)	20.6 (30.4)	0.136
Change in reactive hyperaemia (%)	85.0 (38.6)	-27.3 (84.5)	0.316
Change in FMD (%)	-0.4 (0.5)	1.6 (0.3)	0.005
Change in NMD (%)	-0.1 (1.4)	0.4 (1.2)	0.771

Values are means  $\pm$  SEM

1.3 mmol/l (SEM 0.1) to 4.8 (0.4),  $p < 0.001$ ; however, there were no alterations ( $p > 0.05$ ) in plasma F<sub>2</sub>-isoprostanes, plasma ORAC activity, glucose, HbA<sub>1c</sub>, plasma lipids, blood pressure or other variables (Table 1). Changes in plasma CoQ concentrations were not correlated with the changes in plasma F<sub>2</sub>-isoprostanes. The FMD of the brachial artery increased with CoQ alone from 2.8% (SEM 0.7) to 4.4% (0.5),  $p < 0.001$  but the post-treatment response tended to stay lower ( $p = 0.07$ ) than the values for the non-diabetic control group. There was no correlation between the change in FMD and change in other variables in the CoQ group. In data pooled from both patient groups, improvement in FMD was only correlated with treatment group assignment to CoQ.

## Discussion

Our randomized, double-blind study shows a favourable effect of oral CoQ supplementation on endothelial dysfunction of the peripheral circulation in patients with Type II diabetes. Coenzyme Q<sub>10</sub> improved abnormal endothelium-dependent vasodilator tone of the brachial artery without altering the vasodilatory response to the endothelium-independent agonist glyceryl trinitrate. This favourable effect of CoQ was independent of changes in resting brachial artery diameter. The improvement in endothelial function also occurred in the presence of dyslipidaemia and was not related to changes in plasma F<sub>2</sub>-isoprostanes, glycated haemoglobin or blood pressure.

That oxidative stress is increased in diabetes is well supported by experimental and clinical observations [5, 6, 11, 29, 30]. Oxidative stress occurs in diabetes as a consequence of several mechanisms related to hyperglycaemia [5, 6]. These include accumulation of AGE, activation of the polyol pathway and stimulation of protein kinase C activity. Diabetes-induced generation of reactive oxygen species, in particular superoxide, decreases the expression of nitric oxide synthase and inactivates nitric oxide [6, 31]. Vascular oxidative stress could explain why our patients had impaired FMD of the brachial artery with preserved vasodilation to glyceryl trinitrate. Unexpectedly, our patients did not show evidence of increased oxidative stress, as measured by plasma concentrations of F<sub>2</sub>-

isoprostanes. Plasma F<sub>2</sub>-isoprostanes reflect non-enzymic, free radical induced lipid peroxidation and might not be sensitive to increases in oxidative stress at the vascular wall. Previously reported increases in plasma F<sub>2</sub>-isoprostanes and their diminution in diabetes with antioxidant vitamins referred to patients with poorer glycaemic control than our study group [29, 30]. We found that plasma antioxidant capacity was decreased in our patients in agreement with previous reports showing that diabetic patients have decreased plasma concentrations of antioxidant vitamins [32]. However, low plasma ORAC was not corrected by CoQ supplementation, suggesting that CoQ might not contribute to plasma antioxidants measured by this assay. A dissociation between changes in endothelial function and plasma F<sub>2</sub>-isoprostane concentrations was found using antioxidant vitamin supplementation in non-diabetic subjects [33]. Our patients were dyslipidaemic and it is possible that increased plasma concentrations of lipoprotein remnants, small-dense LDL and low HDL cholesterol might have contributed to endothelial dysfunction in the absence of a systemic increase in oxidative stress [6].

There is evidence in Type II diabetes for and against improvement in vasodilator function of forearm resistance arteries in response to the muscarinic agonist acetylcholine with vitamin E supplementation [10, 11]. One positive study using a vitamin E analogue was not placebo controlled but did show reduction in plasma F<sub>2</sub>-isoprostanes in a small number of patients [11]. Vitamin E supplementation was reported to improve forearm microcirculatory function in a larger sample of Type II diabetic patients in a well-controlled trial [10]. Improvement in methacholine-mediated vasodilator function of forearm resistance vessels was also reported in Type II diabetes following the intra-arterial administration of vitamin C [12]. However, that study used a small sample size, involved an acute intervention, and did not use a placebo arm. A recent report has shown that intra-arterial administration of the powerful antioxidant  $\alpha$ -lipoic acid improved forearm blood flow responses to acetylcholine to the same extent as ascorbic acid in patients with Type II diabetes [34]. The greatest benefit was seen in patients with low-plasma concentration of CoQ, supporting an important role of CoQ in vascular endothelial dysfunction in Type II diabetes. We

have extended previous reports by investigating intervention with the antioxidant CoQ in a larger sample of patients and in a peripheral conduit artery. In contrast to forearm microcirculatory blood flow responses to acetylcholine, post-*ischaemic* FMD of the brachial artery has been shown to be a surrogate for the coronary circulation and to predict coronary events in patients with angina [35, 36].

The beneficial properties of CoQ could relate not only to its antioxidant effect [15, 16], but also to improvements in glycaemic control [18] and blood pressure [14]. Coenzyme Q<sub>10</sub> is a powerful antioxidant that might decrease superoxide generation from endothelial cells [13, 14]. Since our patients remained dyslipidaemic, their increased plasma concentration of small-dense LDL and low HDL would have contributed to endothelial dysfunction [6]. By decreasing vascular oxidative stress, CoQ could decrease the oxidative modification of LDL and HDL in the sequestered environment of the arterial wall, thereby increasing the synthesis and/or action of endogenously derived nitric oxide [6, 37, 38]. This effect might particularly extend to triglyceride-rich lipoproteins in the post-prandial phase, when oxidative stress could be maximal in diabetes [39]. Coenzyme Q<sub>10</sub> supplementation has been reported to improve glycaemic control [14, 18, 21, 22] and blood pressure [22] in patients with diabetes. However, along with other studies [40, 41], we found no evidence to support this finding. Other potential mechanisms whereby CoQ could have improved endothelial function in the brachial artery involve reduction in the cellular levels of asymmetric dimethyl-arginine and AGEs [31, 42], as well as an increase in the bioavailability of tetrahydrobiopterin [43] and glutathione [44]. Normalization of mitochondrial superoxide production could be central to these mechanisms and to an anti-inflammatory effect of CoQ [45].

We did not rigorously study the mechanism of action of CoQ with pharmacological agents, such as acetylcholine with and without N<sup>G</sup>-monomethyl-L-arginine. Post-*ischaemic* FMD of human conduit arteries has predominately been shown to be mediated by nitric oxide, although studies to date have mostly referred to the radial artery [46]. We cannot infer that the vascular benefit of CoQ extends to microcirculatory function, where the mediators of shear-stress induced increase in blood flow could be different from conduit vessels [4]. Because NMD of the brachial artery was tested maximally with a high dose of GTN, we cannot strictly exclude that our diabetic patients had non-endothelial vascular dysfunction, and that this abnormality improved with CoQ. Acute hyperglycaemia has been shown to impair FMD of the brachial artery [47] and we did not study patients at isoglycaemia. However, at the time of ultrasound, patients were on average near-normoglycaemic and there was no difference in blood glucose concentra-

tions before and after intervention. Metformin has also been reported to improve brachial artery endothelial function in diabetes [48], but the number of patients included in our study on this agent were small and did not differ between intervention and control groups. The clinical relevance of the small but significant 1.6% increase in FMD in our patients with CoQ supplementation is not clear, given that FMD is only weakly correlated with coronary responses and that NMD did not change with treatment. We were able to detect a small improvement in FMD, however, because of the enhanced precision of our computerized method for assessing luminal diameter changes.

If our findings reflect the favourable effect of CoQ on the bioavailability and action of nitric oxide, they have implications for the prevention and reversal of atherogenesis, procoagulopathy and myocardial dysfunction in diabetic patients [2, 4, 6]. That a benefit in endothelial function was seen in the presence of diabetic dyslipidaemia without complete restoration of FMD to normal, raises the possibility of further investigating the synergistic effects of CoQ with other agents that could improve vascular function in diabetes, such as angiotensin-converting enzyme inhibitors, fish-oils and lipid regulators [4, 6].

*Acknowledgements.* This study was supported by research grants from Diabetes Australia, The National Health and Medical Research Council of Australia and The Medical Research Foundation, Royal Perth Hospital. We thank Blackmores (Australia) for providing the CoQ and matching placebo and Professor J. Best for estimating the particle size of LDL. We are also grateful to the research nurses and technical staff of the University Department of Medicine, in particular Ms M-A Powell, for her excellent assistance in carrying out the study. We acknowledge the assistance of Ms L. Rich and Mr R. Woodman in assuring the quality of the analyses of the ultrasound scans.

## References

1. Semenkovich CF, Heinecke JW (1997) The mystery of diabetes and atherosclerosis. Time for a new plot. *Diabetes* 46: 327–334
2. Rubanyi GM (1993) The role of endothelium in cardiovascular homeostasis and diseases. *J Cardiovasc Pharmacol* 22 [Suppl 4]: S1-S14
3. Cohen RA (1993) Dysfunction of vascular endothelium in diabetes mellitus. *Circulation* 87 [Suppl V]:V67–V76
4. De Vriese AS, Verbeuren TJ, Van der Voorde J, Lamiere NH, Vanhoutte PM (2000) Endothelial dysfunction in diabetes. *Br J Pharmacol* 130: 963–974
5. Giugliano D, Ceriello A, Paolisso G (1996) Oxidative stress and diabetic vascular complications. *Diabetes Care* 19: 257–267
6. Watts GF, Playford D (1998) Dyslipoproteinaemia and hyperoxidative stress in the pathogenesis of endothelial dysfunction in NIDDM: an hypothesis. *Atherosclerosis* 141: 17–31
7. Stocker R (1999) Dietary and pharmacological antioxidants in atherosclerosis. *Curr Opin Lipidol* 10: 589–597

8. Yusuf S, Dagenais G, Pogue J, Bosch J, Sleight P (2000) Vitamin E supplementation and cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Eng J Med* 342: 154–160
9. Skyrme-Jones RAP, O'Brien RC, Berry KL, Meredith IT (2000) Vitamin E supplementation improves endothelial function in type 1 diabetes mellitus: a randomized, placebo-controlled study. *J Am Coll Cardiol* 36: 94–102
10. Gazis A, White DJ, Page SR, Calcroft JR (1999) Effect of oral vitamin E ( $\alpha$ -tocopherol) supplementation in vascular endothelial function in type 2 diabetes mellitus. *Diabet Med* 16: 304–311
11. Chowienzyk PJ, Brett SE, Gopaul NK et al. (2000) Oral treatment with an antioxidant (radoxoflast) reduces oxidative stress and improves endothelial function in men with Type II diabetes. *Diabetologia* 43: 974–977
12. Ting HH, Timimi K, Boles KS, Creager SJ, Ganz P, Creager MA (1996) Vitamin C improves endothelium-dependent vasodilation in patients with non-insulin dependent diabetes mellitus. *J Clin Invest* 97: 22–28
13. Beyer RE (1990) The participation of coenzyme Q in free radical production and anti-oxidation. *Free Radic Biol Med* 8: 545–465
14. McCarty MF (1999) Coenzyme Q versus hypertension: does CoQ decrease endothelial superoxide generation? *Med Hypotheses* 53: 300–304
15. Stocker R, Bowry VW, Frei B (1991) Ubiquinol-10 protects human low-density lipoprotein more efficiently against lipid peroxidation than does  $\alpha$ -tocopherol. *Proc Natl Acad Sci USA* 88: 1646–1650
16. Thomas SR, Neuzil J, Stocker R (1996) Cosupplementation with coenzymeQ prevents the prooxidant effect of  $\alpha$ -tocopherol and increase the resistance of LDL to transition metal-dependent oxidation initiation. *Arterioscler Thromb Vasc Biol* 16: 687–696
17. DeFronzo RA, Bonadonna RC, Ferrannini E (1992) Pathogenesis of NIDDM: a balanced overview. *Diabetes Care* 15: 318–368
18. McCarty MF (1999) Can correction of sub-optimal coenzymeQ staus improve  $\beta$ -cell function in type II diabetes? *Med Hypotheses* 52: 397–400
19. Jameson S (1991) Coenzyme Q<sub>10</sub> alpha-tocopherol, and free cholesterol levels in sera from diabetic patients. In: Folkers K, Littarru G, Yamagami T (eds). *Biomedical and Clinical Aspects of Coenzyme Q*. Elsevier Science, Amsterdam pp 151–158
20. Werbach MR (1993) *Nutritional Influences on Illness 2<sup>nd</sup> edn Third Line Press, Tarzona* pp 250–251
21. Shigeta Y, Izumei K, Abe H (1966) Effect of coenzyme Q<sub>7</sub> treatment on blood sugar and ketone bodies of diabetics. *J Vitaminol (Kyoto)* 12: 293–298
22. Singh RB, Niaz MA, Rastogi SS, Shukla PK, Thakur AS (1999) Effect of hydrosoluble coenzyme Q<sub>10</sub> on blood pressures and insulin resistance in hypertensive patients with coronary artery disease. *J Hum Hypertens* 13: 203–208
23. Thomas SR, Witting PK, Stocker R (1999) A role for reduced coenzymeQ in atherosclerosis. *Biofactors* 9: 207–224
24. Woodman RA, Playford DA, Watts GF et al. (2001) Improved analysis of brachial artery ultrasound using a novel edge detection software system. *J Appl Physiol* 91: 929–937
25. Long JK, Gohil K, Packer L (1986) Simultaneous determination of tocopherols, ubiquinols and ubiquinones in blood, plasma, tissue homogenates and subcellular fractions. *Anal Biochem* 157: 106–116
26. Mori TA, Croft KD, Puddey IB, Beilin LJ (1999) An improved method for the measurement of urinary and plasma F2-isoprostanes using Gas Chromatography Mass Spectrometry. *Anal Biochem* 268: 117–125
27. Cao G, Alessio HM, Cutler RG (1993) Oxygen-radical absorbance capacity assay for antioxidants. *Free Radic Biol Med* 14: 303–311
28. Dogra G, Rich L, Stanton K, Watts GF (2001) Endothelium-dependent and independent vasodilation studied at normoglycaemia in Type I diabetes mellitus with and without microalbuminuria. *Diabetologia* 44: 593–601
29. Gopaul NK, Anngard EE, Mallet AI, Betteridge DJ, Wolfe SP, Nourouz-Zadeh J (1995) Plasma 8-epi-PGF<sub>2 $\alpha$</sub>  levels are elevated in individuals with non-insulin dependent diabetes mellitus. *FEBS Lett* 368: 225–229
30. Davi G, Ciabattini G, Consoli A et al. (1999) In vivo formation of 8-Iso-prostaglandin F<sub>2 $\alpha$</sub>  and platelet activation in diabetes mellitus: effects of improved metabolic control and vitamin E supplementation. *Circulation* 99: 224–229
31. Pieper GM (1998) Review of alterations in endothelial nitric oxide production in diabetes: protective effect of arginine on endothelial dysfunction. *Hypertension* 31: 1047–1060
32. Sundaram RK, Bhaskar A, Vijayalingam S, Viswanathan M, Mohan R, Shanmugasundaram KR (1996) Antioxidant status and lipid peroxidation in type 2 diabetes mellitus with and without complications. *Clin Sci (Colch)* 90: 255–260
33. Simons LA, von Konigsmark M, Simons J, Stocker R, Cellermaier DS Vitamin E (1999) ingestion does not improve arterial endothelial dysfunction in older adults. *Atherosclerosis* 143: 1193–1199
34. Heitzer T, Finckh B, Albers S, Krohn K, Kohlschutter A, Meinertz T (2001) Beneficial effects of  $\alpha$ -lipoic acid and ascorbic acid on endothelium-dependent, nitric oxide-mediated vasodilation in diabetic patients: relation to parameters of oxidative stress. *Free Radic Biol Med* 31: 53–61
35. Anderson TJ, Uehata A, Gerhard MD et al. (1995) Close relation of endothelial function in the human coronary and peripheral circulations. *J Am Coll Cardiol* 26: 1235–1241
36. Neunteufl T, Heher S, Katzenschlager R et al. (2000) Late prognostic value of flow-mediated dilation in the brachial artery of patients with chest pain. *Am J Cardiol* 86: 207–210
37. Liao JK, Shin WS, Lee WY, Clark SL (1995) Oxidized low-density lipoprotein decreases the expression of endothelial nitric oxide synthase. *J Biol Chem* 270: 319–324
38. McNeill KL, Fontana L, Russell-Jones DL, Rajmau I, Ritter JM, Chowienzyk PJ (2000) Inhibitory effects of low-density lipoproteins from men with type 2 diabetes on endothelium-dependent relaxation. *J Am Cardiol* 35: 1622–1627
39. Anderson RA, Evans ML, Ellis GR et al. (2001) The relationship between post-prandial lipaemia, endothelial function and oxidative stress in healthy individuals and patients with type 2 diabetes. *Atherosclerosis* 154: 475–483
40. Eriksson JG, Forsen TJ, Mortensen SA, Rooohde M (1999) The effect of coenzyme Q<sub>10</sub> administration on metabolic control in patients with type 2 diabetes mellitus. *Biofactors* 9: 315–318
41. Henriksen JE, Andersen CB, Hother-Nielsen O, Vaag A, Mortensen SA, Beck-Nielsen H. Impact of ubiquinone (coenzyme Q<sub>10</sub>) treatment on glycaemic control, insulin requirements and well-being in patients with Type 1 diabetes mellitus. *Diabetic Medicine* 199; 16: 312–318
42. Miyazaki H, Matsuoka H, Cooke JP et al. (1999) Endogenous nitric oxide synthase inhibitor: A novel marker of atherosclerosis. *Circulation* 99: 1141–1146

43. Huang A, Vita JA, Venema RC, Kearney JF (2000) Ascorbic acid enhances endothelial nitric oxide activity by increasing intracellular tetrahydrobiopterin. *J Biol Chem* 275: 17399–17406
44. Prasad A, Andrews NP, Padder FA, Hussain M, Quyyumi AA (1999) Glutathione reverses endothelial dysfunction and improves nitric oxide bioavailability. *J Am Coll Cardiol* 34: 507–514
45. Nishikawa T, Edelstein D, Du XL et al. (2000) Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature* 404: 787–790
46. Mullen MJ, Kharbanda RK, Cross J et al. (2001) Heterogeneous nature of flow-mediated dilatation in human conduit arteries in vivo. *Circ Res* 88: 145–151
47. Kawano H, Motoyama T, Hirashima O et al. (1999) Hyperglycaemia rapidly suppresses flow-mediated endothelium-dependent vasodilation of brachial artery. *J Am Coll Cardiol* 34: 146–154
48. Mather KJ, Verma S, Anderson TJ (2001) Improved endothelial function with metformin in type 2 diabetes mellitus. *J Am Coll Cardiol* 37: 1344–1350