Errata

Orosomucoid in urine predicts cardiovascular and over-all mortality in patients with Type II diabetes

M. S. Christiansen¹, E. Hommel², E. Magid¹, B. Feldt-Rasmussen³

¹ Department of Clinical Biochemistry, Amager Hospital, Copenhagen S, Denmark

² Department of Internal Medicine, Bispebjerg Hospital, Copenhagen, Denmark

³ Department of Nephrology and Endocrinology, Rigshospitalet, Copenhagen, Denmark

Diabetologia (2002) 45: 115–120

On page 118, third paragraph, line one: the first '14' should be deleted and the sentence should read: "Of patients with normal UAER ($n = 251$) 14 died compared with 27 patients with increased UAER ($n = 179$; $p < 0.001$)."

Butyrylcholinesterase K variant on chromosome 3q is associated with Type II diabetes in white Caucasian subjects

Y. Hashim¹, D. Shepherd², S. Wiltshire³, R. R. Holman⁴, J. C. Levy¹, A. Clark¹, C. A. Cull⁴

¹ Diabetes Research Laboratories, Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford, UK

² Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, UK

³ Wellcome Trust Centre for Human Genetics, University of Oxford, UK

⁴ Diabetes Trials Unit, Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Infirmary, Oxford, UK

Diabetologia (2001) 44: 2227–2230

In Table 1, the p value quoted as 0.00079 should read 0.0079 and that quoted as 0.00017 should read 0.0017. Furthermore, on page 2228, right column, line 14 and on page 2230, left column, line 11, 'means $\pm$ SEM' should read 'means SD'.