I. Shai · M. B. Schulze · J. E. Manson · M. J. Stampfer ·
N. Rifai · F. B. Hu

A prospective study of lipoprotein(a) and risk of coronary heart disease among women with type 2 diabetes

Received: 12 September 2005 / Published online: 17 November 2005
© Springer-Verlag 2005

Diabetologia (2005) 48:1469–1476

Received: 11 October 2004
Accepted: 9 March 2005
Published online: 22 June 2005

Unfortunately, the values for relative risks given in Tables 2 and 3 were incorrect. The corrected tables follow.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00427-005-0002-2>.

I. Shai · M. B. Schulze · M. J. Stampfer · F. B. Hu
Department of Nutrition, Harvard School of Public Health,
Boston, MA, USA

I. Shai (✉) · J. E. Manson · M. J. Stampfer · F. B. Hu
Department of Epidemiology, Harvard School of Public Health,
677 Huntington Avenue,
Boston, MA 02115, USA
e-mail: irish@bgu.ac.il

I. Shai
S. Daniel Abraham International Center for Health
and Nutrition, Department of Epidemiology,
Ben-Gurion University,
Beer-Sheva, Israel

J. E. Manson · M. J. Stampfer · F. B. Hu
Division of Preventive Medicine, Department of Medicine,
Brigham and Women's Hospital and Harvard Medical School,
Boston, MA, USA

J. E. Manson · M. J. Stampfer · F. B. Hu
Channing Laboratory, Department of Medicine,
Brigham and Women's Hospital and Harvard Medical School,
Boston, MA, USA

N. Rifai
Department of Laboratory Medicine, Children's Hospital
and Harvard Medical School,
Boston, MA, USA

Table 2 Relative risks (95% CIs) of CHD events across quintiles of Lp(a) among 921 women with type 2 diabetes: the Nurses' Health Study

	Quintiles of Lp(a)					<i>p</i> for trend
	1	2	3	4	5	
Median ($\mu\text{mol/l}$)	0.05	0.14	0.26	0.54	1.89	
Cases (<i>n</i>)	22	19	27	24	30	
Age-adjusted	1	0.82 (0.44–1.52)	1.23 (0.70–2.15)	1.08 (0.60–1.93)	1.37 (0.79–2.38)	0.145
MV*	1	1.00 (0.53–1.88)	1.64 (0.91–2.94)	1.16 (0.64–2.11)	1.76 (0.99–3.13)	0.058
MV+TGs	1	1.03 (0.55–1.94)	1.78 (0.98–3.24)	1.25 (0.69–2.29)	1.97 (1.10–3.55)	0.030
MV+ TGs+LDL-c+ HDL-c	1	1.02 (0.54–1.94)	1.78 (0.97–3.26)	1.15 (0.62–2.13)	1.95 (1.07–3.56)	0.035

*Multivariate model, Cox regression, adjusted for age (<49, 50–54, 55–59, 60–64, 65+ years), smoking (current, past, never), BMI (<23, 23–25, 25–28, 28–30, 30–34, 35+ kg/m²), alcohol intake (0, 0.1–4.9, 5+ g/day), physical activity (0–0.9, 1–1.9, 2–3.9, 4+ h/w), postmenopausal hormone use (premenopausal, current, past, never, missing), aspirin use (non-daily, daily), parental history of MI, history of hypertension and levels of HbA_{1c} (quintiles)

MV, multivariate

Table 3 Relative risks* (95% CIs) of CHD events across quintiles of Lp(a), stratified by selected covariates among 921 women with type 2 diabetes: the Nurses' Health Study

	Cases	Quintiles of Lp(a)					<i>p</i> for trend
		1	2	3	4	5	
Median ($\mu\text{mol/l}$)		0.05	0.14	0.26	0.54	1.89	
No aspirin use	77	1	1.74 (0.68–4.48)	2.84 (1.18–6.86)	1.85 (0.71–4.82)	3.84 (1.59–9.29)	0.006
Aspirin use (at least once/week)	45	1	0.52 (0.19–1.45)	1.01 (0.38–2.69)	1.08 (0.44–2.65)	0.63 (0.21–1.86)	0.578
							<i>p</i> for interaction=0.064
Past or never postmenopausal hormone use	84	1	0.80 (0.36–1.78)	1.44 (0.67–3.10)	1.12 (0.53–2.36)	1.94 (0.93–4.07)	0.026
Current postmenopausal hormone use	38	1	1.04 (0.31–3.50)	2.05 (0.64–6.51)	0.79 (0.22–2.90)	1.17 (0.33–4.14)	0.979
							<i>p</i> for interaction=0.105
No insulin use	86	1	0.75 (0.36–1.57)	1.55 (0.78–3.09)	1.14 (0.57–2.30)	1.54 (0.77–3.08)	0.151
Insulin use	36	1	1.53 (0.40–5.84)	1.85 (0.45–7.69)	0.93 (0.23–3.72)	1.61 (0.43–6.08)	0.662
							<i>p</i> for interaction=0.747
Alcohol abstainers	80	1	1.07 (0.47–2.41)	1.80 (0.84–3.87)	1.18 (0.53–2.62)	2.57 (1.19–5.55)	0.009
Alcohol consumers	42	1	1.17 (0.37–3.68)	1.82 (0.62–5.34)	2.02 (0.67–6.10)	1.29 (0.43–3.88)	0.978
							<i>p</i> for interaction=0.2261
Duration of diabetes <median (4.4 years)	39	1	0.53 (0.17–1.64)	0.92 (0.31–2.72)	1.06 (0.40–2.83)	1.14 (0.40–3.26)	0.448
≥median	83	1	1.56 (0.69–3.51)	2.54 (1.17–5.53)	1.29 (0.57–2.94)	2.84 (1.34–6.16)	0.029
							<i>p</i> for interaction=0.561
LDL-c<median	52	1	0.55 (0.22–1.34)	0.71 (0.28–1.79)	0.55 (0.21–1.44)	0.81 (0.33–2.01)	0.842
LDL-c≥median	70	1	1.85 (0.67–5.12)	3.62 (1.42–9.22)	2.65 (1.02–6.89)	3.87 (1.50–9.97)	0.036
							<i>p</i> for interaction=0.648
TGs<2.26 mmol/l (200mg/dl)	64	1	1.08 (0.42–2.80)	1.66 (0.67–4.12)	1.18 (0.48–2.91)	1.57 (0.66–3.71)	0.399
TGs≥2.26 mmol/l	58	1	0.67 (0.27–1.65)	1.69 (0.74–3.87)	1.03 (0.42–2.49)	2.17 (0.93–5.01)	0.030
							<i>p</i> for interaction=0.212

* Adjusted for (if not stratified by the variable) age (<49, 50–54, 55–59, 60–64, 65+ years), smoking (current, past, never), BMI (<23, 23–25, 25–28, 28–30, 30+ kg/m²), alcohol intake (less than/more than 5 g/day), physical activity (0–0.9, 1–1.9, 2–3.9, 4+ h/w), postmenopausal hormone use (premenopausal, current, past, never, missing), aspirin use (non-daily, daily), parental history of MI and history of hypertension and levels of HbA_{1c} (quintiles), TGs (quintiles), LDL-c (quintiles) and HDL-c (quintiles)