H. Elouil · A. K. Cardozo · D. L. Eizirik ·
J. C. Henquin · J. C. Jonas

High glucose and hydrogen peroxide increase c-Myc and haeme-oxygenase 1 mRNA levels in rat pancreatic islets without activating NFκB

Received: 6 October 2005 / Published online: 10 November 2005
© Springer-Verlag 2005

The online version of the original article can be found at: <http://dx.doi.org/10.1007/s00125-004-1664-4>

H. Elouil · J. C. Henquin · J. C. Jonas (✉)
Endocrinology and Metabolism,
Université Catholique de Louvain (UCL),
55.30, Avenue Hippocrate, 55,
1200 Brussels, Belgium
e-mail: jonas@endo.ucl.ac.be
Tel.: +32-2-7649575
Fax: +32-2-7645532

A. K. Cardozo · D. L. Eizirik
Laboratory of Experimental Medicine,
Université Libre de Bruxelles (ULB),
Brussels, Belgium

Diabetologia (2005) 48:496–505

The concentration of hydrogen peroxide (H₂O₂) used in this study was incorrectly given as 5 μmol/l. The actual concentration was 58 μmol/l throughout the paper (Table 3, Fig. 3b and c, Fig. 4). This does not affect the discussion or the conclusion of the paper.