Replication of the association between variants in *WFS1* and risk of type 2 diabetes in European populations

P. W. Franks · O. Rolandsson · S. L. Debenham ·
K. A. Fawcett · F. Payne · C. Dina · P. Froguel ·
K. L. Mohlke · C. Willer · T. Olsson · N. J. Wareham ·
G. Hallmans · I. Barroso · M. S. Sandhu

Published online: 15 January 2008
© Springer-Verlag 2007

Unfortunately there were some minor errors in this paper, as outlined below:

1. The reference to glucose levels in the legend of Table 2 and in the last paragraph of the Methods is redundant.
2. At the end of the Discussion the numbers of cases and controls are stated as 12,979 and 14,937 respectively, whereas they should be 14,139 and 16,109 respectively, as correctly indicated elsewhere in the paper.
3. The overall p value reported on the penultimate line of the Results is given as $p=5.4\times 10^{-11}$, whereas it should read 4.9×10^{-11} , as correctly indicated elsewhere in the paper.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-007-0887-6>.

P. W. Franks (✉) · O. Rolandsson · T. Olsson · G. Hallmans
Department of Public Health and Clinical Medicine,
Umeå University Hospital,
Umeå, Sweden
e-mail: paul.franks@medicin.umu.se
URL: <http://www.umu.se/phmed/medicin/paul.franks/>

P. W. Franks · N. J. Wareham · M. S. Sandhu
MRC Epidemiology Unit, Institute of Metabolic Science,
Cambridge, UK

S. L. Debenham · M. S. Sandhu
Department of Public Health and Primary Care,
University of Cambridge,
Cambridge, UK

K. A. Fawcett · F. Payne · I. Barroso
Metabolic Disease Group, The Wellcome Trust Sanger Institute,
Hinxton, UK

C. Dina · P. Froguel
CNRS 8090-Institute of Biology, Pasteur Institute,
59019 Lille Cedex, France

K. L. Mohlke
Department of Genetics, School of Medicine,
University of North Carolina at Chapel Hill,
Chapel Hill, NC, USA

C. Willer
Department of Biostatistics and Center for Statistical Genetics,
University of Michigan,
Ann Arbor, MI 48109, USA