Enhancement of the incretin pathway in response to bariatric surgery is important for restoration of beta cell function

M. Komatsu · T. Aizawa

Received: 25 August 2008 / Accepted: 5 November 2008 / Published online: 27 November 2008
© Springer-Verlag 2008

Keywords Bariatric surgery · Incretins · NEFA · Type 2 diabetes

Abbreviations

GLP-1 glucagon-like peptide 1
GSIS glucose-stimulated insulin secretion
ISR insulin secretion rate

To the Editor: In a recent paper, Salinari and colleagues reported normalisation of glucose metabolism associated with improved pancreatic beta cell sensitivity to nutrients derived from a mixed meal, 1 week after malabsorptive bariatric surgery (bilio-pancreatic diversion), in morbidly obese patients with type 2 diabetes [1]. This was observed before the patients had lost a significant amount of weight. On the basis of timed serum insulin, C-peptide, NEFA and plasma glucose measurements, they calculated the effects of NEFA and glucose on the insulin secretion rate (ISR). The plasma glucose level was decreased to within the normal range after the surgery, but the serum NEFA level was unchanged. Using a mathematical model, the authors

concluded that the improved ISR after bariatric surgery was due to the restoration of normal NEFA oxidation in the beta cell, which enhanced ISR despite the significant decrease in the plasma glucose level. This study is of great interest to us because we consider the malonyl-CoA pathway and the resultant NEFA augmentation of insulin exocytosis to be vital for glucose-stimulated insulin secretion (GSIS), especially for the replenishment of the readily releasable pool of insulin granules during second-phase secretion [2, 3]. Prior to the visual demonstration and molecular elucidation of intracellular granule trafficking, this process was known conceptually as ‘potentiation’ [4], ‘augmentation’ [5] or ‘amplification’ [6].

The paper makes no mention of factors other than glucose and NEFA that are also involved in meal-induced insulin secretion, such as incretins [7], parasympathetic input and amino acids, to name but a few. Of note, an increase in the serum level of the incretin glucagon-like peptide 1 (GLP-1) is documented 1 week after bilio-pancreatic diversion, and this is thought to be responsible for long-term effective weight reduction through satiety [7]. GLP-1 acts on the beta cells to increase the level of cyclic AMP, leading to replenishment of the readily releasable pool of insulin granules during GSIS. This is an ATP-sensitive K^+ channel-independent insulinotropic action [8]. Interestingly, NEFA also augments GSIS via a mechanism that does not involve this channel [5].

Salinari et al. use a mathematical model of glucose–NEFA comodulation to estimate the ISR [1]. Therefore, given the lowered basal glucose, lowered glucose excursion after the meal, unchanged NEFA and increased ISR, they will naturally conclude that the effect of NEFA has been enhanced/amplified. However, in our opinion, the increased beta cell potentiation [8], achieved through the

---

M. Komatsu (✉)
Department of Ageing Medicine and Geriatrics,
Shinshu University Graduate School of Medicine,
3-1-1 Asahi,
Matsumoto 390-8621, Japan
e-mail: mitsuk@shinshu-u.ac.jp

T. Aizawa
Centre for Medical Education, Shinshu University,
Matsumoto, Japan

GLP-1–cyclic AMP pathway after surgery, is expected in their patients, in line with previous findings in patients who underwent the same operation, at the same postoperative time-point [7]. A different conclusion might have been reached if the incretin effect had been taken into account. The inclusion of the positive effect of the incretin pathway would have at least reduced, and at most cancelled out, the calculated effect of restored β -oxidation on ISR after surgery [1].

In conclusion, by not considering factors other than glucose and NEFA in meal-induced insulin secretion, Salinari et al. [1] may have overestimated the effect of restored NEFA oxidation on ISR after bariatric surgery. The data itself are very important and have considerably expanded our understanding of the physiology of bariatric surgery. However, a more comprehensive analysis of the ISR, one that at least takes account of the incretin effect, would be desirable.

Duality of interest The authors declare that there is no duality of interest associated with this manuscript.

References

1. Salinari S, Bertuzzi A, Iaconelli A, Manco M, Mingrone G (2008) Twenty-four hour insulin secretion and beta cell NEFA oxidation in type 2 diabetic, morbidly obese patients before and after bariatric surgery. *Diabetologia* 51:1276–1284
2. Aizawa T, Komatsu M, Asanuma N, Sato Y, Sharp GWG (1998) Glucose action ‘beyond ionic events’ in the pancreatic β cell. *Trends Pharmacol Sci* 19:496–499
3. Aizawa T, Komatsu M (2005) Rab27a: a new face in beta cell metabolism-secretion coupling. *J Clin Invest* 115:227–230
4. Neshier R, Cerasi E (2002) Modeling phasic insulin release: immediate and time-dependent effects of glucose. *Diabetes* 55:S53–S59
5. Komatsu M, Sharp GWG (1998) Palmitate and myristate selectively mimic the effect of glucose in augmenting insulin release in the absence of extracellular Ca^{2+} . *Diabetes* 47:352–357
6. Henquin JC (2000) Triggering and amplifying pathways of regulation of insulin secretion by glucose. *Diabetes* 49:1751–1760
7. Guidone C, Manco M, Valera-Mora E et al (2006) Mechanisms of recovery from type 2 diabetes after malabsorptive bariatric surgery. *Diabetes* 55:2025–2031
8. Yajima H, Komatsu M, Shermerhorn T et al (1999) cAMP enhances insulin secretion by an action on the ATP-sensitive K^{+} channel-independent pathway of glucose signaling in rat pancreatic islets. *Diabetes* 48:1006–1012