Impaired glucagon-like peptide-1-induced insulin secretion in carriers of transcription factor 7-like 2 (TCF7L2) gene polymorphisms

S. A. Schäfer · O. Tschritter · F. Machicao · C. Thamer ·
N. Stefan · B. Gallwitz · J. J. Holst · J. M. Dekker ·
L. M. t'Hart · G. Nijpels · T. W. van Haften ·
H. U. Häring · A. Fritsche

Published online: 8 January 2009
© Springer-Verlag 2008

Erratum to: Diabetologia
DOI 10.1007/s00125-007-0753-6

Unfortunately the dose of the GLP-1 bolus during the hyperglycaemic clamp was incorrectly given as 0.6 pmol/kg body weight, instead of 4.5 pmol/kg body weight.

The authors apologise for this mistake.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-007-0753-6>.

S. A. Schäfer · O. Tschritter · F. Machicao · C. Thamer ·
N. Stefan · B. Gallwitz · H. U. Häring · A. Fritsche (✉)
Division of Endocrinology, Diabetology, Vascular Medicine,
Nephrology and Clinical Chemistry,
Department of Internal Medicine,
Eberhard-Karls University of Tübingen,
Otfried-Müller-Str. 10,
72076 Tübingen, Germany
e-mail: andreas.fritsche@med.uni-tuebingen.de

J. J. Holst
Department of Medical Physiology, Panum Institute,
University of Copenhagen,
Copenhagen, Denmark

J. M. Dekker · G. Nijpels
On behalf of the Hoorn study, EMGO Institute,
Vrije Universiteit Medical Center,
Amsterdam, the Netherlands

L. M. t'Hart
Department of Molecular Cell Biology,
Leiden University Medical Center,
Leiden, the Netherlands

T. W. van Haften
Department of Internal Medicine,
Utrecht University Medical Center,
Utrecht, the Netherlands