Erratum to: Expression analysis of loci associated with type 2 diabetes in human tissues

C. Cotsapas · L. Prokunina-Olsson · C. Welch ·
R. Saxena · C. Weaver · N. Usher · C. Guiducci ·
S. Bonakdar · N. Turner · B. LaCroix · J. L. Hall

Published online: 16 October 2010
© Springer-Verlag 2010

Erratum to: *Diabetologia* DOI 10.1007/s00125-010-1861-2

Unfortunately, in Fig. 2, the legend did not match the figure panels. The legend should have read:

Fig. 2 Immunohistochemical localisation of HHEX. Moderate to strong immunoreactivity was found in pancreatic acini and islets (c). Localisation was nuclear and cytoplasmic. Pancreatic ducts (b) exhibited moderate, exclusively nuclear staining. Liver parenchyma (a) and colonic epithelia (d) showed little immunoreactivity or staining

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-010-1861-2>.

C. Cotsapas · R. Saxena · C. Guiducci · S. Bonakdar
Broad Institute,
Cambridge, MA, USA

L. Prokunina-Olsson
Laboratory of Translational Genomics, Division of Cancer
Epidemiology and Genetics, National Cancer Institute, NIH,
Bethesda, MD, USA

C. Welch · C. Weaver · N. Usher · N. Turner · B. LaCroix ·
J. L. Hall (✉)
Department of Medicine, Lillehei Heart Institute,
4-280 NHH, 312 Church St SE,
Minneapolis, MN 55455, USA
e-mail: jlhall@umn.edu

C. Cotsapas · R. Saxena
Department of Medicine, Harvard Medical School,
Boston, MA, USA

J. L. Hall
Developmental Biology Center, University of Minnesota,
Minneapolis, MN, USA

C. Cotsapas · R. Saxena
Center for Human Genetic Research, Department of Medicine,
Massachusetts General Hospital,
Boston, MA, USA

R. Saxena
Department of Anesthesia, Critical Care and Pain Medicine,
Massachusetts General Hospital,
Boston, MA, USA