Erratum to: Dynamic evaluation of renal resistive index in normoalbuminuric patients with newly diagnosed hypertension or type 2 diabetes

R. M. Bruno · E. Daghini · L. Landini · D. Versari ·
A. Salvati · E. Santini · I. Di Paco · A. Magagna ·
S. Taddei · L. Ghiadoni · A. Solini

Published online: 6 July 2011
© Springer-Verlag 2011

Erratum to: Diabetologia
DOI 10.1007/s00125-011-2148-y

The first sentence of the second paragraph of the Results should have read: 'RI values were significantly higher in participants with type 2 diabetes compared with EH and control participants (0.65 ± 0.06 , 0.59 ± 0.05 and 0.58 ± 0.05 respectively), while those participants with EH showed values superimposable to controls (Fig. 2a)' to match the data presented in Fig. 2a.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-011-2148-y>.

R. M. Bruno · E. Daghini · L. Landini · D. Versari · A. Salvati ·
E. Santini · I. Di Paco · A. Magagna · S. Taddei ·
L. Ghiadoni (✉) · A. Solini
Department of Internal Medicine, University of Pisa,
Via Roma 67,
56100 Pisa, Italy
e-mail: l.ghiadoni@med.unipi.it