Increased metabolic rate and insulin sensitivity in male mice lacking the carcino-embryonic antigen-related cell adhesion molecule 2

P. R. Patel · S. K. Ramakrishnan · M. K. Kaw · C. K. Raphael · S. Ghosh ·
J. S. Marino · G. Heinrich · S. J. Lee · R. E. Bourey · J. W. Hill · D. Y. Jung ·
D. A. Morgan · J. K. Kim · S. K. Rahmouni · S. M. Najjar

Received: 22 August 2011 / Accepted: 7 November 2011 / Published online: 11 December 2011
© Springer-Verlag 2011

Abstract

Aims/hypothesis The carcino-embryonic antigen-related cell adhesion molecule (CEACAM)2 is produced in many feeding control centres in the brain, but not in peripheral insulin-targeted tissues. Global *Ceacam2* null mutation causes insulin resistance and obesity resulting from hyperphagia and hypometabolism in female *Ceacam2* homozygous null mutant mice (*Cc2* [also known as *Ceacam2*]^{−/−}) mice. Because male mice are not obese, the current study examined their metabolic phenotype.

Methods The phenotype of male *Cc2*^{−/−} mice was characterised by body fat composition, indirect calorimetry, hyperinsulinaemic–euglycaemic clamp analysis and direct recording of sympathetic nerve activity.

Results Despite hyperphagia, total fat mass was reduced, owing to the hypermetabolic state in male *Cc2*^{−/−} mice. In contrast to females, male mice also exhibited insulin sensitivity with elevated β -oxidation in skeletal muscle, which is likely to offset the effects of increased food intake. Males and females had increased brown adipogenesis. However, only males had increased activation of sympathetic tone regulation of adipose tissue and increased spontaneous activity. The mechanisms underlying sexual dimorphism in energy balance with the loss of *Ceacam2* remain unknown.

Conclusions/interpretation These studies identified a novel role for CEACAM2 in the regulation of metabolic rate and insulin sensitivity via effects on brown adipogenesis,

Electronic supplementary material The online version of this article (doi:10.1007/s00125-011-2388-x) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

P. R. Patel · S. K. Ramakrishnan · M. K. Kaw · C. K. Raphael ·
S. Ghosh · J. S. Marino · G. Heinrich · S. J. Lee · R. E. Bourey ·
J. W. Hill · S. M. Najjar (✉)
Center for Diabetes and Endocrine Research,
College of Medicine and Life Sciences, University of Toledo,
Health Science Campus, 3000 Arlington Avenue, Mail Stop 1009,
Toledo, OH 43614, USA
e-mail: sonia.najjar@utoledo.edu

P. R. Patel · S. K. Ramakrishnan · M. K. Kaw · C. K. Raphael ·
S. Ghosh · J. S. Marino · G. Heinrich · S. J. Lee · J. W. Hill ·
S. M. Najjar
Department of Physiology and Pharmacology
at the College of Medicine and Life Sciences, University of Toledo,
Health Science Campus,
Toledo, OH, USA

R. E. Bourey
Department of Internal Medicine at the College of Medicine
and Life Sciences, University of Toledo, Health Science Campus,
Toledo, OH, USA

D. Y. Jung · J. K. Kim
Department of Cellular and Molecular Physiology,
Pennsylvania State University College of Medicine,
Hershey, PA, USA

D. Y. Jung · J. K. Kim
Program in Molecular Medicine,
University of Massachusetts Medical School,
Worcester, MA, USA

D. A. Morgan · S. K. Rahmouni
Department of Internal Medicine,
University of Iowa Carver College of Medicine,
Iowa City, IA, USA

sympathetic nervous outflow to brown adipose tissue, spontaneous activity and energy expenditure in skeletal muscle.

Keywords Brown adipogenesis · CEACAM2 · Energy balance · Energy dissipation · Hypermetabolism · Hyperphagia · Insulin resistance · Insulin sensitivity · Sexual dimorphism · Sympathetic nervous activation

Abbreviations

BAT	Brown adipose tissue
<i>Cc2</i> ^{−/−}	<i>Ceacam2</i> homozygous null mutant mice
<i>Cc2</i> ^{+/+}	Wild-type littermate mice
CEACAM	Carcino-embryonic antigen-related cell adhesion molecule
FATP1	Fatty acid transport protein 1
α-IR _β	β-Subunit of the insulin receptor
MRS	¹ H-Magnetic resonance spectroscopy
SNA	Sympathetic nerve activity
UCP	Uncoupling protein
VMH	Ventromedial nucleus of the hypothalamus
$\dot{V}\text{CO}_2$	CO ₂ production
$\dot{V}\text{O}_2$	Oxygen consumption
WAT	White adipose tissue

Introduction

The carcino-embryonic antigen-related cell adhesion molecule (CEACAM)2 is highly homologous to the well characterised CEACAM1, a plasma membrane glycoprotein with ubiquitous production and multiple functions. These include cell adhesion [1], hepatic insulin clearance [2–4], angiogenesis [5, 6], tumour suppression [7–10] and anti-inflammation [11].

Except for cell adhesion, which requires an intact extracellular domain [1], all other functions ascribed to CEACAM1 are regulated by phosphorylation events on its cytoplasmic tail [11, 12]. Because CEACAM1 and CEACAM2 share 70 of 73 amino acids in the cytoplasmic tail, it is likely that these two proteins exert overlapping functions. However, in light of their different tissue-specific distribution, it is also reasonable to speculate that each performs unique functions.

Unlike CEACAM1, CEACAM2 production is limited to brain, kidney, uterus, spleen and small intestine [13–17]. Moreover, its function has not been as well elucidated. In contrast to CEACAM1, CEACAM2 does not act as a cell adhesion molecule, a fact correlated to divergent amino acid sequences on the extracellular tails of these proteins [16]. Using *Ceacam2* homozygous null mutant (*Cc2* [also known as *Ceacam2*]^{−/−}) mice, with global null deletion of *Ceacam2*, we recently identified a role for CEACAM2 in the regulation of energy balance, consistent with its production in brain centres involved in food intake behaviour [17]. Female, but

not male *Cc2*^{−/−} mice display obesity [17]. This results from hyperphagia and reduced energy expenditure. Hyperphagia causes peripheral insulin resistance and defective glucose disposal in skeletal muscle, as determined by pair-feeding analysis. Given that CEACAM2 is produced in the ventromedial nucleus of the hypothalamus (VMH) among other brain nuclei involved in the regulation of food intake, but is not produced in skeletal muscle, which is the main site of energy expenditure, the altered energy balance and glucose disposal in female *Cc2*^{−/−} mice are attributed to deranged central control [18–22].

The present studies further investigated the mechanism underlining the lean phenotype of male *Cc2*^{−/−} mice. They reveal that in contrast to that in females, hyperphagia in male mice does not cause insulin resistance or obesity, owing to increased energy expenditure that could be, at least in part, mediated by increased brown adipogenesis and activated sympathetic nervous outflow to adipose tissues.

Methods

Generation of *Cc2*^{−/−} mice The generation of the mice has been briefly described [17]. A 345 bp XbaI-SacI fragment containing exon 1 and flanking regions in the promoter and the proximal region of intron 1 was replaced with sequences derived from plasmid pGK Neo^r (electronic supplementary material [ESM] Fig. 1a). The ApaI-linearised targeting vector was electroporated into J-1 embryonic stem cells (ES-129 Sv; Genome Systems, St Louis, MO, USA). We screened 200 embryonic stem cell clones by Southern blot analysis. As expected, digestion of genomic DNA with XbaI/NsiI produced a 6.7- and 8.2-kb fragment from the wild-type and recombinant (knockout) allele, respectively (ESM Fig. 1b). Embryonic stem cells positive for homologous recombination events were injected into mouse blastocysts (C57BL/6J). Three resultant chimeric mice that gave rise to germ line transmission were bred with C57BL/6 mice, and *Cc2*^{+/−} heterozygous mice were identified by PCR, using gDNA as template and (as indicated in ESM Fig. 1a) an antisense primer from exon 2 (α-123) with the (S-107) sense primer from exon 1 to detect the wild-type sequence, or the (S-Neo^r) sense primer from the Neo^r cassette to detect the knockout sequence (not shown). F1 heterozygous mice were bred to homozygosity by brother–sister mating to produce wild-type littermate (*Cc2*^{+/+}) and *Cc2*^{−/−} homozygous mice, identified by the 950 bp and the 1.9 kb PCR products, respectively, using the primer sets described above (ESM Fig. 1c). Interestingly, the *Ceacam1* allele was intact in genomic isolates from *Cc2*^{+/+} and *Cc2*^{−/−} mice (ESM Fig. 1c).

Ceacam2 gene deletion was confirmed by RT-PCR, followed by Southern blotting, using mRNA from the liver

of wild-type, $Cc2^{+/-}$ and $Cc2^{-/-}$ mice (ESM Fig. 1d). A 50% and 100% loss of *Ceacam2* cDNA content was detected in $Cc2^{+/-}$ and $Cc2^{-/-}$ mice, respectively (ESM Fig. 1d). The specificity of the probes was examined by DNA blotting and hybridising with a $\gamma^{32}\text{P}$ -labelled ATP *Cc2*-specific probe (ESM Fig. 1d).

All animals were housed in a 12 h dark–light cycle and had free access to standard chow (Harlan Teklad 2016; Harlan, Haslett, MI, USA). All procedures were approved by the Institutional Animal Care and Utilization Committee.

Metabolic analysis Following an overnight fast, mice were anaesthetised with sodium pentobarbital at 11:00 hours. Whole venous blood was drawn from the retro-orbital sinuses to carry out radioimmunoassay analysis of serum levels of insulin and leptin (Linco Research, Billerica, MA, USA). Serum NEFA and serum triacylglycerol were determined using kits (NEFA C; Wako, Richmond, VA, USA; Triacylglycerol Kit; Pointe Scientific, Canton, MI, USA). Tissue triacylglycerol content was assayed, as described previously [23].

To determine oestrogen levels, daily vaginal smears in 50 μl normal saline were taken at 09:00 hours. The cytology was immediately examined in lavage samples under a light microscope at magnification $\times 10$ to determine the dioestrus stage of the oestrus cycle (predominantly leucocytes). At this stage, mice were anaesthetised with isoflurane and retro-orbital blood was drawn to determine serum 17-beta oestradiol levels following the manufacturer's instructions (mouse/rat oestradiol ELISA, catalogue number ES180S-100; Calbiotech, Spring Valley, CA, USA).

Body composition and hyperinsulinaemic–euglycaemic clamp Gonadal visceral white adipose tissue (WAT) was excised and weighed, and visceral adiposity expressed as percentage of total body weight. Whole-body fat and lean mass were measured by ^1H -magnetic resonance spectroscopy (MRS) (Echo Medical Systems, Houston, TX, USA), and a 2-h hyperinsulinaemic–euglycaemic clamp was performed in overnight-fasted awake mice ($n=10$ –12) with primed and continuous infusion of human regular insulin (Humulin; Eli Lilly, Indianapolis, IN, USA) at a rate of $2.5 \text{ mU kg}^{-1} \text{ min}^{-1}$, as previously described [3, 23]. Glucose metabolism was estimated with a continuous infusion of $[3\text{-}^3\text{H}]\text{glucose}$ (PerkinElmer Life and Analytical Sciences, Waltham, MA, USA) for 2 h prior to (1,850 Bq/min) and throughout the clamps (3,700 Bq/min).

Daily food intake Mice ($n \geq 5$ per age group) were housed individually in a 12 h dark–light cycle with free access to standard chow. Food was weighed daily at 12:00 hours for seven consecutive days. Daily food intake was calculated as food mass from previous day minus remaining food in the cage.

Indirect calorimetry Indirect calorimetry was performed in awake mice (4 months old; $n=6$ per group), as previously described [24]. Mice were placed in metabolism cages (TSE Systems, Bad Homburg, Germany) for 3 days and had free access to food and water. Spontaneous locomotor activity was measured with an optical beam device used for quantifying horizontal and vertical movement (x -, y - and z -axes). Oxygen consumption ($\dot{V}\text{O}_2$) and CO_2 production ($\dot{V}\text{CO}_2$) were sampled every 30 min for 5 s and data normalised per fat-free lean mass. Respiratory exchange ratio was calculated from the $\dot{V}\text{CO}_2:\dot{V}\text{O}_2$ ratio.

Recording of sympathetic nerve activity Baseline sympathetic nerve activity (SNA) was measured directly from the nerves subserving brown adipose tissue (BAT) or WAT as described previously [25, 26]. Briefly, mice were anaesthetised by intraperitoneal injection of ketamine (91 mg/kg) and xylazine (9.1 mg/kg), and a nerve fibre innervating interscapular BAT or gonadal WAT was identified, placed on the bipolar platinum–iridium electrode and secured with silicone gel (Kwik-Cast; World Precision Instruments, Sarasota, FL, USA). Each electrode was attached to a high-impedance probe (HIP-511; Grass Instruments, West Warwick, RI, USA) and the nerve signal was amplified 10^5 times with an AC pre-amplifier (P5; Grass Instruments), and filtered at 100- and 1,000-Hz cut-offs using a nerve traffic analysis system (model 706C; University of Iowa Bioengineering, Iowa City, IA, USA). The signal was then routed (1) to an oscilloscope (model 54501A; Hewlett-Packard, Palo Alto, CA, USA) to monitor the quality of the sympathetic nerve recording and (2) to a resetting voltage integrator (model B600c; University of Iowa Bioengineering). SNA measurements were made for 30 min. To exclude electrical noise in the assessment of SNA, each recording was corrected for post mortem (death induced by lethal dose of ketamine/xylazine) background BAT and WAT sympathetic activities.

Immunohistochemistry Gonadal WAT was formalin-fixed and paraffin-embedded. After deparaffinising and hydration, slides were placed in antigen retrieval solution, quenched in hydrogen peroxide solution for 30 min and blocked with normal goat serum (Vectastain Elite ABC peroxidase enzyme systems kit PK-6101; Vector Laboratories, Burlingame, CA, USA). Slides were then incubated overnight at 4°C with rabbit polyclonal α -uncoupling protein (UCP)1 antibody (1:100) (ab23841; Abcam, Cambridge, UK). After three washes with TBS (5 min each), slides were incubated at room temperature in biotinylated goat anti-rabbit IgG secondary antibody (ABC Vectastain) for 30 min, treated with ABC solution for 30 min at room temperature and stained with DAB Vectastain kit SK-4100 (Vector Laboratories).

Ex vivo palmitate oxidation Ex vivo palmitate oxidation was analysed as previously described [17]. Briefly, mice (7 months old, $n=5$) were fasted overnight and anaesthetised before soleus and gastrocnemius muscles were removed, weighed, mixed, homogenised and injected into a sealed beaker at 30°C to initiate the reaction in incubation buffer containing 0.2 mmol/l [$1\text{-}^{14}\text{C}$]palmitate at 18,500 Bq/ml in the presence of 2 mmol/l L-carnitine, 0.1 mmol/l malic acid, 2 mmol/l ATP and 0.05 mmol/l coenzyme A. After 45 min, the reaction was terminated with glacial acetic acid. Radioactive oxidised CO_2 was trapped in a well, which was suspended above the medium and filled with benzothionium hydroxide, and counted by liquid scintillation (CytoCint; MP Biomedicals, Solon, OH, USA).

Western analysis and ex vivo phosphorylation Tissues were lysed and the protein concentrations determined by BCA protein assay (Pierce Biotechnology, Rockford, IL, USA) prior to analysis by 7% or 4% to 12% gradient SDS-PAGE (Invitrogen, Carlsbad, CA, USA) and immunoprobings with specific antibodies. Antibodies included polyclonal antibodies against fatty acid transporter1 (FATP1) (I-20; Santa Cruz Biotechnology, Santa Cruz, CA, USA), AKT and phospho-AKT (Cell Signaling Technology, Danvers, MA, USA), in addition to monoclonal antibodies against actin (Sigma-Aldrich, St Louis, MO, USA) to normalise for the amount of proteins analysed. Blots were incubated with horseradish peroxidase-conjugated anti-goat IgG (Santa Cruz Biotechnology), anti-mouse IgG (Amersham Biosciences, Sunnyvale, CA, USA) or anti-rabbit IgG (Amersham) antibodies prior to detection by enhanced chemiluminescence (Amersham) and quantification by densitometry. For CEACAM2 quantification in hypothalamus, 30 µg protein of tissue lysates from overnight-fasted mice were immunoprecipitated with custom-made anti-peptide mouse CEACAM2 antibody ($\alpha\text{-mCC2}$; Bethyl Laboratories, Montgomery, TX, USA) prior to SDS-PAGE analysis and immunoblotting with the same antibody.

For ex vivo phosphorylation experiments, 50 µg tissue lysates were treated with 100 nmol/l insulin for 5 min prior to immunoprecipitation with antibodies against the β -subunit of the insulin receptor ($\alpha\text{-IR}_\beta$) (Santa Cruz Biotechnology), SDS-PAGE analysis and immunoblotting with α -phosphotyrosine antibody (Upstate Biotechnology, Billerica, MA, USA), followed by reprobing with $\alpha\text{-IR}_\beta$ to normalise against the amount of insulin receptor in the immunopellet.

Semi-quantitative real-time PCR analysis Total RNA was prepared using a kit (PerfectPure RNA Tissue kit; 5Prime, Gaithersburg, MD, USA) following the manufacturer's instructions. cDNA was synthesised with oligo dT primers and Improm II Reverse Transcriptase (Promega, Madison, WI, USA), using 1 µg total RNA and primers (ESM Table 1). cDNA was evaluated with semi-quantitative real-time PCR (Fast SYBR Green PCR mix on a StepOne Plus Real-Time PCR system; Applied Biosystems, Foster City, CA, USA). Standard curves for each primer set were generated using serial 1:4 dilutions of pooled samples. The relative amounts of mRNA were calculated by comparison with the corresponding standards and normalised relative to *Gapdh*. Results are expressed as mean $\pm$ SEM in fold change relative to controls.

Statistical methods Statistical significance was determined by two-tailed Student's *t* test unless otherwise indicated. Welch's correction was used if variances were significantly different. Statistical significance was set at 1% to 5%.

Results

Body composition of male $Cc2^{-/-}$ mice Unlike female $Cc2^{-/-}$ mice, which exhibit increased body weight, male $Cc2^{-/-}$ mice maintain a body mass similar to that of $Cc2^{+/+}$ counterparts throughout their life [17]. ^1H -MRS imaging revealed that 5-month-old male $Cc2^{-/-}$ mice had similar lean mass, but reduced fat mass compared with $Cc2^{+/+}$ mice

Table 1 Metabolic variables of male $Cc2^{+/+}$ and $Cc2^{-/-}$ mice

Variable	At 1 month, by mouse group		At 5–6 months by mouse group	
	$Cc2^{+/+}$	$Cc2^{-/-}$	$Cc2^{+/+}$	$Cc2^{-/-}$
Visceral adiposity (% BWT)	N/A	N/A	3.13 $\pm$ 0.26	2.38 $\pm$ 0.46
Lean mass (g)	N/A	N/A	23.9 $\pm$ 0.81	22.1 $\pm$ 0.82
Fat mass (g)	N/A	N/A	5.00 $\pm$ 0.30	4.01 $\pm$ 0.21 *
Serum insulin (pmol/l)	114 $\pm$ 18.7	114 $\pm$ 9.11	62.5 $\pm$ 3.43	76.0 $\pm$ 7.15
Random blood glucose (mmol/l)	7.33 $\pm$ 0.29	7.66 $\pm$ 0.33	6.49 $\pm$ 0.27	6.33 $\pm$ 0.27
Serum triacylglycerol (mmol/l)	0.54 $\pm$ 0.04	0.59 $\pm$ 0.04	0.90 $\pm$ 0.09	0.52 $\pm$ 0.10 *
Serum NEFA (mEq/l)	0.83 $\pm$ 0.03	0.67 $\pm$ 0.05	0.85 $\pm$ 0.06	0.76 $\pm$ 0.02 *
Leptin (ng/ml)	2.38 $\pm$ 0.23	2.13 $\pm$ 0.37	12.8 $\pm$ 2.28	18.9 $\pm$ 5.90

Mice were fasted overnight prior to blood drawing, except for leptin analysis (mice fasted for 6 h) and random blood glucose (blood removed at 23:00 hours)

* $p<0.05$ for $Cc2^{-/-}$ vs $Cc2^{+/+}$ mice; $n\geq 7$

BWT, body weight; N/A, not applicable

(Table 1). The decrease in fat mass did not reflect a change in body weight, perhaps owing to compensation by minor changes in body water and/or lean mass that would fall below the detection limit. Male mice also exhibited no difference in their visceral adipose mass compared with

wild-type controls (Table 1). Their leptin levels were consistently normal at all ages examined (Table 1).

Male $Cc2^{-/-}$ mice are insulin-sensitive Male $Cc2^{-/-}$ mice displayed normal serum insulin and blood glucose levels

Fig. 1 Peripheral insulin action. Overnight-fasted, awake wild-type $Cc2^{+/+}$ and $Cc2^{-/-}$ mice (5 months old; $n \geq 8$ per group) were subjected to a 2 h hyperinsulinaemic-euglycaemic clamp. **a** Whole blood glucose at basal (white $Cc2^{+/+}$, black $Cc2^{-/-}$ all panels) and clamp (striped $Cc2^{+/+}$, checked $Cc2^{-/-}$ all panels) states, and **b** at time 0 and up to 120 min of clamp (white, $Cc2^{+/+}$; black $Cc2^{-/-}$). **c** Steady-state glucose infusion rate during clamp. **d** Hepatic glucose production as above (**a**). **e–g** Glucose uptake as indicated; WAT, white adipose tissue and Gastro., gastrocnemius muscle. **h** Whole-body glucose turnover (R_d). **i** Glycolysis during clamp. **j** Whole-body glycogen synthesis. Values are all expressed as mean $\pm$ SEM in milligrams per kilogram per

minute, except for blood glucose levels (mmol/l). * $p < 0.05$ vs basal. **k** Ex vivo analysis of insulin signalling. Liver was excised from 5 months old male mice, lysed, and treated with insulin (+) or buffer alone (-) prior to immunoprecipitating (Ip) with an antibody against α -IR $_{\beta}$ and immunoblotting (Ib) with anti-phosphotyrosine (α -pTyr) antibody to assess the amount of tyrosine phosphorylated insulin receptor. Gels were reprobbed (relb) with α -IR $_{\beta}$ to normalise to the amount of insulin receptor loaded. The gel is representative of three experiments, performed on one pair of mice from each group in each experiment. The band density was scanned and analysed relative to loaded protein, and is quantified in arbitrary units (AU); * $p < 0.05$ vs untreated

Fig. 2 Biochemical analysis of lipid metabolism in skeletal muscle of male mice. Mixed skeletal muscle was isolated (gastrocnemius and soleus) from overnight-fasted 5 to 7 months old male mice (white bars, *Cc2^{+/+}*; black bars, *Cc2^{-/-}* all panels). **a** mRNA expression of genes involved in fatty acid uptake and oxidation was determined by semi-quantitative real-time PCR and normalised to *Gapdh*. **b** FATP1 abundance by western blot analysis, using α-FATP1 antibody to immunoblot (Ib) and α-actin antibody to reprobe (reIb) and normalise against the amount of protein loaded. **c** Ex vivo palmitate oxidation. **d** triacylglycerol content. Values are expressed as mean ± SEM; *n* = 4–8 per group; **p* < 0.05 vs *Cc2^{+/+}*

(Table 1). To assess insulin sensitivity in vivo, a 2 h hyperinsulinaemic–euglycaemic clamp was performed on 5-month-old mice. Basal glucose at the beginning of the clamp was similar in both groups of mice (Fig. 1a). During the

clamp, which maintains comparable glucose levels in awake overnight-fasted mice (Fig. 1b), the glucose infusion rate required to maintain euglycaemia was normal (Fig. 1c), indicating overall insulin sensitivity. Hepatic glucose production at the beginning and during the clamp was similar to that in controls (Fig. 1d), indicating normal suppression by insulin. Liver responsiveness to insulin was shown to be normal by the intact tyrosine phosphorylation (and activation) of insulin receptor β in response to insulin in liver lysates, as shown by western blot analysis of insulin receptor immunoprecipitates with α-phosphotyrosine antibody (Fig. 1k).

Glucose uptake in the whole body (Fig. 1e), including WAT (Fig. 1f) and gastrocnemius muscle (Fig. 1g), was unaltered in male *Cc2^{-/-}* mice. Consistently, whole-body glucose turnover was normal (Fig. 1h). Whole-body glycolysis (Fig. 1i) and glycogen synthesis (Fig. 1j) were also intact.

Increased fat metabolism in skeletal muscle of male *Cc2^{-/-}* mice Liver triacylglycerol content in *Cc2^{-/-}* mice was similar to that in controls (16.5 ± 1.6 μmol/g wet tissue [mean ± SEM] vs 13.5 ± 1.2 in *Cc2^{+/+}*). However, fasting serum triacylglycerol and NEFA levels were reduced significantly, starting at 2 to 3 months of age (triacylglycerol 0.48 ± 0.06 vs 0.86 ± 0.07 mmol/l in *Cc2^{+/+}*; NEFA 0.50 ± 0.07 vs 0.72 ± 0.04 mEq/l in *Cc2^{+/+}*). Skeletal muscle *Fatp1* (also known as *Slc27a1*) mRNA (Fig. 2a) and protein (Fig. 2b) levels were also increased. Together with reduced fed serum triacylglycerol (0.40 ± 0.05 vs 1.20 ± 0.08 mmol/l in *Cc2^{+/+}*; *p* < 0.05) and NEFA levels (0.40 ± 0.03 vs 0.60 ± 0.03 mEq/l in *Cc2^{+/+}*; *p* < 0.05), with no change in visceral obesity (Table 1), this points to increased fatty acid uptake in muscle relative to adipose tissue. Increased uptake in skeletal

Fig. 3 Characterisation of the metabolic state of male mice. *Cc2^{+/+}* (white bars) and *Cc2^{-/-}* (black bars) mice were individually caged, given free access to food and subjected to: **a** analysis of daily food intake over seven consecutive days (*n* > 6 per group) and **b–f** indirect calorimetry analysis (performed over a 24 h period at age 4 months, and values as indicated (**b–f**) were measured). T, total light–dark cycle; D, dark period; L, light period; *n* = 5–6 mice per group. Values are expressed as mean ± SEM; **p* < 0.05 and †*p* < 0.005 vs *Cc2^{+/+}*

muscle was followed by increased β -oxidation, as determined by $\dot{V}\text{CO}_2$ from palmitate (Fig. 2c), and as supported by increased expression of *Ppara* mRNA and its target genes, including *Cpt1* (also known as *Cpt1a*), which is involved in fatty acid transport to mitochondria (Fig. 2a). Increased β -oxidation lowered triacylglycerol content in the skeletal muscle of male *Cc2^{-/-}* mice (Fig. 2d).

Male *Cc2^{-/-}* mice display a hypermetabolic state Beginning at 2 months of age, *Cc2^{-/-}* mice displayed an increase in daily food intake, tested over a period of 7 days (Fig. 3a).

Indirect calorimetry analysis over a 24 h period (Fig. 3b–f) revealed a 7% increase in total $\dot{V}\text{O}_2$ (Fig. 3b) and $\dot{V}\text{CO}_2$ (Fig. 3c) during a dark–light cycle in 4-month-old *Cc2^{-/-}* mice compared with controls. Due to the combined increases

of both variables, the respiratory exchange ratio remained unchanged (Fig. 3d). Moreover, total spontaneous locomotor activity was increased by 31% (Fig. 3e) and heat generation by 7% (Fig. 3f). Taken together, these data show that *Cc2^{-/-}* mice are in a hypermetabolic state.

Male *Cc2^{-/-}* mice display elevated brown adipogenesis We next examined whether the hypermetabolic state and increased heat production in males is associated with increased brown adipogenesis. mRNA levels of the key BAT genes *Ucp1* and *Dio2* were elevated by three- and sixfold, respectively, in BAT derived from male *Cc2^{-/-}* mice (Fig. 4a). In females, however, mRNA levels of BAT-specific genes tended to increase by around twofold without reaching statistical significance (Fig. 4a). Levels of *Prdm16*

Fig. 4 Elevated brown adipogenesis and SNA in male *Cc2^{-/-}* mice. **a** mRNA content was assessed by semi-quantitative real-time PCR and normalised to *Gapdh* in BAT from 3-month-old *Cc2^{+/+}* (white bars) and *Cc2^{-/-}* (black bars) mice; $n \geq 6$ per group. **b** Immunohistochemical analysis of WAT from four of each mouse group at age 3 months. Tissues were fixed and immunostained with α -UCP1 antibody (brown). Representative images from three sections per mouse are shown. Magnification $\times 20$. Scale bar, 100 μm . **c, d** Baseline SNA, recorded directly from the nerve subserving **(c)** WAT or **(d)** BAT, was compared between male (M) and female (F) mouse groups as above **(a)**, but at 5 months of age; $n = 9$ –10 per group. Values **(a, c, d)** are expressed as mean $\pm$ SEM; * $p < 0.05$ vs *Cc2^{+/+}* and ** $p < 0.01$ vs *Cc2^{+/+}*

function of CEACAM2 in the sympathetic regulation of brown adipogenesis and in the oxidative state of skeletal muscle, which is a major site of energy expenditure.

Like their female counterparts [17], male *Cc2^{-/-}* mice develop hyperphagia, suggesting a primary role for CEACAM2 in the regulation of feeding behaviour. The normal circulating levels of leptin at the onset of hyperphagia in both sexes suggest that hyperphagia in *Cc2^{-/-}* mice is not secondary to changes in leptin sensitivity and/or signals [33]. Given that brain *Ceacam2* mRNA is induced by fasting and is markedly reduced upon refeeding [17], it is likely that hypothalamic CEACAM2 is involved in food intake suppression and that hyperphagia in these mice is due primarily to CEACAM2 deletion from brain centres involved in feeding behaviour and rewards, including the VMH, the globus and ventral pallidus, the striatum, the olfactory bulb and the hippocampus [17, 34, 35].

Like female mice [17], male mice exhibit increased fatty acid uptake in skeletal muscle and in the mitochondria, as suggested by increased levels of FATP1 and of *Ppara* mRNA plus its target genes. However, while in females β -oxidation of fatty acids is incomplete, and lipotoxicity develops to cause insulin resistance and reduced glucose uptake in skeletal muscle [36, 37], in male mice fatty acid oxidation in skeletal muscle is increased, leading to insulin sensitivity and lower total fat mass. Thus, complete β -oxidation in skeletal muscle, together with increased brown adipogenesis in BAT and WAT [31], and increased sympathetic tone to adipose tissue, is likely to offset the effect of hyperphagia and render male *Cc2^{-/-}* mice hypermetabolic and insulin-sensitive.

Given that CEACAM2 is not endogenously produced in adipocytes, it is likely that the changes in adipose depots of male *Cc2^{-/-}* mice are driven by a cell non-autonomous effect brought about by *Ceacam2* gene deletion. The enrichment of CEACAM2 content in metabolic sensing centres of the brain, including VMH, a nucleus associated with critical involvement in the sympathetic regulation of fatty acid mobilisation from adipose tissue [38, 39], points to loss of central CEACAM2 as a likely cause of the preferential increase in brown adipogenesis in male *Cc2^{-/-}* mice. Consistent with such a possibility, direct assessment of sympathetic nervous system activity revealed elevated sympathetic traffic to adipose tissues in male *Cc2^{-/-}* mice. This is further supported by a strong induction of key genes in BAT (*Ucp1*, *Dio2* and *Adb3r* [also known as *Adrb3*]) and WAT (*Ucp1*) of male *Cc2^{-/-}* mice, providing additional evidence that CEACAM2 participates in the central regulation of brown adipogenesis, energy dissipation and sympathetic nervous tone in adipose tissues [40–42]. The underlying mechanism remains to be determined, but the sexually dimorphic phenotype is consistent with the well-

documented sex difference with respect to obesity in rats and mice with VMH lesions, as reviewed by others [21].

Future studies using targeted deletion of *Ceacam2* are under way to dissect the mechanisms involved in the sexually dimorphic central regulation of peripheral insulin action by CEACAM proteins. Nonetheless, the current study emphasises a novel role for CEACAM2 in the hypothalamic autonomic control of brown adipogenesis and energy dissipation.

Acknowledgements The authors thank A.M. DeAngelis and T. Dai from the Najjar Laboratory at the University of Toledo College of Medicine for their assistance in genotyping, and Z. Zhang from the Kim Laboratory at Pennsylvania State University College of Medicine for his assistance in metabolic phenotyping. We also thank J. Kalisz at the Najjar Laboratory for excellent technical assistance in the generation, genotyping and maintenance of *Cc2^{-/-}* null mice, in addition to the carrying out of routine RNA analyses and the oestrogen assay. We especially thank D. Accili from Columbia University Diabetes Center for frequent scientific discussions and critical reading of the manuscript. This work was supported by grants from the National Institutes of Health as follows: R01DK054254 and R01DK083850 (to S.M. Najjar), R01-DK80756 and U24-DK59635 (to J.K. Kim), HL084207 (to K. Rahmouni) and R00 HD056491 (to J. W. Hill). Grants were also received from the United States Department of Agriculture (USDA 38903-19826, to S.M. Najjar) and the American Diabetes Association (to J.K. Kim and K. Rahmouni).

Contribution statement PRP researched data, contributed to conception and design, and to analysis and interpretation of data, and also wrote the manuscript. SKRam, MKK, CKR, SG, JSM, JWH, DYJ, DAM, GH and SJL researched and analysed data, and contributed to the drafting of the article. REB contributed to data analysis, discussion and reviewed/edited the manuscript. SKRah researched data and reviewed/edited the manuscript. JKK researched data and contributed to discussion and revision of the manuscript. SMN oversaw the work, including its conception and design, contributed to analysis and interpretation of data, and also reviewed/edited the manuscript. All authors approved the final version of the manuscript.

Duality of interest The authors declare that there is no duality of interest associated with this manuscript.

References

- Öbrink B (1997) CEA adhesion molecules: multifunctional proteins with signal-regulatory properties. *Curr Opin Cell Biol* 9:616–626
- Poy MN, Yang Y, Rezaei K et al (2002) CEACAM1 regulates insulin clearance in liver. *Nat Genet* 30:270–276
- DeAngelis AM, Heinrich G, Dai T et al (2008) Carcinoembryonic antigen-related cell adhesion molecule 1: a link between insulin and lipid metabolism. *Diabetes* 57:2296–2303
- Xu E, Dubois MJ, Leung N et al (2009) Targeted disruption of carcinoembryonic antigen-related cell adhesion molecule 1 promotes diet-induced hepatic steatosis and insulin resistance. *Endocrinology* 150:3503–3512
- Ergun S, Kilik N, Ziegeler G et al (2000) CEA-related cell adhesion molecule 1: a potent angiogenic factor and a major effector of vascular endothelial growth factor. *Mol Cell* 5:311–320

6. Horst AK, Bickert T, Brewig N et al (2009) CEACAM1+ myeloid cells control angiogenesis in inflammation. *Blood* 113:6726–6736
7. Huang J, Simpson JF, Glackin C, Riethorff L, Wagener C, Shively JE (1998) Expression of biliary glycoprotein (CD66a) in normal and malignant breast epithelial cells. *Anticancer Res* 18:3203–3212
8. Lin SH, Pu YS (1999) Function and therapeutic implication of C-CAM cell-adhesion molecule in prostate cancer. *Semin Oncol* 26:227–233
9. Laurie NA, Comegys MM, Carreiro MP et al (2005) Carcinoembryonic antigen-related cell adhesion molecule 1a-4L suppression of rat hepatocellular carcinomas. *Cancer Res* 65:11010–11017
10. Leung N, Turbide C, Balachandra B, Marcus V, Beauchemin N (2008) Intestinal tumor progression is promoted by decreased apoptosis and dysregulated Wnt signaling in *Ceacam1*^{-/-} mice. *Oncogene* 27:4943–4953
11. Gray-Owen SD, Blumberg RS (2006) CEACAM1: contact-dependent control of immunity. *Nat Rev Immunol* 6:433–446
12. Najjar SM (2002) Regulation of insulin action by CEACAM1. *Trends Endocrinol Metab* 13:240–245
13. Nedellec P, Dveksler GS, Daniels E et al (1994) Bgp2, a new member of the carcinoembryonic antigen-related gene family, encodes an alternative receptor for mouse hepatitis viruses. *J Virol* 68:4525–4537
14. Han E, Phan D, Lo P et al (2001) Differences in tissue-specific and embryonic expression of mouse *Ceacam1* and *Ceacam2* genes. *Biochem J* 355:417–423
15. Zebhauser R, Kammerer R, Eisenried A, McLellan A, Moore T, Zimmermann W (2005) Identification of a novel group of evolutionarily conserved members within the rapidly diverging murine *Cea* family. *Genomics* 86:566–580
16. Robitaille J, Izzi L, Daniels E, Zelus B, Holmes KV, Beauchemin N (1999) Comparison of expression patterns and cell adhesion properties of the mouse biliary glycoproteins Bbgp1 and Bbgp2. *Eur J Biochem* 264:534–544
17. Heinrich G, Ghosh S, Deangelis AM et al (2010) Carcinoembryonic antigen-related cell adhesion molecule 2 controls energy balance and peripheral insulin action in mice. *Gastroenterology* 139:644–652
18. Sudo M, Minokoshi Y, Shimazu T (1991) Ventromedial hypothalamic stimulation enhances peripheral glucose uptake in anesthetized rats. *Am J Physiol* 261:E298–E303
19. Haque MS, Minokoshi Y, Hamai M, Iwai M, Horiuchi M, Shimazu T (1999) Role of the sympathetic nervous system and insulin in enhancing glucose uptake in peripheral tissues after intrahypothalamic injection of leptin in rats. *Diabetes* 48:1706–1712
20. Minokoshi Y, Okano Y, Shimazu T (1994) Regulatory mechanism of the ventromedial hypothalamus in enhancing glucose uptake in skeletal muscles. *Brain Res* 649:343–347
21. King BM (2006) The rise, fall, and resurrection of the ventromedial hypothalamus in the regulation of feeding behavior and body weight. *Physiol Behav* 87:221–244
22. Shiuchi T, Haque MS, Okamoto S et al (2009) Hypothalamic orexin stimulates feeding-associated glucose utilization in skeletal muscle via sympathetic nervous system. *Cell Metab* 10:466–480
23. Park SY, Cho YR, Kim HJ et al (2006) Mechanism of glucose intolerance in mice with dominant negative mutation of CEACAM1. *Am J Physiol Endocrinol Metab* 291:E517–E524
24. Hong EG, Ko HJ, Cho YR et al (2009) Interleukin-10 prevents diet-induced insulin resistance by attenuating macrophage and cytokine response in skeletal muscle. *Diabetes* 58:2525–2535
25. Ramadori G, Fujikawa T, Fukuda M et al (2010) SIRT1 deacetylase in POMC neurons is required for homeostatic defenses against diet-induced obesity. *Cell Metab* 12:78–87
26. Zhang Z, Liu X, Morgan DA et al (2011) Neuronal receptor activity-modifying protein 1 promotes energy expenditure in mice. *Diabetes* 60:1063–1071
27. Seale P, Kajimura S, Yang W et al (2007) Transcriptional control of brown fat determination by PRDM16. *Cell Metab* 6:38–54
28. de Jesus LA, Carvalho SD, Ribeiro MO et al (2001) The type 2 iodothyronine deiodinase is essential for adaptive thermogenesis in brown adipose tissue. *J Clin Invest* 108:1379–1385
29. Onai T, Kilroy G, York DA, Bray GA (1995) Regulation of beta 3-adrenergic receptor mRNA by sympathetic nerves and glucocorticoids in BAT of Zucker obese rats. *Am J Physiol* 269:R519–R526
30. Ravussin E, Kozak LP (2009) Have we entered the brown adipose tissue renaissance? *Obes Rev* 10:265–268
31. Rosenbaum M, Leibel RL (2010) Adaptive thermogenesis in humans. *Int J Obes* 34(Suppl 1):S47–S55
32. Chernogubova E, Hutchinson DS, Nedergaard J, Bengtsson T (2005) Alpha1- and beta1-adrenoceptor signaling fully compensates for beta3-adrenoceptor deficiency in brown adipocyte norepinephrine-stimulated glucose uptake. *Endocrinology* 146:2271–2284
33. Bingham NC, Anderson KK, Reuter AL, Stallings NR, Parker KL (2008) Selective loss of leptin receptors in the ventromedial hypothalamic nucleus results in increased adiposity and a metabolic syndrome. *Endocrinology* 149:2138–2148
34. Morton GJ, Cummings DE, Baskin DG, Barsh GS, Schwartz MW (2006) Central nervous system control of food intake and body weight. *Nature* 443:289–295
35. Cota D, Barrera JG, Seeley RJ (2006) Leptin in energy balance and reward: two faces of the same coin? *Neuron* 51:678–680
36. Koves TR, Ussher JR, Noland RC et al (2008) Mitochondrial overload and incomplete fatty acid oxidation contribute to skeletal muscle insulin resistance. *Cell Metab* 7:45–56
37. Wang H, Knaub LA, Jensen DR et al (2009) Skeletal muscle-specific deletion of lipoprotein lipase enhances insulin signaling in skeletal muscle but causes insulin resistance in liver and other tissues. *Diabetes* 58:116–124
38. Bray GA, Nishizawa Y (1978) Ventromedial hypothalamus modulates fat mobilisation during fasting. *Nature* 274:900–902
39. Steffens AB, Scheurink AJ, Luiten PG, Bohus B (1988) Hypothalamic food intake regulating areas are involved in the homeostasis of blood glucose and plasma FFA levels. *Physiol Behav* 44:581–589
40. Saito M, Minokoshi Y, Shimazu T (1985) Brown adipose tissue after ventromedial hypothalamic lesions in rats. *Am J Physiol* 248:E20–E25
41. Giordano A, Frontini A, Murano I et al (2005) Regional-dependent increase of sympathetic innervation in rat white adipose tissue during prolonged fasting. *J Histochem Cytochem* 53:679–687
42. Song CK, Schwartz GJ, Bartness TJ (2009) Anterograde trans-neuronal viral tract tracing reveals central sensory circuits from white adipose tissue. *Am J Physiol Regul Integr Comp Physiol* 296:R501–R511