

# ABO and Rhesus blood groups and risk of type 2 diabetes: evidence from the large E3N cohort study

Guy Fagherazzi · Gaëlle Gusto · Françoise Clavel-Chapelon ·  
Beverley Balkau · Fabrice Bonnet

Received: 27 October 2014 / Accepted: 25 November 2014 / Published online: 23 December 2014  
© Springer-Verlag Berlin Heidelberg 2014

## Abstract

**Aims/hypothesis** The objective of this study was to evaluate the relationship of ABO blood type (A, B, AB and O), Rhesus factor (positive or negative) and a combination of the two (ABO×Rhesus) with type 2 diabetes mellitus risk.

**Methods** In total, 82,104 women from the large prospective E3N cohort were followed between 1990 and 2008. Multivariate Cox regression models were used to estimate HRs and 95% CIs.

**Results** Those with either the A (HR 1.10 [95% CI 1.02, 1.18]) or B (HR 1.21 [95% CI 1.07, 1.36]) group were at increased risk of type 2 diabetes mellitus compared with those with the O group. The association with the AB group did not reach statistical significance (HR 1.17 [95% CI 0.99, 1.39]). There

was no difference in type 2 diabetes mellitus risk between Rhesus positive and negative groups (HR 0.96 [95% CI 0.88, 1.05]). When the universal donors (O<sup>-</sup>) were taken as the reference category, we observed an increased risk for both A<sup>+</sup> (HR 1.17 [95% CI 1.02, 1.35]) and A<sup>-</sup> (HR 1.22 [95% CI 1.03, 1.45]) individuals. The greatest increase in risk was seen for those with the B<sup>+</sup> blood group (HR 1.35 [95% CI 1.13, 1.60]). We also observed a greater type 2 diabetes mellitus risk for those with the AB<sup>+</sup> group (HR 1.26 [95% CI 1.02, 1.57]). Adjustment for fasting plasma glucose and lipid concentrations in a case–control subsample did not alter the associations. **Conclusions/interpretation** This study suggests that people with the O blood type have a lower risk of developing type 2 diabetes mellitus. Therefore, blood group should be investigated in future clinical and epidemiological studies on diabetes, and further pathophysiological research is needed to determine why individuals with blood type O have a lower risk of type 2 diabetes mellitus.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00125-014-3472-9) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

G. Fagherazzi (✉) · G. Gusto · F. Clavel-Chapelon  
Team 9, Nutrition, Hormones and Women's Health, Inserm U1018,  
Centre for Research in Epidemiology and Population Health,  
Gustave Roussy Institute, 114 rue Edouard Vaillant,  
94805 Villejuif Cedex, France  
e-mail: guy.fagherazzi@gustaveroussy.fr

G. Fagherazzi · G. Gusto · F. Clavel-Chapelon · B. Balkau  
Paris-South University, Villejuif, France

B. Balkau  
Epidemiology of Diabetes, Obesity and Chronic Kidney Disease  
over the Lifecourse, Inserm U1018, Centre for Research in  
Epidemiology and Population Health, Villejuif, France

F. Bonnet  
Service d'endocrinologie diabétologie et nutrition,  
Centre Hospitalier Universitaire de Rennes, Rennes, France

F. Bonnet  
Paris-Cardiovascular Research Center (PARCC),  
Inserm UMR 970, Paris, France

**Keywords** ABO · Blood group · Cohort · Diabetes · Risk factor

## Abbreviations

E3N Etude Epidémiologique auprès des femmes  
de la Mutuelle Générale de l'Education  
Nationale  
ICAM-1 Intercellular adhesion molecule 1  
TNF-R2 TNF receptor 2

## Introduction

ABO blood type and Rhesus factors are inherited traits that have been associated with cardiovascular [1] and cancer outcomes. In particular, the AB blood type is suggested to be

associated with a high risk of stroke compared with the O blood type [2]. In the same study, an over-representation of diabetes cases was reported among individuals with the AB blood group compared with other groups.

A few studies have investigated the relationship between blood group and type 2 diabetes. Most were small cross-sectional studies of specific hospital-based populations [3, 4]. Only one study of white women of European ancestry found that blood group B was associated with a decreased risk of diabetes compared with blood group O, but this study did not evaluate type 2 diabetes risk in cross categories of ABO and Rhesus groups [5].

Therefore, we investigated the association of ABO blood group, Rhesus factor and a combination of the two (ABO×Rhesus) with type 2 diabetes risk in the large prospective E3N (Etude Epidémiologique auprès des femmes de la Mutuelle Générale de l'Education Nationale) cohort. In a nested case–control study, we also evaluated the influence of fasting plasma glucose and lipid concentration on the relationship between blood group and type 2 diabetes risk.

## Research design and methods

**Study population** The E3N study is a French prospective cohort study of 98,995 female teachers initiated in 1990. It was approved by the French institutional review board (Comité de protection des personnes d'Ile-de France Paris VII; 3 December 2008). Potential diabetes cases were identified via nine follow-up questionnaires through self-reporting of diabetes, diabetes diet, diabetes drugs and hospitalisation for diabetes, and then through the file listing drugs reimbursed by their medical insurance between January 2004 and February 2012. Type 2 diabetes cases were validated if confirmed by at least two of the following sources: self-reported diabetes in the follow-up questionnaires; ascertained diabetes in a specific diabetes questionnaire; and drugs reimbursed by health insurance.

Of the 98,995 women in the cohort, we excluded those who did not complete any questionnaires after inclusion ( $n=3,506$ ), those with prevalent diabetes ( $n=938$ ), those with prevalent cancer or cardiovascular diseases ( $n=4,509$ ) and those lacking information on ABO or Rhesus blood group ( $n=7,938$ ). Thus, a total of 82,104 women were included in the analysis, of whom 3,553 had a validated diagnosis of type 2 diabetes during follow-up (1990–2008).

**Nested case–control study** Blood samples were collected from a subcohort of the E3N study between 1995 and 1997. Concentrations of biomarkers potentially involved in mechanisms leading to diabetes were measured in 271 type 2 diabetes cases and 526 control participants matched for age, fasting status and blood collection centre.

**Statistical analysis** Cox regression models with age as the timescale were used to estimate HRs and 95% CIs in the entire cohort. Time at entry was defined as age at the beginning of follow-up and exit time was defined as the age at which participants were diagnosed with diabetes, lost to follow-up or censored at the end of the follow-up period, or died, whichever came first.

Conditional logistic regression models were used in the nested case–control study. The main exposure variables were ABO blood group (O, A, B or AB; O was used as the reference), Rhesus factor (positive or negative; positive was used as the reference) and a combination of ABO×Rhesus ( $O^+$ ,  $O^-$ ,  $A^+$ ,  $A^-$ ,  $B^+$ ,  $B^-$ ,  $AB^+$  or  $AB^-$ ;  $O^-$  was used as the reference). Both crude ( $M_0$ ) and multivariate models ( $M_1$ ) adjusted for established type 2 diabetes risk factors (see Table 1 footnote) were calculated. Crude and multivariate models gave similar results; therefore, only  $M_1$  models will be discussed.

## Results

**Baseline characteristics of the study population** There was little difference between baseline characteristics across the blood group categories (electronic supplementary material

**Table 1** HRs and 95% CIs for type 2 diabetes by ABO, Rhesus and ABO×Rhesus groups

Blood group	Cases ( $n$ )	Model $M_0$	Model $M_1$
ABO			
O	1,440	1 (reference)	1 (reference)
A	1,622	1.10 (1.03, 1.18)	1.10 (1.02, 1.18)
B	344	1.23 (1.09, 1.38)	1.21 (1.07, 1.36)
AB	147	1.19 (1.00, 1.41)	1.17 (0.99, 1.39)
Rhesus			
Positive	2,958	1 (reference)	1 (reference)
Negative	595	0.96 (0.88, 1.05)	0.96 (0.88, 1.05)
ABO×Rhesus			
$O^-$	233	1 (reference)	1 (reference)
$O^+$	1,207	1.07 (0.93, 1.23)	1.09 (0.95, 1.25)
$A^+$	1,332	1.16 (1.01, 1.33)	1.17 (1.02, 1.35)
$A^-$	290	1.19 (1.00, 1.41)	1.22 (1.03, 1.45)
$B^+$	297	1.37 (1.16, 1.63)	1.35 (1.13, 1.60)
$B^-$	47	0.97 (0.71, 1.32)	1.04 (0.76, 1.42)
$AB^+$	122	1.26 (1.01, 1.57)	1.26 (1.02, 1.57)
$AB^-$	25	1.24 (0.82, 1.87)	1.23 (0.82, 1.86)

E3N cohort data;  $n=82,104$

Data are HR (95% CI)

$M_0$ , crude model;  $M_1$ ,  $M_0$  adjusted for family history of diabetes, education level, physical activity, hypertension, smoking and BMI

[ESM] Table 1), even in the two cases that showed some significant differences.

*ABO, Rhesus and type 2 diabetes risk* Mean follow-up was 13.20 years for type 2 diabetes cases and 16.48 years for non-diabetes cases. Compared with individuals with the O blood group, those with either A or B groups were at an increased risk of type 2 diabetes (HR 1.10 [95% CI 1.02, 1.18] and HR 1.21 [95% CI 1.07, 1.36], respectively; see Table 1). The increased type 2 diabetes risk associated with the AB group did not reach statistical significance (HR 1.17 [95% CI 0.99, 1.39]). There was no association between Rhesus group and type 2 diabetes risk. When ABO and Rhesus groups were combined in the same multivariate model (data not shown), associations with diabetes risk were unchanged.

We observed an increased diabetes risk for individuals in both A<sup>+</sup> and A<sup>-</sup> groups (HR 1.17 [95% CI 1.02, 1.35] and 1.22 [95% CI 1.03, 1.45], respectively) compared with universal donors (O<sup>-</sup>). The highest risk was seen for those with the B<sup>+</sup> blood group (HR 1.35, [95% CI 1.13, 1.60]). We also observed an increased risk for those with the AB<sup>+</sup> group (HR 1.26 [95% CI 1.02, 1.57]). No difference was seen for the other categories. Trends in risk were identical when prevalent cases were included and logistical regression analyses performed.

*Influence of fasting plasma glucose and lipid concentrations* There was no association of either fasting plasma glucose or lipid concentration with ABO blood group or Rhesus factor (ESM Table 2) among the 526 control participants of the nested case–control study. Moreover, as observed in the entire cohort, we found higher risks of type 2 diabetes for individuals in the non-O blood type group, with ORs between 1.21 and 1.95, even after adjusting for metabolic covariates such as fasting plasma glucose and lipid concentrations (Table 2, models M<sub>1</sub> and M<sub>2</sub>).

## Discussion

The present study shows for the first time in a large prospective cohort that specific ABO blood groups are associated with an increased type 2 diabetes risk. Individuals with the O blood group were found to have the lowest risk of developing type 2 diabetes. When the Rhesus factor was taken into account, those with A<sup>+</sup>, A<sup>-</sup>, B<sup>+</sup> and AB<sup>+</sup> blood groups were found to be at a higher risk of developing type 2 diabetes compared with universal donors (O<sup>-</sup> group). Although the associations did not reach statistical significance, adjusting for fasting plasma glucose and lipid concentrations did not substantially alter the observed associations in the nested case–control study.

A few studies have examined the relationship between ABO blood groups and the risk of diabetes, but the results have been contradictory. Higher total cholesterol levels, glucose levels and BP have been reported in individuals with the O blood group in an Iraqi population, with a decreasing trend from those with blood group A to B, and then to AB [3]. Some of the differences with our results may be explained by the different study designs, our larger set of covariates and the different genetic backgrounds. Another cross-sectional study reported results consistent with ours [4]. A higher frequency of the B blood group was found in individuals in the type 2 diabetes group than in the general population, whereas a lower frequency of the O blood group was reported. A study of white women of European ancestry found the B blood group to be associated with a decreased risk of diabetes compared with the O blood group [5]. The difference from our results could be caused by the smaller sample size: only 1,005 women and 501 cases of diabetes were included, compared with 3,553 cases in our study [5].

In contrast to previous studies, we analysed the combined effect of ABO and Rhesus blood groups on diabetes risk. These differences highlight the importance of using a large

**Table 2** Association between ABO blood group and type 2 diabetes risk in the nested case–control study, with adjustment for several biomarkers

Blood group	Cases (n)	Model M <sub>0</sub>	Model M <sub>1</sub>	Model M <sub>2</sub>	Model M <sub>3</sub>
<b>ABO</b>					
O	123	1 (reference)	1 (reference)	1 (reference)	1 (reference)
A	13	1.21 (0.89, 1.66)	1.23 (0.87, 1.74)	1.21 (0.82, 1.78)	1.22 (0.78, 1.89)
B	25	1.33 (0.76, 2.31)	1.37 (0.75, 2.52)	1.26 (0.64, 2.48)	1.13 (0.53, 2.43)
AB	110	1.37 (0.64, 2.94)	1.49 (0.64, 3.49)	1.95 (0.75, 5.11)	2.02 (0.70, 5.87)
<b>Rhesus</b>					
Positive	233	1 (reference)	1 (reference)	1 (reference)	1 (reference)
Negative	38	0.72 (0.47, 1.10)	0.74 (0.46, 1.18)	0.78 (0.47, 1.28)	0.72 (0.41, 1.28)

n=271 cases; n=526 control participants

Data are OR (95% CI)

M<sub>0</sub>, crude model; M<sub>1</sub>, M<sub>0</sub> adjusted for fasting plasma glucose; M<sub>2</sub>, M<sub>1</sub> adjusted for total cholesterol, LDL- and HDL-cholesterol and triacylglycerol; M<sub>3</sub>, M<sub>2</sub> adjusted for family history of diabetes, education level, physical activity, hypertension, smoking and BMI

sample size with appropriate adjustment for possible confounding factors to delineate the precise effect of each ABO blood group and the Rhesus factor on diabetes risk.

The mechanisms underlying the observed association are unknown. It has been suggested that the human ABO locus might influence endothelial or inflammation markers, such as the factor VIII–von Willebrand factor (vWF) complex, which is present in higher levels in non-O individuals [6]. In addition, the ABO blood groups have been associated with plasma soluble intercellular adhesion molecule 1 (ICAM-1) and TNF receptor 2 (TNF-R2) levels [5]. These markers have both been associated with an increased type 2 diabetes risk, thus providing a potential explanation for the observed relationships [7, 8]. Finally, a recent paper suggested that the ABO blood group is one of the genetically determined host factors that modulate the composition of the intestinal microbiota [9], which participates in metabolism by affecting the energy balance, glucose metabolism and low-grade inflammation [10].

Although residual confounding cannot be completely ruled out, most of the known type 2 diabetes risk factors were controlled for. Our study population included only women but, to our knowledge, no biological mechanisms are likely to explain a sex-dependent association. Information on the participants was self-reported but this is unlikely to substantially affect the results. This is the first study to investigate a relationship between blood type and type 2 diabetes risk using such a large cohort size and a prospective design. Type 2 diabetes cases were validated; some residual misclassifications with respect to diabetes status may exist but are likely to be non-differential.

## Conclusion

Our findings support a strong relationship between blood group and diabetes risk, with participants with the O blood type having a lower risk of developing type 2 diabetes. Therefore, the effects of blood groups should be investigated in future clinical and epidemiological studies on diabetes. Further pathophysiological research is also needed to determine why the individuals with blood type O have a lower risk of type 2 diabetes.

**Acknowledgements** We are indebted to all study participants and are grateful to the E3N-EPIC group.

**Funding** The study is supported by the Mutuelle Générale de l'Éducation Nationale, the Institut de Cancérologie Gustave Roussy and the Institut National de la Santé et de la Recherche Médicale. The validation of potential diabetes cases was supported by the European Union (Integrated Project LSHM-CT-2006–037197 in Framework Programme 6 of the European Community) InterAct project. Study sponsors had no role in designing the study, data analysis or interpretation, writing the manuscript, or the decision to submit the manuscript for publication.

**Duality of interest** The authors declare that there is no duality of interest associated with this manuscript.

**Contribution statement** Author contributions were as follows: GF designed the research, conducted the research and analysed data; GF, GG, FB and BB interpreted the data; GF drafted the article, and BB, GG, FB and FC-C revised it critically; FC-C contributed substantially to data acquisition and had primary responsibility for the final content of the manuscript; all authors read and approved the final manuscript.

## References

1. He M, Wolpin B, Rexrode K et al (2012) ABO blood group and risk of coronary heart disease in two prospective cohort studies. *Arterioscler Thromb Vasc Biol* 32:2314–2320
2. Zakai NA, Judd SE, Alexander K et al (2014) ABO blood type and stroke risk: the reasons for geographic and racial differences in stroke study. *J Thromb Haemost* 12:564–570
3. Jassim WE (2012) Association of ABO blood group in Iraqis with hypercholesterolaemia, hypertension and diabetes mellitus. *East Mediterr Health J* 18:888–891
4. Qureshi MA, Bhatti R (2003) Frequency of ABO blood groups among the diabetes mellitus type 2 patients. *J Coll Phys Surg Pak* 13:453–455
5. Barbalic M, Dupuis J, Dehghan A et al (2010) Large-scale genomic studies reveal central role of ABO in sP-selectin and sICAM-1 levels. *Hum Mol Genet* 19:1863–1872
6. Gill JC, Endres-Brooks J, Bauer PJ, Marks WJ Jr, Montgomery RR (1987) The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood* 69:1691–1695
7. Thorand B, Baumert J, Chambless L et al (2006) Elevated markers of endothelial dysfunction predict type 2 diabetes mellitus in middle-aged men and women from the general population. *Arterioscler Thromb Vasc Biol* 26:398–405
8. Meigs JB, Hu FB, Rifai N, Manson JE (2004) Biomarkers of endothelial dysfunction and risk of type 2 diabetes mellitus. *JAMA* 291:1978–1986
9. Makivuokko H, Lahtinen SJ, Wacklin P et al (2012) Association between the ABO blood group and the human intestinal microbiota composition. *BMC Microbiol* 12:94
10. Cani PD, Osto M, Geurts L, Everard A (2012) Involvement of gut microbiota in the development of low-grade inflammation and type 2 diabetes associated with obesity. *Gut Microbes* 3:279–288