



## Correction to: One hour post-load plasma glucose and 3 year risk of worsening fasting and 2 hour glucose tolerance in the RISC cohort

Melania Manco<sup>1</sup> · Andrea Mari<sup>2</sup> · John Petrie<sup>3</sup> · Geltrude Mingrone<sup>4,5,6</sup> · Beverley Balkau<sup>7</sup> · for the EGIR-RISC study group

Published online: 21 March 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

**Correction to: Diabetologia**  
<https://doi.org/10.1007/s00125-018-4798-5>

The affiliation details for Geltrude Mingrone are corrected below.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00125-018-4798-5>

---

✉ Melania Manco  
melania.manco@opbg.net

✉ John Petrie  
john.petrie@glasgow.ac.uk

<sup>1</sup> Research Area for Multifactorial Diseases and Complex Phenotypes, Bambino Gesù Children's Hospital, Viale Ferdinando Baldelli 38, 00146 Rome, Italy

<sup>2</sup> CNR Institute of Neuroscience, Padua, Italy

<sup>3</sup> Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, 126 University Place, Glasgow G12 8TA, UK

<sup>4</sup> Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italia

<sup>5</sup> Università Cattolica del Sacro Cuore, Roma, Italia

<sup>6</sup> Department of Diabetes, King's College London, London, UK

<sup>7</sup> CESP Centre for Research in Epidemiology and Population Health, Univ Paris-Saclay, Univ Paris Sud, Villejuif, France