2. Van den Ouweland JMW, Lemkes HHPJ, Ruitenbeek W et al (1992) Mutation in mitochondrial tRNA Leu gene in a large pedigree with maternally transmitted type II diabetes mellitus and deafness. *Nature Genetics* 1: 369–371
3. Suzuki Y, Muramatsu T, Taniyama M et al (1996) Association of aldehyde dehydrogenase with inheritance of NIDDM. *Diabetologia* 39: 1115–1118
4. Suzuki Y, Muramatsu T, Taniyama M et al (1996) Mitochondrial aldehyde dehydrogenase in diabetes mellitus associated with mitochondrial tRNA^{Leu(UUR)} mutation at position 3243. *Diabetes Care* 19: 1423–1425
5. Taniyama M, Suzuki Y, Enomoto S et al (1995) Specific PCR amplification for detection of the mutation of mitochondrial DNA in diabetic patients. *J Japan Diab Soc* 38: 401–404
6. Suzuki Y, Goto Y, Taniyama M et al (1997) Muscle histopathology in diabetes mellitus associated with mitochondrial tRNA^{Leu(UUR)} mutation at position 3243. *J Neurol Sci* 145: 49–63
7. Odawara M, Yamashita K (1995) Are MELAS and diabetes mellitus caused solely by the same mutation at np 3243 of the mitochondrial gene? *Diabetologia* 38: 1488–1489 (Letter)

Response from the author

Dear Sir,

The pathogenic mutation in mitochondrial DNA at position 3243 exhibits enigmatic properties when its phenotypic expression is considered. Many carriers of this mutation develop mitochondrial diabetes and deafness (MIDD). In contrast, other carriers of the 3243 mutation develop the severe neurological syndrome of MELAS [1, 2]. The data presented by Suzuki and collaborators in their letter and in their previous publication [3] indicate the association of inactive ALDH2 with the 3243 mutation in patients with mitochondrial diabetes. Remarkably, this association is absent in patients with MELAS. The authors suggest that the expected accumulation of acetaldehyde in carriers of inactive ALDH2 may be a factor in developing a diabetic phenotype. In addition, the authors propose that high heteroplasmy levels of the 3243 mutation may favour the development of MELAS. Though the data are interesting a note of caution is warranted with respect to genetic association studies. They are vulnerable to misinterpretation when the study groups are not fully matched for genetic background. The presented data are based on allele frequencies of mutant ALDH2 in a group of MELAS patients and a group of diabetic patients with the 3243 mutation. For comparison, the authors have included a large control group of non-3243 carriers. Because of the intrinsic situation of relatively small numbers of 3243 carriers in each group, which are obtained through clinics, it is difficult to find groups of 3243 carriers fully matched to the control group. Besides, spurious relationships may be present in selected patients from a specific geographical region. Despite this word of caution, the results by Suzuki and collaborators are certainly of interest and will trigger studies in other populations.

An aspect on which different points of view exist is whether variations in heteroplasmy levels alone are responsible for the development of the MELAS phenotype. Studies have shown clustering of MELAS in only some pedigrees with the 3243 mutation. Other studies found no clear relation between the phenotype and heteroplasmy levels and this has led to the hypothesis proposed by Odawara and Yamashita [4] that

additional factors are required for the development of MELAS. Heteroplasmy levels are generally determined on a population of cells in a tissue sample from patients. Studies by Matthews et al. [5] have provided evidence that there is variation of heteroplasmy between individual cells which is not adequately reflected by the mean heteroplasmy value for a particular tissue. These authors suggested that the distribution of the mutation over individual cells rather than the overall level of heteroplasmy may contribute to the development of the clinical phenotype. The existence of large differences in heteroplasmy levels between individual cells has been corroborated [6] by direct visualization of mutant mtDNA in individual cells by fluorescence in situ hybridization. It is my feeling that heterogeneity in heteroplasmy of the 3243 mutation at the level of individual cells has to be taken into consideration for the evaluation of phenotype-genotype relations.

Yours sincerely,

J. A. Maassen

References

1. Goto Y, Nonaka I, Horai S (1990) A mutation in the tRNA-Leu (UUR) gene associated with the Melas subgroup of mitochondrial encephalomyopathies. *Nature* 348: 651–653
2. Maassen JA, Kadowaki T (1996) Maternally inherited diabetes and deafness: a new diabetic subtype. *Diabetologia* 39: 375–382
3. Suzuki Y, Muramatsu T, Taniyama M et al (1996) Mitochondrial aldehyde dehydrogenase in diabetes associated with mitochondrial tRNA^{Leu(UUR)} mutation at position 3243. *Diabetes Care* 19: 1423–1425
4. Odawara M, Yamashita K (1995) Are MELAS and diabetes mellitus caused solely by the same mutation at np 3243 of the mitochondrial gene? *Diabetologia* 38: 1488–1489
5. Matthews PM, Brown RM, Morten K et al (1995) Intracellular heteroplasmy for disease-associated point mutations in mtDNA: implications for disease expression and evidence for mitotic segregation of heteroplasmic units of mtDNA. *Hum Genet* 96: 261–268
6. Van de Corput MPC, Van den Ouweland JMW, Dirks RW et al (1997) Detection of mitochondrial DNA deletions in human skin fibroblasts of patients with Pearson's syndrome by two-color fluorescence in situ hybridization. *J Histochem Cytochem* 45: 55–61

Corresponding author: Dr. J. A. Maassen, Department of Medical Biochemistry, Sylvius Laboratories, Wassenaarseweg 72, NL-2330 RA Leiden, The Netherlands