Korrespondenzadresse

Dr. J.F. Funk

Klinik für Orthopädie,
Sektion Kinder- und
Neuroorthopädie, Centrum für
Muskuloskeletale Chirurgie,
Charité-Universitätsmedizin
Berlin
Charitéplatz 1, 10117 Berlin
julia.funk@charite.de

Interessenkonflikt. Die korrespondierende Autorin gibt für sich und ihre Koautoren an, dass kein Interessenkonflikt besteht.

Literatur

- Boehm S, Limpaphayom N, Alaei F et al (2008) Early results of the ponseti method for the treatment of clubfoot in distal arthrogryposis. *J Bone Joint Surg [Am]* 90:1501–1507
- Bosse HJ van, Marangoz S, Lehman WB et al (2009) Correction of arthrogryptic clubfoot with a modified Ponseti technique. *Clin Orthop Relat Res* 467(5):1283–1293
- Chung C, Nemecek R, Larsen I et al (1969) Genetic and epidemiological studies of clubfoot in Hawaii: general and medical considerations. *Hum Hered* 19:321–342
- Dobbs MB, Rudzki JR, Purcell DB et al (2004) Factors predictive of outcome after use of the Ponseti method for the treatment of idiopathic clubfeet. *J Bone Joint Surg [Am]* 86:22–27
- Gerlach DJ, Gurnett CA, Limpaphayom N et al (2009) Early results of the Ponseti method for the treatment of clubfoot associated with myelomeningocele. *J Bone Joint Surg [Am]* 91:1350–1359
- Janicki JA, Narayanan UG, Harvey B et al (2009) Treatment of neuromuscular and syndrome-associated (nonidiopathic) clubfeet using the ponseti method. *J Pediatr Orthop* 29(4):393–397
- Legaspi J, Li HY, Chow W et al (2001) Talectomy in patients with recurrent deformity in club foot. A long-term follow-up study. *J Bone Joint Surg [Br]* 83:384–387
- Menelaus MB (1971) Talectomy for equinovarus deformity in arthrogryposis and spina bifida. *J Bone Joint Surg [Br]* 53:468–473
- Mirzayan R, Early SD, Matthys GA et al (2001) Single-stage talectomy and tibio-calcaneal arthrodesis as a salvage of severe, rigid equinovarus deformity. *Foot Ankle Int* 22:209–213
- Morcuende JA, Dolan LA, Dietz FR et al (2004) Radical reduction in the rate of extensive corrective surgery for clubfoot using the Ponseti method. *Pediatrics* 113:376–380
- Morcuende JA, Dobbs MB, Frick SL (2008) Results of the Ponseti method in patients with clubfoot associated with arthrogryposis. *Iowa Orthop J* 28:22–26
- Pirani S, Outerbridge HK, Moran M et al (1995) A method of assessing the virgin clubfoot. *Pediatric Orthopaedic Society of North America. Annual Meeting, Miami*
- Ponseti IV, Smoley EN (1963) Congenital club foot: the results of treatment. *J Bone Joint Surg [Am]* 45:261–344
- Ponseti IV (1996) Congenital clubfoot: fundamentals of treatment. Oxford University Press, New York
- Siapkara A, Duncan R (2007) Congenital talipes equinovarus: a review of current management. *J Bone Joint Surg [Br]* 89:995–1000
- Thacker MM, Scher DM, Sala DA et al (2005) Use of the foot abduction orthosis following Ponseti casts: is it essential? *J Pediatr Orthop* 25:225–228
- Widmann RF, Do TT, Burke SW (2005) Radical soft-tissue release of the arthrogryptic clubfoot. *J Pediatr Orthop B* 14:111–115
- Wynne-Davies R (1964) Family studies and the cause of congenital club foot. Talipes equinovarus, talipes calcaneo-valgus and metatarsus varus. *J Bone Joint Surg [Br]* 46:445–463
- Zionts LE, Zhao G, Hitchcock K et al (2010) Has the rate of extensive surgery to treat idiopathic club-foot declined in the united states? *J Bone Joint Surg [Am]* 92:882–889

Orthopäde 2012 · 41:983
DOI 10.1007/s00132-012-2030-8
Online publiziert: 25. November 2012
© Springer-Verlag Berlin Heidelberg 2012

M. Bänfer

Springer-Verlag GmbH, Heidelberg

Erratum zu: Bibliographischen Angaben aus Der Orthopäde 05/2012

Bei folgenden Beiträgen waren die bibliographischen Angaben fehlerhaft:

- Roth A (2012) Der minimal-invasive anterolaterale Zugang. Eine Literaturübersicht. *Orthopäde* 41:377–381. DOI: 10.1007/s00132-011-1892-5
- Fink B, Mittelstaedt A (2012) Minimal-invasiver posteriorer Zugang für die Hüfttotalendoprothesenimplantation. *Orthopäde* 41:382–389. DOI: 10.1007/s00132-011-1893-4
- Nöth U, Nedopil A, Holzapfel BM et al (2012) Der minimal-invasive anteriore Zugang. *Orthopäde* 41:390–398. DOI: 10.1007/s00132-011-1894-3
- Jung J, Anagnostakos K, Kohn D (2012) Klinische Ergebnisse nach minimal-invasiver Hüftendoprothetik. *Orthopäde* 41:399–406. DOI: 10.1007/s00132-011-1895-2
- Heisel J (2012) Rehabilitation nach minimal-invasiver Hüftendoprothesenimplantation. *Orthopäde* 41:407–412. DOI: 10.1007/s00132-011-1896-1

Bitte berücksichtigen Sie die korrigierten Angaben.

Die Redaktion

Korrespondenzadresse

M. Bänfer

Springer-Verlag GmbH
Tiergartenstr. 17, 69120 Heidelberg
michal.baenfer@springer.com