Corinne Alberti
Christian Brun-Buisson
Hilmar Burchardi
Claudio Martin
Sergey Goodman
Antonio Artigas
Alberto Sicignano
Mark Palazzo
Rui Moreno
Ronan Boulmé
Eric Lepage
Jean Roger Le Gall

Epidemiology of sepsis and infection in ICU patients from an international multicentre cohort study

---

Published online: 13 March 2002
© Springer-Verlag 2002

Intensive Care Med (2002) 28:108–121

Due to an unfortunate error Table 6 appeared incorrect.
The correct table is printed below.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00134-001-1143-z>

C. Alberti (✉)

Department of Biostatistics, Saint-Louis Hospital,
U444-INSERM, 1 Avenue Claude Vellefaux,
75475 Paris Cedex 10, France
e-mail: corinne.alberti@rdb.ap-hop-paris.fr

C. Alberti · J.R. Le Gall

Intensive Care Unit, Saint-Louis Hospital, Paris 7 University,
1 Avenue Claude Vellefaux, 75475 Paris Cedex 10, France

C. Brun-Buisson

Intensive Care Unit, Henri-Mondor Hospital, Créteil, France

H. Burchardi

Intensive Care Unit, Göttingen Hospital, Göttingen, Germany

C. Martin

Critical Care–Trauma Centre, London Health Sciences Centre,
London, Canada

S. Goodman

Intensive Care Unit, Hadassah Hebrew University Medical Center,
Jerusalem, Israel

A. Artigas

Intensive Care Unit, Parc Tauli Hospital, Sabadell Barcelona, Spain

A. Sicignano

Anestesia e Rianimazione, Maggiore Hospital, Milan, Italy

M. Palazzo

Intensive Care Unit, Charing Cross Hospital, London, UK

R. Moreno

Intensive Care Unit, Santo Antonio dos Capuchos Hospital,
Lisbon, Portugal

R. Boulmé · E. Lepage

Medical Information Unit, Henri-Mondor Hospital,
Créteil, France

Table 6 Long-stay group: ICU procedures and outcomes according to infected status. ICU and hospital mortality are presented as percentage by column (95% confidence intervals). For patients with infection on ICU admission and ICU-acquired infection, ACCP criteria were evaluated in the first 24 h of the first infection

	Not infected at ICU admission		Infected at ICU admission			
	No ICU-acquired infection (n=4789)	ICU-acquired infection (n=868)	Community		Hospital	
			No ICU-acquired infection (n=1165)	ICU-acquired infection (n=339)	No infection ICU-acquired (n=818)	ICU-acquired infection (n=374)
SAPS II at ICU admission	30 (12–62)	40 (19–70)	38 (15–70)	45 (24–78)	43 (20–75)	44 (23–79)
Whole population						
ICU mortality (%; 95%CI)	12.1 (11.2–13.0)	32.1 (29.0–35.2)	22.1 (19.7–24.5)	38.4 (33.2–43.6)	35.7 (32.4–39.0)	43.9 (38.9–48.9)
ICU LOS (days; 5th–95th)	4 (3–13)	16 (5–56)	7 (3–24)	24 (7–73)	7 (3–23)	20 (7–67)
After excluding five units ^a						
ICU mortality (%; 95%CI)	11.5 (10.5–12.5)	31.9 (28.2–35.6)	22.1 (19.6–24.6)	38.9 (33.4–44.4)	35.6 (32.1–39.1)	43.6 (38.3–48.9)
ICU LOS (days; 5th–95th)	4 (3–14)	18 (6–60)	7 (3–24)	23 (7–75)	7 (3–22)	21 (7–68)
Hospital mortality (%; 95%CI)	16.9 (15.7–18.1)	39.6 (35.7–43.5)	27.8 (25.1–30.5)	46.3 (40.6–52.0)	45.7 (42.1–49.3)	53.6 (48.2–59.0)
Hospital LOS (days; 5th–95th)	14 (3–51)	30 (8–99)	16 (3–58)	30 (8–101)	15 (3–73)	30 (7–114)
According to ACCP classification						
No SIRS (%; 95%CI)						
ICU mortality		23.6 (16.1–31.1)	15.9 (10.7–21.1)	36.8 (21.5–52.2)	22.4 (14.8–30.0)	42.9 (26.5–59.3)
Hospital mortality		30.9 (22.7–39.1)	20.6 (14.9–26.4)	44.7 (28.9–60.5)	29.3 (21.0–37.6)	48.6 (32.0–65.1)
Sepsis (%; 95%CI)	–					
ICU mortality	–	17.3 (11.5–23.0)	13.2 (9.6–16.8)	32.3 (20.6–43.9)	16.2 (10.7–21.7)	37.5 (26.3–48.7)
Hospital mortality	–	26.8 (20.1–33.5)	17.0 (13.0–21.0)	38.7 (26.6–50.8)	28.3 (21.6–35.0)	50.0 (38.5–61.5)
Severe sepsis (%; 95%CI)	–					
ICU mortality	–	33.6 (25.7–41.4)	19.6 (14.2–25.1)	40.3 (28.6–52.0)	35.3 (28.1–42.5)	49.3 (37.7–60.9)
Hospital mortality	–	40.7 (32.6–48.9)	25.5 (19.5–31.5)	47.8 (35.8–59.7)	46.5 (39.0–54.0)	56.3 (44.8–67.9)
Septic shock (%; 95%CI)	–					
ICU mortality	–	57.4 (48.8–65.9)	41.9 (35.8–48.1)	38.8 (29.9–47.7)	58.3 (51.6–64.9)	46.0 (37.6–54.3)
Hospital mortality	–	62.8 (54.4–71.1)	50.0 (43.8–56.2)	45.7 (36.6–54.8)	66.8 (60.5–73.2)	55.5 (47.2–63.8)

^a No hospital outcome measures