Corsino Rey
Marta Los Arcos
Andrés Concha
Alberto Medina
Soledad Prieto
Pablo Martínez
Belen Prieto

Procalcitonin and C-reactive protein as markers of systemic inflammatory response syndrome severity in critically ill children

Published online: 26 April 2007

© Springer-Verlag 2007

The online version of the original article can be found at:
<http://dx.doi.org/10.1007/s00134-006-0509-7>.

C. Rey (✉) · A. Concha · A. Medina · S. Prieto
University of Oviedo, Department of Paediatrics, Paediatric
Intensive Care Unit, Hospital Universitario Central de Asturias,
Celestino Villamil s/n., 33006 Oviedo, Spain
e-mail: crey@uniovi.es

M. Los Arcos
Hospital Alvarez-Buylla,
Belonga s/n., 33600 Mieres, Spain

P. Martínez
University of Tamaulipas, Academic Multidisciplinary Unit of
Agronomy and Science,
Adolfo López Mateos, 87149 CD, Victoria, Tamaulipas, Mexico

B. Prieto
University of Oviedo, Biochemical Laboratory, Hospital
Universitario Central de Asturias,
Celestino Villamil s/n., 33006 Oviedo, Spain

In Fig. 4, the cut-off values of group 2 are PCT 0.89–1.64 (*not* 0.89–1.16) and the cut-off values of group 3 are PCT 1.64–9.33 (*not* 1.16–9.33). The rest of the results are correct, including number and percentages of the different groups (see Fig. 1). This error does not affect the significance of the results or the validity of the comments in the discussion. The authors regret their mistake.

Fig. 1 Cut-off values of the five procalcitonin groups and percentage of SIRS groups, according to the definitions of the American College of Chest Physicians/Society of Critical Care Medicine [6], modified for use in paediatrics [3, 7], correctly classified in each group