FOCUS EDITORIAL

Measuring physical function after ICU: one step at a time

Carol L. Hodgson^{1,2} and Linda Denehy^{3*}

© 2017 Springer-Verlag GmbH Germany and ESICM

Long-term recovery after critical illness is important to patients, families and clinicians [1]. Powerful messages from patients highlight the persistent functional legacy of an intensive care unit (ICU) stay and describe the difficulties in recovering physical strength, functional capacity, and resuming domestic roles [2, 3]. In the last 10 years, clinicians and researchers have focused on both measuring these long-term patient outcomes and intervening in an attempt to attenuate them. This research has provided us with mixed messages relating to the best methods of measuring long-term outcomes, and of identifying effective interventions and the patients who may most benefit from them. It is in this context that we review physical function outcomes through the lens of four papers published in *Intensive Care Medicine* which have focused on outcomes from 6 months to 5 years after an ICU admission and which outline the possible next steps of our inquiry.

Physical function can be mapped to the World Health Organization's International Classification of Functioning [4], with outcome measures that include organ impairment (e.g. strength), activity limitation (e.g. walking distance or activities of daily living), participation restriction (e.g. return to work or leisure activities) and measures of health status or health-related quality of life that separate physical status. Each of these domains has several specific outcome assessment tools that can be used to track patient recovery.

Across physical function domains, the relationship between organ impairment (strength), activity limitation and participation restriction following critical illness is described in several studies [5, 6]. Physical function

and activity may be reduced for months to years following critical illness compared with the healthy normal population, and poor physical function is associated with reduced health status [6, 7]. In a multicentre longitudinal study of ARDS survivors over 5 years, 86% ($n = 166$) experienced a decline in physical status from their hospital discharge functional state [6]. Patients who were stable or showed improvement over the first year after discharge commonly experienced physical decline in subsequent years which was related to age and pre-morbid co-morbidities. More than half of the cohort experienced declines in both strength and either exercise capacity or physical functioning, as measured by the Medical Research Council Manual Muscle Test (MRC), the 6-min walking distance (6MWD) and the self-reported generic health status instrument, SF-36 Physical Function domain, respectively. Similarly, in a study of ICU survivors following severe sepsis and septic shock, at 3 months there was a decline in physical function with a reduction in both muscle strength and physical activity levels (measured by accelerometers) [8].

Measuring activity limitation is particularly complex in ICU survivors. Techniques include performance-based in-person tests of activity such as the 6MWT or the timed up and go test (TUG), wearable technology such as accelerometers or wireless activity trackers that measure data such as the number of steps walked, metabolic equivalents or steps climbed [8] and patient-reported outcome measures such as the activities of daily living and physical function domain or physical component scores (PCS) of health-related quality of life tools such as the SF-36. Performance-based tests are particularly difficult since they require the patient to return to hospital or the assessor to visit the patient. This uses resources, and patient loss to follow-up can be an issue [9]. Conversely, using patient-reported measures is less difficult as these

*Correspondence: l.denehy@unimelb.edu.au

³ Department of Physiotherapy, The University of Melbourne, Melbourne, Australia

Full author information is available at the end of the article

are generally questionnaires and can be administered by phone or mail. Patient-reported outcomes are important in assessing value-based health care and are recommended by the International Consortium of Health Outcomes [10].

To date, the most common measure of physical function using patient-reported outcomes is a health-related quality of life questionnaire. Across four international studies, and at different time points, the correlations between the 6MWD, 4-m walk and the SF-36 PCS were reported to be moderately strong [11, 12]. In contrast, performance-based 6MWD and TUG explained only 54 and 33% of the variance, respectively, in the SF-36 physical function domain at 3 months after ICU in 177 patients, although correlations were also moderate between these variables [13]. Researchers should be cognisant that patient report and performance tests may measure different constructs; however, the benefits of questionnaire-based data are the minimization of both loss to follow-up and use of resources. Interestingly, in a recent Delphi process to identify a core outcome set for long-term outcomes of survivors of acute respiratory failure, there was no consensus for a measure of performance-based physical function which should be used in all survivors of acute respiratory failure [1]. The 6MWD came closest to consensus at 54% agreement, although the SF-36 was suggested as a core outcome measure [1].

In a recent publication of the impact of disability after ICU, patients who reported moderate to severe disability

at 6 months after ICU were most likely to report problems with physical function, and this was associated with a significant reduction in health-related quality of life [2]. A follow-up study of health-related quality of life in survivors 5 years after ICU discharge also demonstrated that the domains which were most reduced were physical functioning, role-physical, general health and social functioning [14]. However, in contrast to Pfoh et al., these authors reported a small change in SF-36 physical function measures in survivors compared with pre-morbid measures that may not be clinically significant at 5 years after adjustment for natural decline. Reasons for these differences may be the variability around the PCS measure, the age of the populations in the two study groups and their pre-morbid disease state. Additionally, resilience, adaptation and environmental factors (such as family support) may also have a large impact on a patient's health status [2, 14].

Early mobilization and rehabilitation is one candidate intervention to improve physical function in ICU survivors [5, 15]. In a single centre study of $n = 50$, patient-reported physical function (SF-36) improved in the ICU exercise group at 6 months compared with control, although no performance-based tests showed between-group differences. In a recent meta-analysis of active mobilization and rehabilitation during ICU, patients who were randomized to early mobilization and rehabilitation had improved muscle strength and were more likely to be able to walk independently at hospital discharge

compared to patients who received standard care [16]. The meta-analysis showed no difference in health status measured with the SF-36 domain of physical function. Other randomized trials across the last 10 years provide inconsistent evidence related to the role of rehabilitation in improving physical function outcomes, perhaps due to differences in timing of the intervention, inadequate dose or heterogeneous patient populations [17].

What are the next steps in solving the dilemma of addressing the functional deficits of ICU survivors (Fig. 1)? We have evidence that physical function is an important problem for ICU patients and their families, but conflicting evidence about how and when to intervene to improve outcomes [9]. First, we need consensus on a measure of physical function that can be used across the continuum of survivorship, which is sensitive to change and agreement on the time-point of measurement. While health status questionnaires are commonly used and SF-36 is regarded as a core outcome [1], results across studies for this outcome are most inconsistent. Second, we need to define the most effective intervention, the timing of commencement, a dose that may need to be individually titrated and the interaction of multiple interventions (e.g. sedation, nutrition, exercise). Third, we must identify the potential confounding variables such as the patient's pre-admission health trajectory, comorbidities and frailty. Finally, we need to understand genetic, biological, environmental and lifestyle factors that may assist us in developing a model of precision medicine to impact functional outcomes after critical illness, such as altered inflammatory response or mitochondrial dysfunction.

While these next steps in unravelling the answer to reduced functional outcomes in ICU survivors appear numerous and daunting, we know that, by addressing these issues, we will find a way to the final step; that is, identifying what we should do to most effectively improve our patients' functional recovery after an admission to the ICU.

Author details

¹ Australian and New Zealand Intensive Care Research Centre, Monash University, Level 3, 553 St Kilda Rd, Melbourne, Victoria 3004, Australia. ² Physiotherapy Department, The Alfred Hospital, Melbourne 3181, Australia. ³ Department of Physiotherapy, The University of Melbourne, Melbourne, Australia.

Compliance with ethical standards

Conflicts of interest

Authors declare that they have no competing interests.

Received: 13 September 2017 Accepted: 14 September 2017

Published online: 21 September 2017

References

1. Needham DM, Sepulveda KA, Dinglas VD, Chessare CM, Aronson Friedman L, Bingham Ili CO, Turnbull AE (2017) Core outcome measures for clinical research in acute respiratory failure survivors: an International Modified Delphi Consensus Study. *Am J Respir Crit Care Med*. doi:10.1164/rccm.201702-0372OC
2. Hodgson CL, Udy AA, Bailey M, Barrett J, Bellomo R, Bucknall T, Gabbe BJ, Higgins AM, Iwashyna TJ, Hunt-Smith J, Murray LJ, Myles PS, Ponsford J, Pilcher D, Walker C, Young M, Cooper DJ (2017) The impact of disability in survivors of critical illness. *Intensive Care Med* 43:992–1001
3. Misak CJ (2011) ICU-acquired weakness: obstacles and interventions for rehabilitation. *Am J Respir Crit Care Med* 183:845–846
4. World Health Organization (2001) ICF: International classification of functioning, disability and health. World Health Organization, Geneva
5. Kayambu G, Boots R, Paratz J (2015) Early physical rehabilitation in intensive care patients with sepsis syndromes: a pilot randomised controlled trial. *Intensive Care Med* 41:865–874
6. Pfoh ER, Wozniak AW, Colantuoni E, Dinglas VD, Mendez-Tellez PA, Shanholtz C, Ciesla ND, Pronovost PJ, Needham DM (2016) Physical declines occurring after hospital discharge in ARDS survivors: a 5-year longitudinal study. *Intensive Care Med* 42:1557–1566
7. Herridge MS, Tansey CM, Matte A, Tomlinson G, Diaz-Granados N, Cooper A, Guest CB, Mazer CD, Mehta S, Stewart TE, Kudlow P, Cook D, Slutsky AS, Cheung AM (2011) Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 364:1293–1304
8. Borges RC, Carvalho CR, Colombo AS, da Silva Borges MP, Soriano FG (2015) Physical activity, muscle strength, and exercise capacity 3 months after severe sepsis and septic shock. *Intensive Care Med* 41:1433–1444
9. Hodgson C, Cuthbertson BH (2016) Improving outcomes after critical illness: harder than we thought! *Intensive Care Med* 42:1772–1774
10. International Consortium of Health Outcomes. <http://www.ichom.org/>. Accessed Sept 2017
11. Chan KS, Friedman LA, Dinglas VD, Hough CL, Morris PE, Mendez-Tellez PA, Jackson JC, Ely EW, Hopkins RO, Needham DM (2016) Evaluating physical outcomes in acute respiratory distress syndrome survivors: validity, responsiveness, and minimal important difference of 4-meter gait speed test. *Crit Care Med* 44:859–868
12. Chan KS, Pfoh ER, Denehy L, Elliott D, Holland AE, Dinglas VD, Needham DM (2015) Construct validity and minimal important difference of 6-minute walk distance in survivors of acute respiratory failure. *Chest* 147:1316–1326
13. Denehy L, Nordon-Craft A, Edbrooke L, Malone D, Berney S, Schenckman M, Moss M (2014) Outcome measures report different aspects of patient function three months following critical care. *Intensive Care Med* 40:1862–1869
14. Hofhuis JG, van Stel HF, Schrijvers AJ, Rommes JH, Spronk PE (2015) ICU survivors show no decline in health-related quality of life after 5 years. *Intensive Care Med* 41:495–504
15. Hodgson C, Bellomo R, Berney S, Bailey M, Buhr H, Denehy L, Harrold M, Higgins A, Presneill J, Saxena M, Skinner E, Young P, Webb S (2015) Early mobilization and recovery in mechanically ventilated patients in the ICU: a bi-national, multi-centre, prospective cohort study. *Crit Care* 19:81
16. Tipping CJ, Harrold M, Holland A, Romero L, Nisbet T, Hodgson CL (2017) The effects of active mobilisation and rehabilitation in ICU on mortality and function: a systematic review. *Intensive Care Med* 43:171–183
17. Denehy L, Lanphere J, Needham DM (2017) Ten reasons why ICU patients should be mobilized early. *Intensive Care Med* 43:86–90