Neovascularization in Achilles tendinopathy: have we been chasing a red herring?

Johannes L. Tol · Filippo Spiezia · Nicola Maffulli

Received: 5 June 2012 / Accepted: 28 July 2012 / Published online: 14 August 2012
© Springer-Verlag 2012

Abstract The concept of neovascularization in tendinopathy seems to have gained nearly mythological proportions and quasi-religious state: it is considered of diagnostic and prognostic value, related to clinical outcome, and the exclusive target of some therapeutic interventions. However, we question whether these assumptions are based on scientific evidence, and we come to the conclusion that, in the light of recent well-performed research, it seems that detecting neovessels has no additional value for the diagnosis, no firmly confirmed prognostic value, and no proven relation with symptoms. The role of neovascularization in this field should be re-thought. *Level of evidence V.*

Keywords Neovascularization · Achilles tendons · Power Doppler · Color Doppler

‘How often have I said to you that when you have eliminated the impossible, whatever remains, *however, improbable*, must be the truth?’ said the most famous

consulting detective of the literature, Sherlock Holmes, to Doctor Watson [17]. The same logically strict deductive process should be applied to neovascularization in Achilles tendinopathy [34, 35]. Indeed, in some quarters, the concept of neovascularization in tendinopathy seems to have gained nearly mythological proportions and quasi-religious state: it is considered of diagnostic and prognostic value, related to clinical outcome, and the exclusive target of some therapeutic interventions. However, are these assumptions based on scientific evidence, or are they as vulnerable as Achilles’ heel, eventually punctured by Paris’ killing arrow?

Historical perspective

Until recently, doctors’ status was unquestioned, and patients fully trusted their opinion. At present, we have to substantiate our diagnoses with (novel) imaging modalities, even when the clinical picture is clear. In the 1990s, blood flow in symptomatic tendons on power Doppler ultrasonography (PDU) was firstly described by Newman et al. [36]. Subsequently, two Scandinavian colleagues, Ohberg and Alfredson [38], defined this blood flow as ‘neovascularization’. The blood vessels were thought to be formed anew and were described as tortuous vessels with a small lumen, predominantly located on the anterior aspect of the Achilles tendon, infiltrating it diffusely. From an etiological perspective, the neovessels were thought to be secondary to the essential abnormality of tendinopathy, the failed healing lesion [10].

Using color Doppler ultrasound (CDU), Öhberg and Alfredson showed, in a case–control study, increased blood flow and neovascularity in all painful tendons and absence of these features in the asymptomatic control tendons [2].

J. L. Tol
Sports Medicine Division, ASPETAR, Qatar Orthopaedic
and Sports Medicine Hospital, Doha, Qatar

F. Spiezia
Department of Orthopaedic and Trauma Surgery,
Campus Biomedico University, Via Alvaro del Portillo,
200, 00128 Trigoria, Rome, Italy

N. Maffulli (✉)
Barts and The London School of Medicine and Dentistry,
William Harvey Research Institute, Centre for Sports
and Exercise Medicine, Queen Mary University of London,
Mile End Hospital, 275 Bancroft Road, London E1 4DG, UK
e-mail: n.maffulli@qmul.ac.uk

They described a five grade ultrasonographic neovascularization score, which has excellent reliability [44].

Histopathology

Healthy tendons are relatively avascular [45]. Recently, neovascularization and the accompanying neoneurones have been hypothesized to be the source of pain in chronic mid-portion Achilles tendinopathy [3, 23, 27, 46, 47, 50, 53]. This hypothesis was advanced after the detection of several sensory neurotransmitters within and around the tendons using microdialysis [24] and staining of pathologic tissue specimens. Also, in the area of neovessels, there are more sympathetic than sensory nerves, with pronounced occurrence of α -adrenoreceptors, which mediate vasoconstriction. In asymptomatic tendons, there may be a physiologically slow blood flow that cannot be detected with PDU. In tendinopathic tendons, Andersson et al. [3] hypothesized that the induced vasoconstriction might lead to increased blood flow, which can be detectable by ultrasonographic Doppler techniques [19].

To the best of our knowledge, the appearance of such vasculo-neural ingrowth has only been studied once in Achilles tendinopathy, without specifying the number of observed nerves, exact relation to the neovessels, reversibility, nor their existence in asymptomatic tendons [3, 35, 43]. Biopsies from tendinopathic patellar tendons have shown increased avascular sensory nerves and decreased vascular sympathetic nerves [25].

Diagnosis

The diagnosis of Achilles tendinopathy is clinical, with pain, swelling, and impaired performance as the cardinal signs [22, 28, 30]. Symptomatic tendinopathic Achilles tendons with neovascularization show evidence of a statistically significant association between site of maximum tenderness on palpation and site of maximum presence of PDU neovessels [16]. It is, however, unknown whether the degree of maximum pain is associated with the degree of neovascularization.

In patients with a clinical diagnosis of Achilles tendinopathy, there is conflicting evidence about the presence of neovessels. In the Swedish studies, the investigators were able to detect neovessels in 100 % of the symptomatic tendons [1, 26, 37, 38]. This observation was not supported by other researchers, who reported a percentage varying from 47 to 88 % [13, 41, 42, 54].

Also, neovessels were detected in 29 % of asymptomatic athletes [44], and in 100 % of subjects after strenuous exercise [6]. Therefore, are these neovessels really newly

formed blood vessels, and does neovascularity reflect a physiological or a pathological response?

Reviewing these studies, a limitation might be the lack of standardization in assessing the degree of neovascularization and degree of prior physical activity [6]. In a cross-sectional study in active athletes, Doppler flow was not associated with pain but might reflect an adaptation to mechanical load [35].

Relation with clinical severity

Only a few studies have examined the association between neovascularization and pain or function. Reiter et al. [42] identified a statistically significant association between the presence of neovascularization and pain and restricted function in 20 patients, but de Vos et al. [13] and Sengerij et al. [44] found no relation between degree of neovascularization and pain or VISA-A [31] score in 63 tendons [13]. As in both these studies the power calculation was not based on change of neovascularization score, no definitive conclusion could be drawn.

More recently, a prospective study with over 500 coupled measurements showed no evidence of an association between neovascularization and pain domain of the VISA-A score. This study also confirmed the earlier findings of absence of correlation between the neovessels and overall VISA-A clinical outcome score [48].

Prognosis

The predictive value of neovascularization has been studied by Zanetti et al., using only a VAS and not a functional score [54]. Nevertheless, these authors reported that the presence of neovascularization at baseline had no role in predicting outcome after three and 6 months.

These findings were confirmed by de Vos et al. [13], who showed that neovascularization at baseline did not predict outcome in patients treated with the Alfredson eccentric exercises [32, 39]. In a recent larger series, it was also confirmed that a reduction from baseline neovascularization is not associated with improvement of VISA-A or decrease of VAS pain score [48].

Neovessels and therapeutic interventions

Several conservative and surgical interventions target neovessel destruction as the key to success. The most direct approach utilizes sclerosing injections of polidocanol. The original promising results have not [51], however, been confirmed in an independent large retrospective series [49].

Another injection modality that targets the area of neovascularization is high volume image guided injections (HVIGI) [11, 20, 21]. In a relatively small cohort of patients, this therapy improved the short-term functional outcome and reduced the neovascularization score, though there was no significant association between a decrease of the neuro-vascular ingrowth and functional outcome [9, 11, 21].

The potential effect of PRP on neovessels has only been studied once in Achilles tendinopathy [4, 5, 8, 12, 14, 15, 29, 40]. It is hypothesized that the formation of neovessels is related to the release of vascular endothelial growth factor (VEGF), a potent stimulator of angiogenesis [7]. VEGF is also released after injection of platelet rich plasma (PRP). An initial increase of neovascularization after PRP injection was indeed reported, but did not differ from what occurred after placebo injection of saline, suggesting a needle instead of content effect [14].

The Achilles heel

The introduction and popularization of the neovascularization theory has given an enormous stimulus to Achilles tendinopathy research. With the increase in number and quality of studies, the Achilles heel of this promising theory has become apparent. In the light of recent well-performed research, it seems that detecting neovessels has no additional value for the diagnosis, no firmly confirmed prognostic value, and no proven relation with symptoms [33]. Also, all these issues lack of standardization of machine settings before expanding on the role of the neovascularization itself in the process of tendinopathy [52]. Hence, as Holmes said before, it should be clear that ‘It is a capital mistake to theorize before one has data. Insensibly one begins to twist facts to suit theories, instead of theories to suit facts’ [18].

References

1. Alfredson H, Ohberg L (2005) Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc* 13(4):338–344
2. Alfredson H, Ohberg L, Forsgren S (2003) Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 11(5):334–338
3. Andersson G, Danielson P, Alfredson H, Forsgren S (2007) Nerve-related characteristics of ventral paratendinous tissue in chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 15(10):1272–1279
4. Andia I, Sanchez M, Maffulli N (2011) Platelet rich plasma therapies for sports muscle injuries: any evidence behind clinical practice? *Expert Opin Biol Ther* 11(4):509–518
5. Andia I, Sanchez M, Maffulli N (2010) Tendon healing and platelet-rich plasma therapies. *Expert Opin Biol Ther* 10(10):1415–1426
6. Boesen MI, Boesen A, Koenig MJ, Bliddal H, Torp-Pedersen S (2006) Ultrasonographic investigation of the Achilles tendon in elite badminton players using color Doppler. *Am J Sports Med* 34(12):2013–2021
7. Bosch G, van Schie HT, de Groot MW, Cadby JA, van de Lest CH, Barneveld A, van Weeren PR (2010) Effects of platelet-rich plasma on the quality of repair of mechanically induced core lesions in equine superficial digital flexor tendons: a placebo-controlled experimental study. *J Orthop Res* 28(2):211–217
8. Castricini R, Longo UG, De Benedetto M, Panfoli N, Pirani P, Zini R, Maffulli N, Denaro V (2011) Platelet-rich plasma augmentation for arthroscopic rotator cuff repair: a randomized controlled trial. *Am J Sports Med* 39(2):258–265
9. Chan O, O'Dowd D, Padhiar N, Morrissey D, King J, Jalan R, Maffulli N, Crisp T (2008) High volume image guided injections in chronic Achilles tendinopathy. *Disabil Rehabil* 30(20–22):1697–1708
10. Cook JL, Khan KM, Purdam C (2002) Achilles tendinopathy. *Man Ther* 7(3):121–130
11. Crisp T, Khan F, Padhiar N, Morrissey D, King J, Jalan R, Maffulli N, Frer OC (2008) High volume ultrasound guided injections at the interface between the patellar tendon and Hoffa's body are effective in chronic patellar tendinopathy: a pilot study. *Disabil Rehabil* 30(20–22):1625–1634
12. de Jonge S, de Vos RJ, Weir A, van Schie HT, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Tol JL (2011) One-year follow-up of platelet-rich plasma treatment in chronic Achilles tendinopathy: a double-blind randomized placebo-controlled trial. *Am J Sports Med* 39(8):1623–1629
13. de Vos RJ, Weir A, Cobben LP, Tol JL (2007) The value of power Doppler ultrasonography in Achilles tendinopathy: a prospective study. *Am Journal Sports Med* 35(10):1696–1701
14. de Vos RJ, Weir A, van Schie HT, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Tol JL (2010) Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 303(2):144–149
15. Del Buono A, Papalia R, Denaro V, Maccauro G, Maffulli N (2011) Platelet rich plasma and tendinopathy: state of the art. *Int J Immunopathol Pharmacol* 24(1 Suppl 2):79–83
16. Divani K, Chan O, Padhiar N, Twycross-Lewis R, Maffulli N, Crisp T, Morrissey D (2010) Site of maximum neovascularisation correlates with the site of pain in recalcitrant mid-tendon Achilles tendinopathy. *Man Ther* 15(5):463–468
17. Doyle A (2001) *The sign of four*. Penguin Classics, London, pp 22–27
18. Doyle A (2007) *The adventures of Sherlock Holmes*. Penguin Classics, Boston
19. Genovese E, Ronga M, Recaldini C, Fontana F, Callegari L, Maffulli N, Fugazzola C (2011) Analysis of achilles tendon vascularity with second-generation contrast-enhanced ultrasound. *J Clin Ultrasound* 39(3):141–145
20. Hamilton B, Remedios D, Loosemore M, Maffulli N (2008) Achilles tendon rupture in an elite athlete following multiple injection therapies. *J Sci Med Sport* 11(6):566–568
21. Humphrey J, Chan O, Crisp T, Padhiar N, Morrissey D, Twycross-Lewis R, King J, Maffulli N (2010) The short-term effects of high volume image guided injections in resistant non-insertional Achilles tendinopathy. *J Sci Med Sport* 13(3):295–298
22. Jhingan SPM, O'Driscoll G, Lewin C, Teatino R, Malliaras P, Maffulli N, Morrissey D (2011) Thicker Achilles tendons are a

- risk factor to develop Achilles tendinopathy in elite professional soccer players. *MLTJ* 1(2):51–56
23. Kader D, Saxena A, Movin T, Maffulli N (2002) Achilles tendinopathy: some aspects of basic science and clinical management. *Br J Sports Med* 36(4):239–249
 24. Knobloch K (2008) The role of tendon microcirculation in Achilles and patellar tendinopathy. *J Orthop Surg Res* 3:18
 25. Lian O, Dahl J, Ackermann PW, Frihagen F, Engebretsen L, Bahr R (2006) Pronociceptive and antinociceptive neuromediators in patellar tendinopathy. *Am J Sports Med* 34(11):1801–1808
 26. Lind B, Ohberg L, Alfredson H (2006) Sclerosing polidocanol injections in mid-portion Achilles tendinosis: remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 14(12):1327–1332
 27. Longo UG, Ronga M, Maffulli N (2009) Achilles tendinopathy. *Sports Med Arthrosc* 17(2):112–126
 28. Maffulli N (1996) Clinical tests in sports medicine: more on Achilles tendon. *Br J Sports Med* 30(3):250
 29. Maffulli N, Del Buono A (2012) Platelet plasma rich products in musculoskeletal medicine: any evidence? *Surgeon* 10(3):148–150
 30. Maffulli N, Kenward MG, Testa V, Capasso G, Regine R, King JB (2003) Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clin J Sport Med* 13(1):11–15
 31. Maffulli N, Longo UG, Testa V, Oliva F, Capasso G, Denaro V (2008) Italian translation of the VISA-A score for tendinopathy of the main body of the Achilles tendon. *Disabil Rehabil* 30(20–22):1635–1639
 32. Maffulli N, Walley G, Sayana MK, Longo UG, Denaro V (2008) Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 30(20–22):1677–1684
 33. Malliaras P, Chan O, Simran G, Martinez de Albornoz P, Morrissey D, Maffulli N (2012) Doppler ultrasound signal in achilles tendinopathy reduces immediately after activity. *Int J Sports Med* 33(6):480–484
 34. Malliaras P, Purdam C, Maffulli N, Cook JL (2010) Temporal sequence of gray-scale ultrasound changes and their relationship with neovascularity and pain in the patellar tendon. *Br J Sports Med* 44(13):944–947
 35. Malliaras P, Richards PJ, Garau G, Maffulli N (2008) Achilles tendon Doppler flow may be associated with mechanical loading among active athletes. *Am J Sports Med* 36(11):2210–2215
 36. Newman JS, Adler RS, Bude RO, Rubin JM (1994) Detection of soft-tissue hyperemia: value of power Doppler sonography. *AJR Am J Roentgenol* 163(2):385–389
 37. Ohberg L, Alfredson H (2002) Ultrasound guided sclerosis of neovessels in painful chronic Achilles tendinosis: pilot study of a new treatment. *Br J Sports Med* 36(3):173–175
 38. Ohberg L, Lorentzon R, Alfredson H (2001) Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: an ultrasonographic investigation. *Knee Surg Sports Traumatol Arthrosc* 9(4):233–238
 39. Ohberg L, Lorentzon R, Alfredson H (2004) Eccentric training in patients with chronic Achilles tendinosis: normalised tendon structure and decreased thickness at follow up. *Br J Sports Med* 38(1):8–11
 40. Parafioriti A, Del Bianco S, Tibalt E, Oliva F, Berardi AC (2011) Single injection of platelet-rich plasma in a rat Achilles tendon tear model. *MLTJ* 1(2):41–47
 41. Peers KH, Brys PP, Lysens RJ (2003) Correlation between power Doppler ultrasonography and clinical severity in Achilles tendinopathy. *Int Orthop* 27(3):180–183
 42. Reiter M, Ulreich N, Dirisamer A, Tscholakoff D, Bucek RA (2004) Colour and power Doppler sonography in symptomatic Achilles tendon disease. *Int J Sports Med* 25(4):301–305
 43. Richards PJ, McCall IW, Day C, Belcher J, Maffulli N (2010) Longitudinal microvascularity in Achilles tendinopathy (power Doppler ultrasound, magnetic resonance imaging time-intensity curves and the Victorian Institute of Sport Assessment-Achilles questionnaire): a pilot study. *Skeletal Radiol* 39(6):509–521
 44. Sengkerij PM, de Vos RJ, Weir A, van Weelde BJ, Tol JL (2009) Interobserver reliability of neovascularization score using power Doppler ultrasonography in midportion achilles tendinopathy. *Am J Sports Med* 37(8):1627–1631
 45. Sharma P, Maffulli N (2005) Basic biology of tendon injury and healing. *Surgeon* 3(5):309–316
 46. Sharma P, Maffulli N (2006) Understanding and managing Achilles tendinopathy. *Br J Hosp Med (Lond)* 67(2):64–67
 47. Sharma P, Maffulli N (2008) Tendinopathy and tendon injury: the future. *Disabil Rehabil* 30(20–22):1733–1745
 48. Tol J, de Jonge S, Weir A, de Vos RJ, Verhaar J (2012) Relationship between neovascularization and clinical severity in Achilles tendinopathy: a prospective analysis of 556 paired measurements. *Knee Surg Sports Traumatol Arthrosc* 20(suppl 1):S63
 49. van Sterkenburg MN, de Jonge MC, Siersevelt IN, van Dijk CN (2010) Less promising results with sclerosing ethoxysclerol injections for midportion achilles tendinopathy: a retrospective study. *Am J Sports Med* 38(11):2226–2232
 50. Vora AM, Myerson MS, Oliva F, Maffulli N (2005) Tendinopathy of the main body of the Achilles tendon. *Foot Ankle Clin* 10(2):293–308
 51. Willberg L, Sunding K, Forssblad M, Alfredson H (2007) Ultrasound- and Doppler-guided arthroscopic shaving to treat jumper's knee: a technical note. *Knee Surg Sports Traumatol Arthrosc* 15(11):1400–1403
 52. Yang X, Pugh ND, Coleman DP, Nokes LD (2010) Are Doppler studies a useful method of assessing neovascularization in human Achilles tendinopathy? A systematic review and suggestions for optimizing machine settings. *J Med Eng Technol* 35(7–8):365–372
 53. Zafar MS, Mahmood A, Maffulli N (2009) Basic science and clinical aspects of achilles tendinopathy. *Sports Med Arthrosc* 17(3):190–197
 54. Zanetti M, Metzendorf A, Kundert HP, Zollinger H, Vienne P, Seifert B, Hodler J (2003) Achilles tendons: clinical relevance of neovascularization diagnosed with power Doppler US. *Radiology* 227(2):556–560