ERRATUM

Erratum to: Two doses of low-dose perioperative dexamethasone improve the clinical outcome after total knee arthroplasty: a randomized controlled study

Bin Xu¹ · Jun Ma¹ · Qiang Huang¹ · Ze-yu Huang¹ · Shao-yun Zhang¹ · Fu-xing Pei¹

Published online: 19 June 2017

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2017

Erratum to: Knee Surg Sports Traumatol Arthrosc DOI 10.1007/s00167-017-4506-x

Unfortunately, the online published article has error in Table 2. In the Metoclopramide, Number of patients requiring, The *P* value should be 0.018. The original article was corrected.

The correct table is given in the following page (Table 2).

Table 2 Comparison of the requirement of rescue treatment between the two groups

	Group dexta	Group placebo	<i>P</i> -value
Oxycodone			
Number of patients requiring [†]	5/54	18/54	0.004
Total dose [§]	250	870	0.001
Median dose [§]	0 (0–70)	0 (0–90)	0.005
Parecoxib			
Number of patients requiring [†]	0	4/54	n.s
Total dose [§]	0	160	0.042
Median dose [§]	0	0 (0–40)	n.s
Metoclopramide			
Number of patients requiring [†]	4/54	14/54	0.018
Total dose [§]	40	180	0.008
Median dose [§]	0 (0–10)	0 (0–20)	0.02
Ondansetron			
Number of patients requiring [†]	0	2/54	n.s
Total dose [§]	0	10	n.s
Median dose [§]	0	0 (0–5)	n.s

[§]Wilcoxon Mann–Whitney *U*, [†]Fisher exact tests

The online version of the original article can be found under doi: [10.1007/s00167-017-4506-x](https://doi.org/10.1007/s00167-017-4506-x).

✉ Fu-xing Pei
peifuxing@vip.163.com

¹ Department of Orthopaedics, West China Hospital, Sichuan University, 37# Wainan Guoxue Road, Chengdu 610041, People's Republic of China